



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:25 PM UTC

PDB ID : 5WRR / pdb_00005wrr
Title : Crystal structure of Fam20A
Authors : Zhu, Q.
Deposited on : 2016-12-03
Resolution : 2.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	438	12% 84% 13% ..
1	B	438	14% 87% 12% .
2	C	2	50% 50%
2	D	2	100%
2	E	2	50% 50%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7137 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pseudokinase FAM20A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	431	Total	C	N	O	S	0	0	0
			3494	2234	615	624	21			
1	B	438	Total	C	N	O	S	0	0	0
			3545	2267	623	634	21			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	332	LYS	ASN	engineered mutation	UNP Q96MK3
B	332	LYS	ASN	engineered mutation	UNP Q96MK3

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	B	1	14	8	1	5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pseudokinase FAM20A



• Molecule 1: Pseudokinase FAM20A



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  50%  50%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	156.85Å 156.85Å 143.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.14 – 2.51 39.14 – 2.51	Depositor EDS
% Data completeness (in resolution range)	99.8 (39.14-2.51) 92.8 (39.14-2.51)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 2.51Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.215 , 0.231 0.218 , 0.233	Depositor DCC
R_{free} test set	1996 reflections (2.84%)	wwPDB-VP
Wilson B-factor (Å ²)	50.4	Xtriage
Anisotropy	0.771	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.024 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7137	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.12	0/3578	0.35	0/4842
1	B	0.12	0/3633	0.34	0/4923
All	All	0.12	0/7211	0.35	0/9765

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3494	0	3475	36	0
1	B	3545	0	3531	38	0
2	C	28	0	25	1	0
2	D	28	0	25	1	0
2	E	28	0	25	0	0
3	B	14	0	13	1	0
All	All	7137	0	7094	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204:ALA:HB1	1:B:204:ALA:HB1	1.58	0.86
1:B:369:GLU:HG2	1:B:374:ASN:HD22	1.46	0.81
1:B:360:ILE:HG23	1:B:361:ARG:HG2	1.76	0.68
1:A:174:TYR:O	1:A:276:ARG:NH1	2.25	0.68
1:A:360:ILE:HG23	1:A:361:ARG:HG2	1.79	0.65
1:A:278:VAL:HG12	1:A:428:HIS:HB2	1.82	0.60
1:B:248:PRO:HG2	1:B:324:LYS:HB3	1.82	0.60
1:A:202:GLU:OE2	1:A:223:LYS:NZ	2.34	0.60
1:A:401:ILE:HG23	1:A:505:ILE:HD13	1.84	0.59
1:B:401:ILE:HG23	1:B:505:ILE:HD13	1.86	0.57
1:B:237:LYS:NZ	1:B:263:GLU:OE2	2.31	0.57
1:B:202:GLU:OE1	1:B:223:LYS:NZ	2.37	0.55
1:B:256:ASP:HB2	1:B:259:ARG:HB2	1.88	0.55
1:B:260:HIS:NE2	1:B:337:GLU:OE2	2.37	0.55
1:B:147:THR:O	1:B:149:LEU:N	2.38	0.54
1:B:141:ARG:HD3	1:B:148:SER:HA	1.91	0.53
1:B:278:VAL:HG12	1:B:428:HIS:HB2	1.90	0.52
1:B:141:ARG:HG2	1:B:146:LEU:O	2.09	0.52
1:A:213:GLN:HA	1:A:216:LYS:HE2	1.93	0.51
1:A:248:PRO:HD2	1:A:324:LYS:HB3	1.93	0.50
1:B:389:ASN:HB3	3:B:603:NAG:H82	1.93	0.50
1:B:160:SER:OG	1:B:187:ASP:OD2	2.30	0.50
1:A:367:GLY:HA3	1:A:368:LYS:HD2	1.94	0.49
1:B:367:GLY:HA3	1:B:368:LYS:HD2	1.95	0.48
1:A:318:LYS:HG2	1:A:322:MET:HE2	1.95	0.47
1:A:90:SER:OG	1:A:479:GLU:OE2	2.19	0.47
1:A:443:ILE:HD11	1:B:364:THR:HG21	1.97	0.47
1:B:207:GLY:HA3	1:B:208:ALA:HB3	1.96	0.46
1:A:367:GLY:HA2	1:A:368:LYS:HA	1.57	0.46
1:B:411:ARG:NH2	1:B:429:LEU:O	2.49	0.46
1:A:237:LYS:NZ	1:A:263:GLU:OE2	2.45	0.46
1:B:141:ARG:HE	1:B:147:THR:C	2.23	0.46
1:B:141:ARG:CZ	1:B:146:LEU:HG	2.45	0.46
1:A:399:MET:SD	1:A:411:ARG:HD3	2.56	0.46
1:A:403:ASP:OD2	1:A:411:ARG:HD2	2.16	0.46
1:B:399:MET:SD	1:B:411:ARG:HD3	2.57	0.45
1:B:395:ASN:HB3	1:B:426:LEU:HG	1.99	0.45
1:B:319:CYS:HB3	1:B:320:PRO:HD3	1.99	0.45
1:A:478:ARG:HB2	1:A:495:LEU:HD13	1.99	0.44
1:A:413:HIS:NE2	1:B:215:VAL:O	2.50	0.44
1:A:488:PRO:O	1:A:490:LEU:N	2.45	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:ASP:HB2	1:A:259:ARG:HB2	1.99	0.44
1:A:422:ASP:C	1:A:424:GLY:H	2.26	0.44
1:A:446:LEU:HD11	1:A:505:ILE:HG12	1.99	0.44
1:B:367:GLY:HA2	1:B:368:LYS:HA	1.58	0.43
1:A:411:ARG:NH2	1:A:429:LEU:O	2.49	0.43
1:B:270:ASP:CG	1:B:278:VAL:HG13	2.44	0.42
1:B:130:VAL:HG21	1:B:155:LEU:HB2	2.00	0.42
1:A:270:ASP:CG	1:A:278:VAL:HG13	2.44	0.42
1:B:519:GLN:CD	1:B:519:GLN:H	2.28	0.42
1:A:413:HIS:HE1	1:B:217:PRO:HD3	1.85	0.42
2:D:1:NAG:O3	2:D:2:NAG:H83	2.20	0.42
1:A:206:LEU:HA	1:A:207:GLY:HA3	1.81	0.41
1:A:211:CYS:O	1:A:215:VAL:HG23	2.20	0.41
1:A:364:THR:HG21	1:B:443:ILE:HD11	2.01	0.41
1:A:519:GLN:CD	1:A:519:GLN:H	2.27	0.41
1:B:374:ASN:OD1	1:B:376:LEU:HB2	2.19	0.41
1:A:287:ASN:HD22	2:C:1:NAG:H83	1.86	0.41
1:A:319:CYS:HB3	1:A:320:PRO:HD3	2.02	0.40
1:A:347:LEU:HD21	1:A:352:ARG:HE	1.86	0.40
1:A:355:VAL:HG22	1:B:249:VAL:HG22	2.03	0.40
1:B:156:ARG:HB3	1:B:158:GLU:OE1	2.20	0.40
1:B:403:ASP:OD2	1:B:411:ARG:HD2	2.21	0.40
1:A:160:SER:OG	1:A:187:ASP:OD2	2.35	0.40
1:B:141:ARG:NE	1:B:148:SER:HA	2.36	0.40
1:B:256:ASP:OD2	1:B:259:ARG:HD3	2.22	0.40
1:A:372:GLU:CD	1:A:372:GLU:H	2.29	0.40
1:A:446:LEU:HD23	1:A:446:LEU:HA	1.94	0.40
1:B:218:SER:HA	1:B:433:ARG:HD3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	425/438 (97%)	413 (97%)	12 (3%)	0	100	100
1	B	436/438 (100%)	422 (97%)	13 (3%)	1 (0%)	43	63
All	All	861/876 (98%)	835 (97%)	25 (3%)	1 (0%)	48	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	148	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	387/394 (98%)	376 (97%)	11 (3%)	38	66
1	B	394/394 (100%)	381 (97%)	13 (3%)	33	61
All	All	781/788 (99%)	757 (97%)	24 (3%)	35	62

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	153	LEU
1	A	160	SER
1	A	225	VAL
1	A	249	VAL
1	A	278	VAL
1	A	369	GLU
1	A	372	GLU
1	A	373	VAL
1	A	486	LEU
1	A	506	LEU
1	A	519	GLN
1	B	147	THR
1	B	149	LEU
1	B	153	LEU
1	B	158	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	160	SER
1	B	163	GLN
1	B	173	LEU
1	B	200	GLN
1	B	225	VAL
1	B	249	VAL
1	B	278	VAL
1	B	337	GLU
1	B	372	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	186	GLN
1	A	190	HIS
1	A	312	ASN
1	A	516	HIS
1	B	516	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2,1	14,14,15	0.34	0	17,19,21	0.50	0
2	NAG	C	2	2	14,14,15	0.34	0	17,19,21	0.58	0
2	NAG	D	1	2,1	14,14,15	0.21	0	17,19,21	0.52	0
2	NAG	D	2	2	14,14,15	0.41	0	17,19,21	0.44	0
2	NAG	E	1	2,1	14,14,15	0.31	0	17,19,21	0.78	0
2	NAG	E	2	2	14,14,15	1.09	1 (7%)	17,19,21	1.43	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	4/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	NAG	D	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	4/6/23/26	0/1/1/1
2	NAG	E	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	E	2	2	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	2	NAG	O5-C1	3.92	1.50	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2	NAG	C1-O5-C5	5.60	119.69	112.19

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	2	NAG	C4-C5-C6-O6
2	E	2	NAG	O5-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
2	D	1	NAG	O5-C5-C6-O6
2	D	1	NAG	C4-C5-C6-O6
2	C	1	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	1	NAG	O7-C7-N2-C2
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2
2	D	2	NAG	C4-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
2	E	1	NAG	C3-C2-N2-C7
2	C	1	NAG	O5-C5-C6-O6
2	E	1	NAG	C1-C2-N2-C7


There are no ring outliers.


3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	NAG	1	0
2	D	2	NAG	1	0
2	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	B	603	1	14,14,15	0.28	0	17,19,21	0.86	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	603	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	603	NAG	C1-O5-C5	3.20	116.47	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	603	NAG	O5-C5-C6-O6
3	B	603	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	603	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	431/438 (98%)	0.63	52 (12%) 8 7	45, 67, 121, 178	0
1	B	438/438 (100%)	0.82	63 (14%) 6 5	51, 71, 123, 185	0
All	All	869/876 (99%)	0.73	115 (13%) 7 6	45, 69, 122, 185	0

All (115) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	217	PRO	10.3
1	B	526	PRO	8.9
1	B	146	LEU	7.9
1	B	149	LEU	7.0
1	A	525	GLY	6.9
1	B	218	SER	6.6
1	B	144	MET	6.1
1	A	208	ALA	5.8
1	B	208	ALA	5.8
1	B	207	GLY	5.7
1	A	216	LYS	5.3
1	A	144	MET	5.0
1	B	147	THR	4.8
1	B	89	SER	4.8
1	B	216	LYS	4.8
1	A	207	GLY	4.7
1	A	151	PRO	4.7
1	B	368	LYS	4.7
1	A	423	ASP	4.6
1	A	111	GLY	4.5
1	B	221	HIS	4.4
1	B	111	GLY	4.4
1	B	106	GLU	4.2
1	B	367	GLY	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	231	PHE	4.0
1	B	143	GLN	4.0
1	A	218	SER	3.9
1	B	145	ASN	3.9
1	A	219	GLY	3.9
1	B	110	LEU	3.8
1	A	89	SER	3.8
1	B	206	LEU	3.7
1	A	368	LYS	3.7
1	B	153	LEU	3.6
1	B	213	GLN	3.5
1	B	136	ARG	3.5
1	A	519	GLN	3.5
1	A	150	ASP	3.5
1	A	110	LEU	3.5
1	B	148	SER	3.4
1	A	221	HIS	3.4
1	A	369	GLU	3.3
1	A	136	ARG	3.2
1	B	370	GLU	3.1
1	A	129	LYS	3.1
1	A	332	LYS	3.1
1	B	322	MET	3.1
1	B	256	ASP	3.0
1	A	128	ARG	3.0
1	B	437	ARG	3.0
1	A	319	CYS	3.0
1	A	220	VAL	3.0
1	B	141	ARG	2.9
1	B	214	ILE	2.9
1	A	153	LEU	2.9
1	A	213	GLN	2.8
1	A	112	ALA	2.8
1	B	222	LEU	2.8
1	B	330	CYS	2.8
1	B	142	GLU	2.7
1	A	139	MET	2.7
1	A	132	ARG	2.7
1	B	204	ALA	2.7
1	B	244	ASP	2.6
1	A	367	GLY	2.6
1	A	330	CYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	376	LEU	2.6
1	A	439	SER	2.6
1	A	512	CYS	2.6
1	B	120	GLN	2.6
1	A	322	MET	2.6
1	B	107	PRO	2.6
1	B	124	ARG	2.5
1	A	156	ARG	2.5
1	B	422	ASP	2.5
1	B	332	LYS	2.5
1	B	215	VAL	2.5
1	B	129	LYS	2.5
1	A	413	HIS	2.5
1	A	135	ARG	2.5
1	A	120	GLN	2.4
1	B	140	TYR	2.4
1	B	390	SER	2.4
1	A	421	GLY	2.4
1	A	154	GLN	2.4
1	A	143	GLN	2.3
1	B	200	GLN	2.3
1	B	423	ASP	2.3
1	B	139	MET	2.3
1	A	206	LEU	2.3
1	B	156	ARG	2.3
1	A	209	CYS	2.3
1	B	138	LYS	2.2
1	A	152	PRO	2.2
1	A	378	CYS	2.2
1	B	201	ASP	2.2
1	B	112	ALA	2.2
1	B	418	THR	2.2
1	B	361	ARG	2.2
1	A	204	ALA	2.1
1	B	209	CYS	2.1
1	A	113	GLU	2.1
1	A	214	ILE	2.1
1	A	390	SER	2.1
1	B	150	ASP	2.1
1	B	378	CYS	2.1
1	A	124	ARG	2.1
1	B	151	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	141	ARG	2.0
1	B	310	ALA	2.0
1	B	512	CYS	2.0
1	A	189	ARG	2.0
1	A	507	ARG	2.0
1	B	412	HIS	2.0
1	B	413	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	D	2	14/15	0.31	0.28	127,148,154,155	0
2	NAG	D	1	14/15	0.70	0.21	101,116,132,139	0
2	NAG	E	2	14/15	0.77	0.18	116,131,139,141	0
2	NAG	E	1	14/15	0.85	0.15	91,97,107,116	0
2	NAG	C	2	14/15	0.86	0.14	79,101,117,124	0
2	NAG	C	1	14/15	0.91	0.12	71,81,95,98	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	B	603	14/15	0.60	0.26	119,130,141,144	0

6.5 Other polymers [i](#)

There are no such residues in this entry.