



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 10:04 AM UTC

PDB ID : 5WRZ / pdb_00005wrz
Title : Structure of human PARP1 catalytic domain bound to a phthalazinone inhibitor
Authors : Cao, R.; Wang, Y.L.; Zhou, J.; Huang, N.; Xu, B.L.
Deposited on : 2016-12-04
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	9% 90% 9%
1	B	352	13% 87% 12%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 5584 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Poly [ADP-ribose] polymerase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	351	Total	C	N	O	S	0	3	0
			2773	1763	470	527	13			
1	B	351	Total	C	N	O	S	0	3	0
			2773	1763	470	527	13			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	660	HIS	-	expression tag	UNP P09874
A	661	MET	-	expression tag	UNP P09874
A	762	ALA	VAL	engineered mutation	UNP P09874
B	660	HIS	-	expression tag	UNP P09874
B	661	MET	-	expression tag	UNP P09874
B	762	ALA	VAL	engineered mutation	UNP P09874

- Molecule 2 is 7-fluoranyl-4-[[[(3R)-pyrrolidin-3-yl]methoxy]-2H-phthalazin-1-one (CCD ID: 7U3) (formula: C₁₃H₁₄FN₃O₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			19	13	1	3	2		
2	B	1	Total	C	F	N	O	0	0
			19	13	1	3	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly [ADP-ribose] polymerase 1



- Molecule 1: Poly [ADP-ribose] polymerase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.77Å 92.35Å 163.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	81.60 – 2.20 81.60 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (81.60-2.20) 99.7 (81.60-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.59 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, R_{free}	0.230 , 0.280 (Not available) , 0.298	Depositor DCC
R_{free} test set	1855 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	19.2	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 24.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5584	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 7U3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.08	2/2837 (0.1%)	1.09	3/3827 (0.1%)
1	B	1.05	1/2837 (0.0%)	1.13	12/3827 (0.3%)
All	All	1.07	3/5674 (0.1%)	1.11	15/7654 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	914	ASP	CA-C	7.24	1.61	1.52
1	A	914	ASP	N-CA	6.59	1.55	1.46
1	B	941	LEU	C-O	-5.18	1.18	1.24

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	749	PRO	CA-C-N	8.23	130.12	119.84
1	B	749	PRO	C-N-CA	8.23	130.12	119.84
1	A	780	GLY	N-CA-C	-7.21	100.22	111.98
1	B	722	GLN	N-CA-C	7.01	118.94	107.23
1	B	744	PHE	N-CA-C	-6.77	103.70	112.23
1	B	751	LEU	N-CA-C	6.50	120.42	109.76

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	747	LYS	N-CA-C	6.49	124.63	110.80
1	B	727	SER	N-CA-C	-6.36	104.27	111.14
1	B	748	LYS	CB-CG-CD	6.32	125.84	111.30
1	B	750	PRO	N-CA-C	6.03	124.90	112.47
1	B	749	PRO	N-CA-C	5.96	117.97	110.70
1	A	785	SER	N-CA-C	5.76	118.33	111.71
1	B	679	VAL	N-CA-C	5.76	116.49	110.62
1	B	748	LYS	CA-CB-CG	5.48	125.05	114.10
1	B	950	GLY	N-CA-C	-5.16	100.96	113.18

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	746	MET	Peptide
1	A	747	LYS	Peptide
1	B	743	ASP	Peptide
1	B	750	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2773	0	2806	9	2
1	B	2773	0	2806	14	2
2	A	19	0	0	0	0
2	B	19	0	0	0	0
All	All	5584	0	5612	22	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:675:MET:HE1	1:B:1004:LYS:HD2	1.68	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:934:HIS:HD1	1:B:983:SER:HG	1.42	0.68
1:A:717:GLN:OE1	1:A:887:THR:HB	2.02	0.60
1:B:770:ASP:OD1	1:B:878:ARG:NH2	2.37	0.57
1:A:841:ARG:HD2	1:A:873:LEU:O	2.06	0.55
1:B:841:ARG:HD2	1:B:873:LEU:O	2.07	0.54
1:A:886:VAL:HG11	1:B:934:HIS:CD2	2.42	0.53
1:B:919:ILE:HG21	1:B:1005:LEU:HD11	1.91	0.52
1:B:729:ILE:HG21	1:B:753:ASN:HA	1.93	0.51
1:B:726:ASP:HA	1:B:729:ILE:HB	1.98	0.46
1:A:770:ASP:OD1	1:A:878:ARG:NH2	2.48	0.46
1:B:747:LYS:O	1:B:748:LYS:HB2	2.16	0.46
1:A:821:THR:HB	1:A:900:MET:HA	1.98	0.45
1:B:674:LYS:HG3	1:B:790:ILE:HD11	2.00	0.44
1:B:723:GLY:O	1:B:725:SER:N	2.53	0.42
1:B:892:GLY:HA3	1:B:938:ILE:O	2.19	0.42
1:A:684:LYS:HA	1:A:687:VAL:HG22	2.02	0.41
1:A:934:HIS:ND1	1:A:983:SER:OG	2.51	0.41
1:B:744:PHE:CD2	1:B:749:PRO:CB	3.03	0.41
1:A:841:ARG:CD	1:A:873:LEU:O	2.69	0.41
1:A:872:ILE:HG21	1:A:920:LEU:HD11	2.04	0.40
1:B:717:GLN:OE1	1:B:887:THR:HB	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:715:GLU:OE1	1:B:705:GLN:NE2[1_455]	2.03	0.17
1:A:911:SER:O	1:B:748:LYS:CG[4_445]	2.11	0.09

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/352 (99%)	340 (97%)	5 (1%)	5 (1%)	9	7
1	B	350/352 (99%)	335 (96%)	10 (3%)	5 (1%)	9	7
All	All	700/704 (99%)	675 (96%)	15 (2%)	10 (1%)	9	7

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	914	ASP
1	B	749	PRO
1	B	748	LYS
1	B	750	PRO
1	A	662	LYS
1	A	911	SER
1	B	721	SER
1	A	748	LYS
1	A	663	SER
1	B	663	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/309 (101%)	302 (97%)	9 (3%)	37	51
1	B	311/309 (101%)	299 (96%)	12 (4%)	28	39
All	All	622/618 (101%)	601 (97%)	21 (3%)	32	44

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	667	LYS
1	A	700	LYS
1	A	778	LEU
1	A	914	ASP
1	A	939	SER
1	A	940	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	962	ILE
1	A	963	SER
1	A	1006	LYS
1	B	667	LYS
1	B	700	LYS
1	B	704	ARG
1	B	713	LEU
1	B	730	LEU
1	B	748	LYS
1	B	750	PRO
1	B	769	LEU
1	B	865	ARG
1	B	912	GLN
1	B	940	LYS
1	B	1011	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	707	GLN
1	A	722	GLN
1	A	855	HIS
1	A	998	ASN
1	B	660	HIS
1	B	759	GLN
1	B	868	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	7U3	B	1101	-	20,21,21	2.71	6 (30%)	24,29,29	2.18	5 (20%)
2	7U3	A	1101	-	20,21,21	2.25	8 (40%)	24,29,29	1.95	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7U3	B	1101	-	-	0/5/12/12	0/3/3/3
2	7U3	A	1101	-	-	1/5/12/12	0/3/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1101	7U3	FAB-CAN	-7.43	1.18	1.36
2	B	1101	7U3	CAQ-CAO	-5.86	1.36	1.46
2	B	1101	7U3	CAG-CAF	-4.59	1.46	1.53
2	A	1101	7U3	NAL-NAJ	-4.36	1.26	1.38
2	A	1101	7U3	CAQ-CAO	-4.07	1.39	1.46
2	A	1101	7U3	CAR-CAP	-3.97	1.40	1.47
2	A	1101	7U3	CAP-NAL	3.64	1.41	1.35
2	B	1101	7U3	CAR-CAP	-3.33	1.41	1.47
2	A	1101	7U3	FAB-CAN	-3.25	1.28	1.36
2	A	1101	7U3	CAG-CAF	-3.02	1.48	1.53
2	B	1101	7U3	NAL-NAJ	-2.81	1.30	1.38
2	A	1101	7U3	OAM-CAO	2.28	1.38	1.34
2	A	1101	7U3	CAO-NAJ	2.24	1.34	1.30
2	B	1101	7U3	CAC-CAN	2.12	1.41	1.37

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1101	7U3	CAP-NAL-NAJ	-6.84	120.35	127.98
2	A	1101	7U3	CAP-NAL-NAJ	-6.39	120.85	127.98
2	A	1101	7U3	CAO-NAJ-NAL	4.79	124.31	117.22
2	B	1101	7U3	CAO-NAJ-NAL	4.64	124.10	117.22
2	B	1101	7U3	FAB-CAN-CAC	3.48	124.12	118.55
2	B	1101	7U3	FAB-CAN-CAE	-2.94	114.10	118.28
2	B	1101	7U3	OAM-CAI-CAS	-2.39	103.11	108.75
2	A	1101	7U3	CAR-CAP-NAL	2.39	117.79	115.04
2	A	1101	7U3	CAI-OAM-CAO	2.33	121.83	115.37

There are no chirality outliers.


All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1101	7U3	OAM-CAI-CAS-CAH

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	351/352 (99%)	0.38	33 (9%) 14 11	6, 17, 41, 84	3 (0%)
1	B	351/352 (99%)	0.84	44 (12%) 8 6	7, 20, 49, 169	3 (0%)
All	All	702/704 (99%)	0.61	77 (10%) 10 8	6, 19, 44, 169	6 (0%)

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1011	THR	11.0
1	B	745	GLY	10.9
1	B	749	PRO	10.2
1	B	726	ASP	9.9
1	B	747	LYS	9.6
1	B	750	PRO	9.0
1	B	725	SER	8.7
1	B	746	MET	8.3
1	A	661	MET	7.9
1	B	661	MET	7.9
1	B	744	PHE	6.7
1	A	913	GLY	6.5
1	B	723	GLY	6.4
1	A	915	PRO	6.3
1	A	747	LYS	5.5
1	A	660	HIS	5.4
1	B	980	ASN	5.2
1	B	981	ASP	5.2
1	B	748	LYS	4.9
1	B	662	LYS	4.8
1	B	660	HIS	4.3
1	B	731	ASP	4.3
1	B	705	GLN	4.3
1	A	914	ASP	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	912	GLN	4.1
1	B	752	LEU	4.1
1	B	722	GLN	4.1
1	B	727	SER	4.0
1	A	662	LYS	3.9
1	A	728	GLN	3.9
1	B	962	ILE	3.9
1	A	780	GLY	3.6
1	A	725	SER	3.4
1	B	730	LEU	3.3
1	B	785	SER	3.3
1	B	694	GLN	3.3
1	B	827	ASN	3.2
1	B	795	GLU	3.2
1	A	1011	THR	3.1
1	B	696	MET	3.0
1	A	743	ASP	2.9
1	A	721	SER	2.9
1	A	722	GLN	2.9
1	B	943	LYS	2.8
1	B	966	GLY	2.8
1	B	751	LEU	2.8
1	B	979	VAL	2.8
1	A	727	SER	2.7
1	A	939	SER	2.7
1	A	795	GLU	2.6
1	B	743	ASP	2.6
1	A	806	ARG	2.6
1	B	783	ASP	2.6
1	A	744	PHE	2.6
1	B	963	SER	2.6
1	A	726	ASP	2.5
1	A	1010	LYS	2.5
1	A	735	ARG	2.4
1	B	687	VAL	2.4
1	B	688	GLU	2.4
1	A	689	TYR	2.3
1	B	735	ARG	2.3
1	B	853	GLN	2.2
1	A	980	ASN	2.2
1	B	757	SER	2.2
1	A	719	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	695	LYS	2.1
1	B	807	ASP	2.1
1	B	914	ASP	2.1
1	A	759	GLN	2.1
1	A	695	LYS	2.1
1	A	718	GLN	2.1
1	B	965	ASP	2.1
1	A	745	GLY	2.1
1	A	723	GLY	2.0
1	A	691	ILE	2.0
1	A	940	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	7U3	A	1101	19/19	0.94	0.09	9,11,27,27	0
2	7U3	B	1101	19/19	0.95	0.07	10,11,23,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 7U3 A 1101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.