



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:28 PM UTC

PDB ID : 4WSJ / pdb_00004wsj
Title : Crystal structure of a bacterial fucodiase in complex with 1-((1R,2R,3R,4R,5R,6R)-2,3,4-trihydroxy-5-methyl-7-azabicyclo[4.1.0]heptan-7-yl)ethan-1-one
Authors : Davies, G.J.; Wright, D.W.
Deposited on : 2014-10-28
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1141 (1.64-1.64)
Clashscore	190562	1171 (1.64-1.64)
Ramachandran outliers	187476	1151 (1.64-1.64)
Sidechain outliers	187428	1150 (1.64-1.64)
RSRZ outliers	180081	1141 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	445	0% 94% 5%
1	B	445	2% 92% 7%
1	C	445	0% 94% • •
1	D	445	3% 92% 7%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 15772 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-L-fucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	439	Total	C	N	O	S	0	0	0
			3495	2248	579	652	16			
1	B	445	Total	C	N	O	S	0	1	0
			3569	2296	598	659	16			
1	C	437	Total	C	N	O	S	0	1	0
			3498	2253	579	650	16			
1	D	440	Total	C	N	O	S	0	0	0
			3511	2261	581	653	16			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is N-[(1S,2R,3R,4S,5R)-3,4,5-trihydroxy-2-methylcyclohexyl]acetamide (CCD ID: 3U3) (formula: C₉H₁₇NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	9	1	4		
3	B	1	Total	C	N	O	0	0
			14	9	1	4		
3	C	1	Total	C	N	O	0	0
			14	9	1	4		
3	D	1	Total	C	N	O	0	0
			14	9	1	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	426	Total	O	0	0
			426	426		
4	B	458	Total	O	0	0
			458	458		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	409	Total 409	O 409	0	0
4	D	325	Total 325	O 325	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-L-fucosidase




- Molecule 1: Alpha-L-fucosidase



- Molecule 1: Alpha-L-fucosidase



- Molecule 1: Alpha-L-fucosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.48Å 186.98Å 98.19Å 90.00° 94.17° 90.00°	Depositor
Resolution (Å)	93.49 – 1.64 93.49 – 1.64	Depositor EDS
% Data completeness (in resolution range)	98.0 (93.49-1.64) 98.0 (93.49-1.64)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 1.64Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.164 , 0.194 0.175 , 0.203	Depositor DCC
R_{free} test set	11928 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.202	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	15772	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 3U3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.21	7/3602 (0.2%)	1.04	2/4906 (0.0%)
1	B	1.28	9/3680 (0.2%)	1.10	2/5005 (0.0%)
1	C	1.25	4/3608 (0.1%)	1.04	2/4911 (0.0%)
1	D	1.19	7/3618 (0.2%)	1.01	6/4925 (0.1%)
All	All	1.23	27/14508 (0.2%)	1.05	12/19747 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	229	ASP	CG-OD2	8.83	1.42	1.25
1	B	229	ASP	CG-OD2	8.29	1.41	1.25
1	B	308	CYS	C-N	-7.37	1.24	1.33
1	A	309	MET	SD-CE	-6.98	1.62	1.79
1	B	272	HIS	N-CA	6.01	1.53	1.46
1	D	229	ASP	CG-OD2	5.80	1.36	1.25
1	B	83	GLY	C-O	5.74	1.30	1.24
1	D	223	VAL	CA-CB	5.70	1.60	1.54
1	A	229	ASP	CG-OD2	5.70	1.36	1.25
1	D	86	ALA	N-CA	5.58	1.53	1.46
1	A	104	MET	N-CA	5.51	1.53	1.46
1	D	413	PRO	C-O	5.46	1.29	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	219	ARG	N-CA	-5.46	1.39	1.46
1	C	134	LYS	N-CA	-5.39	1.40	1.46
1	B	76	GLU	N-CA	5.34	1.52	1.46
1	B	285	SER	CA-CB	5.33	1.60	1.53
1	B	137	GLU	N-CA	5.31	1.52	1.46
1	C	38	LEU	N-CA	5.30	1.52	1.46
1	B	434	LEU	C-O	5.20	1.30	1.24
1	A	272	HIS	N-CA	5.18	1.52	1.46
1	C	163	GLU	CA-C	-5.16	1.46	1.52
1	A	361	PRO	CA-CB	5.14	1.61	1.53
1	A	406	TYR	N-CA	-5.11	1.39	1.46
1	D	183	ASP	N-CA	-5.08	1.40	1.46
1	A	407	MET	N-CA	5.02	1.52	1.46
1	D	181	VAL	CA-CB	-5.01	1.48	1.54
1	B	53	MET	N-CA	5.00	1.52	1.46

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	288	GLU	CA-CB-CG	6.82	127.73	114.10
1	B	478	ILE	CB-CA-C	5.82	120.83	111.29
1	D	219	ARG	CG-CD-NE	-5.75	99.35	112.00
1	D	378	GLY	N-CA-C	5.53	121.07	113.99
1	B	376	ARG	CG-CD-NE	-5.43	100.06	112.00
1	D	260	ASN	N-CA-C	5.33	117.75	110.55
1	A	309	MET	CG-SD-CE	-5.21	89.45	100.90
1	C	429	GLU	CA-CB-CG	5.14	124.38	114.10
1	D	264	ARG	N-CA-C	5.14	118.00	109.46
1	D	219	ARG	N-CA-CB	-5.11	102.43	110.30
1	C	260	ASN	N-CA-C	5.08	117.23	110.53
1	A	313	GLU	N-CA-C	5.02	116.44	111.07

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	128	TYR	Sidechain
1	B	128	TYR	Sidechain
1	B	477	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3495	0	3265	6	0
1	B	3569	0	3371	15	0
1	C	3498	0	3278	9	0
1	D	3511	0	3283	11	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
2	C	5	0	0	0	0
3	A	14	0	16	0	0
3	B	14	0	16	0	0
3	C	14	0	16	0	0
3	D	14	0	16	0	0
4	A	426	0	0	0	0
4	B	458	0	0	2	0
4	C	409	0	0	2	0
4	D	325	0	0	2	0
All	All	15772	0	13261	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:444:GLU:HG3	4:D:703:HOH:O	1.84	0.75
1:C:123:GLU:HG2	4:C:830:HOH:O	1.92	0.69
1:B:477:SER:O	1:B:478:ILE:HB	1.98	0.64
1:B:158:ARG:HD2	1:B:163:GLU:OE2	1.98	0.64
1:C:158:ARG:HD2	1:C:163:GLU:OE2	2.00	0.61
1:D:158:ARG:HD2	1:D:163:GLU:OE2	2.03	0.59
1:B:379:LYS:HE2	1:B:406:TYR:OH	2.02	0.58
1:B:176:HIS:CG	1:B:225:ASP:HB3	2.41	0.56
1:C:176:HIS:CG	1:C:225:ASP:HB3	2.43	0.54
1:D:176:HIS:CG	1:D:225:ASP:HB3	2.43	0.54
1:B:320:LYS:HE2	4:B:969:HOH:O	2.07	0.53
1:A:176:HIS:CG	1:A:225:ASP:HB3	2.44	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:188:ASP:OD2	1:D:204:ARG:NH1	2.43	0.52
1:C:132:THR:HA	1:C:178:TYR:HB3	1.92	0.51
1:B:429:GLU:OE2	1:B:470:ARG:HD2	2.12	0.50
1:A:178:TYR:C	1:A:178:TYR:CD1	2.92	0.48
1:D:116:LYS:HE2	4:D:692:HOH:O	2.13	0.47
1:B:178:TYR:C	1:B:178:TYR:CD1	2.92	0.47
1:C:291:LEU:HD21	1:C:347:MET:HE1	1.95	0.47
1:A:132:THR:HA	1:A:178:TYR:HB3	1.97	0.47
1:D:178:TYR:CD1	1:D:178:TYR:C	2.92	0.47
1:A:291:LEU:HD21	1:A:347:MET:HE1	1.97	0.47
1:D:291:LEU:HD21	1:D:347:MET:HE1	1.96	0.46
1:B:475:THR:HG22	1:B:476:LYS:N	2.31	0.46
1:C:145:LYS:NZ	1:D:252:GLU:OE1	2.49	0.46
1:A:305:TRP:CE2	1:A:345:GLY:HA3	2.51	0.46
1:B:291:LEU:HD21	1:B:347:MET:HE1	1.99	0.45
1:B:132:THR:HA	1:B:178:TYR:HB3	2.00	0.44
1:C:158:ARG:NH2	4:C:602:HOH:O	2.44	0.44
1:A:311:ILE:HB	1:A:312:PRO:HD3	2.01	0.42
1:D:132:THR:HA	1:D:178:TYR:HB3	2.01	0.42
1:B:449:GLU:CB	4:B:1037:HOH:O	2.68	0.42
1:C:178:TYR:C	1:C:178:TYR:CD1	2.98	0.42
1:D:309:MET:HB3	1:D:309:MET:HE3	1.66	0.41
1:B:311:ILE:HB	1:B:312:PRO:HD3	2.02	0.41
1:B:130:LYS:HE2	1:B:227:TRP:CD1	2.56	0.41
1:B:399:ARG:NH1	1:B:478:ILE:HG23	2.35	0.41
1:B:476:LYS:O	1:B:477:SER:CB	2.69	0.41
1:C:311:ILE:HB	1:C:312:PRO:HD3	2.03	0.40
1:D:311:ILE:HB	1:D:312:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	437/445 (98%)	424 (97%)	13 (3%)	0	100	100
1	B	444/445 (100%)	435 (98%)	7 (2%)	2 (0%)	24	9
1	C	436/445 (98%)	425 (98%)	11 (2%)	0	100	100
1	D	437/445 (98%)	429 (98%)	8 (2%)	0	100	100
All	All	1754/1780 (98%)	1713 (98%)	39 (2%)	2 (0%)	48	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	477	SER
1	B	476	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	352/379 (93%)	349 (99%)	3 (1%)	70	54
1	B	361/379 (95%)	359 (99%)	2 (1%)	78	66
1	C	352/379 (93%)	349 (99%)	3 (1%)	70	54
1	D	353/379 (93%)	349 (99%)	4 (1%)	65	44
All	All	1418/1516 (94%)	1406 (99%)	12 (1%)	73	58

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	PHE
1	A	110	THR
1	A	267	ASP
1	B	372	LYS
1	B	478	ILE
1	C	64	PHE
1	C	123	GLU
1	C	372	LYS
1	D	110	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	123	GLU
1	D	232	TRP
1	D	288	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	169	ASN
1	A	314	ASN
1	B	314	ASN
1	B	411	ASN
1	B	448	ASN
1	C	314	ASN
1	C	375	ASN
1	C	411	ASN
1	C	448	ASN
1	D	314	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	502	-	4,4,4	0.50	0	6,6,6	0.70	0
2	SO4	B	502	-	4,4,4	0.40	0	6,6,6	0.55	0
2	SO4	C	501	-	4,4,4	0.48	0	6,6,6	0.94	0
3	3U3	A	503	1	14,14,14	2.67	6 (42%)	18,20,20	2.02	6 (33%)
3	3U3	D	501	1	14,14,14	2.18	5 (35%)	18,20,20	1.89	5 (27%)
2	SO4	A	501	-	4,4,4	0.31	0	6,6,6	0.83	0
3	3U3	B	503	1	14,14,14	2.40	8 (57%)	18,20,20	1.81	6 (33%)
2	SO4	B	501	-	4,4,4	0.51	0	6,6,6	0.94	0
3	3U3	C	502	1	14,14,14	1.58	2 (14%)	18,20,20	1.86	6 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	3U3	D	501	1	-	0/4/24/24	0/1/1/1
3	3U3	B	503	1	-	0/4/24/24	0/1/1/1
3	3U3	A	503	1	-	0/4/24/24	0/1/1/1
3	3U3	C	502	1	-	0/4/24/24	0/1/1/1

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	501	3U3	CAE-CAD	-6.59	1.42	1.52
3	A	503	3U3	CAE-CAD	-6.22	1.43	1.52
3	B	503	3U3	CAE-CAD	-5.23	1.44	1.52
3	C	502	3U3	CAF-CAE	4.62	1.61	1.52
3	A	503	3U3	OAH-CAE	4.39	1.52	1.43
3	A	503	3U3	CAF-CAA	3.36	1.58	1.53
3	B	503	3U3	CAF-CAA	-3.31	1.48	1.53
3	C	502	3U3	OAH-CAE	3.11	1.49	1.43
3	B	503	3U3	OAH-CAE	3.04	1.49	1.43
3	A	503	3U3	CAA-NAG	2.80	1.50	1.46
3	B	503	3U3	OAI-CAD	2.52	1.49	1.43
3	B	503	3U3	CAN-CAL	2.50	1.55	1.50
3	D	501	3U3	OAH-CAE	2.49	1.48	1.43
3	B	503	3U3	CAD-CAC	-2.41	1.46	1.52
3	A	503	3U3	OAJ-CAC	2.40	1.48	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	501	3U3	CAA-NAG	2.36	1.50	1.46
3	A	503	3U3	CAF-CAE	-2.35	1.48	1.52
3	B	503	3U3	CAB-CAC	2.31	1.57	1.53
3	B	503	3U3	CAF-CAE	2.17	1.56	1.52
3	D	501	3U3	CAB-CAA	-2.16	1.49	1.54
3	D	501	3U3	CAF-CAE	2.12	1.56	1.52

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	502	3U3	CAF-CAA-CAB	5.10	119.69	114.49
3	A	503	3U3	CAF-CAE-CAD	4.42	117.03	110.67
3	B	503	3U3	CAA-NAG-CAL	-3.96	117.60	122.90
3	D	501	3U3	CAF-CAE-CAD	3.88	116.26	110.67
3	A	503	3U3	CAA-NAG-CAL	-3.68	117.97	122.90
3	A	503	3U3	CAB-CAA-NAG	-3.07	105.50	110.42
3	D	501	3U3	CAK-CAB-CAA	-2.92	107.57	111.47
3	B	503	3U3	CAK-CAB-CAA	-2.91	107.59	111.47
3	B	503	3U3	CAB-CAA-NAG	-2.84	105.86	110.42
3	A	503	3U3	CAF-CAA-CAB	2.80	117.34	114.49
3	A	503	3U3	OAM-CAL-NAG	2.68	126.71	121.98
3	B	503	3U3	CAF-CAE-CAD	2.68	114.52	110.67
3	C	502	3U3	CAF-CAA-NAG	-2.63	106.76	110.78
3	D	501	3U3	CAA-NAG-CAL	-2.60	119.42	122.90
3	D	501	3U3	OAM-CAL-NAG	2.49	126.38	121.98
3	C	502	3U3	CAE-CAF-CAA	2.45	117.42	112.59
3	D	501	3U3	CAF-CAA-NAG	-2.44	107.06	110.78
3	B	503	3U3	OAI-CAD-CAE	-2.31	105.35	110.05
3	C	502	3U3	CAB-CAA-NAG	-2.12	107.02	110.42
3	C	502	3U3	OAI-CAD-CAE	-2.11	105.76	110.05
3	A	503	3U3	CAE-CAF-CAA	2.05	116.63	112.59
3	C	502	3U3	CAK-CAB-CAA	-2.03	108.76	111.47
3	B	503	3U3	CAE-CAF-CAA	2.01	116.55	112.59

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/445 (98%)	-0.07	4 (0%) 81 85	13, 23, 37, 56	1 (0%)
1	B	445/445 (100%)	-0.29	7 (1%) 70 75	11, 19, 34, 55	2 (0%)
1	C	437/445 (98%)	-0.07	6 (1%) 73 79	16, 24, 37, 52	2 (0%)
1	D	440/445 (98%)	0.41	15 (3%) 48 52	17, 28, 47, 64	1 (0%)
All	All	1761/1780 (98%)	-0.01	32 (1%) 67 73	11, 24, 40, 64	6 (0%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	478	ILE	4.2
1	D	200	ILE	4.2
1	D	98	ASP	3.3
1	D	37	PRO	3.0
1	A	43	THR	3.0
1	D	479	TYR	3.0
1	D	36	ILE	3.0
1	B	477	SER	2.9
1	D	197	GLU	2.9
1	A	36	ILE	2.9
1	C	37	PRO	2.9
1	D	110	THR	2.7
1	D	156	TYR	2.7
1	D	473	LYS	2.6
1	D	77	TRP	2.6
1	C	36	ILE	2.5
1	D	35	GLU	2.4
1	D	191	TYR	2.3
1	D	193	ILE	2.3
1	A	37	PRO	2.3
1	B	479	TYR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	446	THR	2.2
1	D	194	LYS	2.2
1	B	309	MET	2.2
1	D	95	VAL	2.2
1	A	98	ASP	2.2
1	B	476	LYS	2.1
1	B	475	THR	2.1
1	C	43	THR	2.1
1	B	474	GLY	2.1
1	C	35	GLU	2.0
1	C	98	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	502	5/5	0.96	0.07	29,30,33,39	0
2	SO4	B	501	5/5	0.96	0.07	42,43,50,52	0
3	3U3	D	501	14/14	0.96	0.07	20,23,34,34	0
3	3U3	B	503	14/14	0.97	0.06	14,15,21,24	0
2	SO4	C	501	5/5	0.98	0.07	24,25,30,31	0
3	3U3	A	503	14/14	0.98	0.05	17,20,23,24	0
2	SO4	A	501	5/5	0.98	0.06	26,27,36,36	0
3	3U3	C	502	14/14	0.98	0.04	15,17,21,22	0
2	SO4	B	502	5/5	0.98	0.06	23,23,29,31	0

6.5 Other polymers ⓘ

There are no such residues in this entry.