



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 01:08 AM UTC

PDB ID : 5WS1 / pdb_00005ws1
Title : Structure of human PARP1 catalytic domain bound to a benzoimidazole inhibitor
Authors : Cao, R.; Wang, Y.L.; Zhou, J.; Huang, N.; Xu, B.L.
Deposited on : 2016-12-04
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	
1	B	352	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 5586 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Poly [ADP-ribose] polymerase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	351	Total	C	N	O	S	0	3	0
			2773	1763	470	527	13			
1	B	351	Total	C	N	O	S	0	3	0
			2773	1763	470	527	13			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	660	HIS	-	expression tag	UNP P09874
A	661	MET	-	expression tag	UNP P09874
A	762	ALA	VAL	engineered mutation	UNP P09874
B	660	HIS	-	expression tag	UNP P09874
B	661	MET	-	expression tag	UNP P09874
B	762	ALA	VAL	engineered mutation	UNP P09874

- Molecule 2 is 2-[(3R)-3-azanylpyrrolidin-1-yl]carbonyl-1H-benzimidazole-4-carboxamide (CCD ID: 7U9) (formula: C₁₃H₁₅N₅O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			20	13	5	2		
2	B	1	Total	C	N	O	0	0
			20	13	5	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly [ADP-ribose] polymerase 1



- Molecule 1: Poly [ADP-ribose] polymerase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.71Å 91.87Å 162.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	81.11 – 1.90 81.11 – 1.90	Depositor EDS
% Data completeness (in resolution range)	95.9 (81.11-1.90) 95.9 (81.11-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.13 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, R_{free}	0.233 , 0.273 0.240 , 0.273	Depositor DCC
R_{free} test set	2751 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.282	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 27.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5586	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 7U9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.16	3/2837 (0.1%)	1.21	13/3827 (0.3%)
1	B	1.18	7/2837 (0.2%)	1.18	12/3827 (0.3%)
All	All	1.17	10/5674 (0.2%)	1.20	25/7654 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	1
All	All	0	4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	749	PRO	CA-C	10.33	1.61	1.52
1	A	781	GLY	N-CA	9.31	1.58	1.45
1	B	748	LYS	CA-C	9.04	1.64	1.52
1	B	748	LYS	CA-CB	8.20	1.66	1.53
1	B	748	LYS	N-CA	7.67	1.56	1.46
1	B	780	GLY	C-O	6.86	1.30	1.23
1	A	841	ARG	CD-NE	-6.61	1.36	1.46
1	B	748	LYS	CB-CG	6.24	1.71	1.52
1	A	780	GLY	C-O	-5.51	1.18	1.23
1	B	841	ARG	CD-NE	-5.47	1.38	1.46

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	747	LYS	N-CA-C	14.00	129.59	108.31
1	B	744	PHE	N-CA-C	-12.07	96.67	111.40
1	A	780	GLY	O-C-N	-10.41	113.86	122.81
1	B	748	LYS	CB-CG-CD	8.52	130.90	111.30
1	A	747	LYS	CB-CA-C	-7.93	107.42	116.54
1	B	748	LYS	CA-CB-CG	7.41	128.92	114.10
1	A	780	GLY	CA-C-N	-6.72	108.23	121.41
1	A	780	GLY	C-N-CA	-6.72	108.23	121.41
1	B	841	ARG	NE-CZ-NH2	-6.62	113.24	119.20
1	A	841	ARG	NE-CZ-NH2	-6.50	113.35	119.20
1	B	749	PRO	N-CA-CB	-6.05	97.21	103.08
1	B	749	PRO	CA-C-N	6.04	127.39	119.84
1	B	749	PRO	C-N-CA	6.04	127.39	119.84
1	A	781	GLY	N-CA-C	5.98	127.34	113.18
1	A	890	MET	N-CA-C	5.90	118.46	111.33
1	A	841	ARG	NE-CZ-NH1	5.66	127.16	121.50
1	B	746	MET	N-CA-C	-5.57	105.29	111.36
1	B	750	PRO	CA-N-CD	-5.52	104.27	112.00
1	A	841	ARG	CD-NE-CZ	5.50	132.11	124.40
1	A	693	LEU	N-CA-C	5.43	119.00	112.38
1	A	796	LYS	CD-CE-NZ	-5.38	94.67	111.90
1	B	748	LYS	CA-C-N	5.25	125.78	120.38
1	B	748	LYS	C-N-CA	5.25	125.78	120.38
1	B	887	THR	OG1-CB-CG2	5.19	119.68	109.30
1	A	962	ILE	CB-CA-C	5.14	118.37	110.55

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	746	MET	Peptide
1	A	747	LYS	Peptide
1	A	841	ARG	Sidechain
1	B	746	MET	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2773	0	2806	10	12
1	B	2773	0	2806	23	11
2	A	20	0	0	0	0
2	B	20	0	0	1	0
All	All	5586	0	5612	31	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:749:PRO:HB2	1:B:750:PRO:HD2	1.52	0.91
1:A:886:VAL:HG11	1:B:934:HIS:CD2	2.10	0.85
1:B:841:ARG:HD2	1:B:873:LEU:O	1.80	0.81
1:A:841:ARG:HD2	1:A:873:LEU:O	1.81	0.81
1:B:743:ASP:HA	1:B:745:GLY:O	1.88	0.73
1:B:746:MET:O	1:B:747:LYS:HG2	1.90	0.71
1:B:717:GLN:OE1	1:B:887:THR:HB	1.92	0.70
1:B:744:PHE:CE2	1:B:749:PRO:HG3	2.30	0.66
1:A:841:ARG:CD	1:A:873:LEU:O	2.44	0.66
1:B:841:ARG:CD	1:B:873:LEU:O	2.46	0.64
1:B:749:PRO:HB2	1:B:750:PRO:CD	2.26	0.62
1:B:766:ASP:OD2	2:B:1101:7U9:NAB	2.33	0.61
1:A:689:TYR:OH	1:A:767:ASN:ND2	2.36	0.59
1:B:752:LEU:HD21	1:B:761:LYS:HE3	1.87	0.57
1:A:723:GLY:O	1:A:725:SER:N	2.40	0.55
1:B:747:LYS:O	1:B:748:LYS:HB2	2.10	0.51
1:B:744:PHE:CE2	1:B:749:PRO:CG	2.94	0.51
1:B:919:ILE:HG21	1:B:1005:LEU:HD11	1.93	0.50
1:A:717:GLN:OE1	1:A:887:THR:HB	2.14	0.47
1:A:934:HIS:CD2	1:B:886:VAL:HG11	2.51	0.45
1:B:744:PHE:CD2	1:B:749:PRO:HG3	2.51	0.45
1:B:719:ALA:HB3	1:B:729:ILE:HD11	1.99	0.45
1:B:675:MET:HE1	1:B:1004:LYS:HD2	1.97	0.45
1:A:676:ILE:HD11	1:A:797:LEU:HD11	1.98	0.45
1:B:744:PHE:CD2	1:B:749:PRO:CB	3.00	0.45
1:A:916:ILE:HD11	1:A:1006:LYS:HD2	2.00	0.44
1:B:712:ILE:HD11	1:B:735:ARG:HD2	2.01	0.43
1:B:705:GLN:OE1	1:B:705:GLN:CA	2.66	0.42
1:A:916:ILE:CD1	1:A:1006:LYS:HD2	2.49	0.42
1:B:705:GLN:OE1	1:B:705:GLN:HA	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:720:VAL:O	1:B:722:GLN:N	2.54	0.40

All (12) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:914:ASP:N	1:B:748:LYS:CB[4_445]	1.27	0.93
1:A:914:ASP:N	1:B:748:LYS:CG[4_445]	1.32	0.88
1:A:913:GLY:C	1:B:748:LYS:CG[4_445]	1.82	0.38
1:A:912:GLN:O	1:B:748:LYS:N[4_445]	1.89	0.31
1:A:913:GLY:C	1:B:748:LYS:CB[4_445]	1.89	0.31
1:A:914:ASP:N	1:B:748:LYS:CD[4_445]	1.97	0.23
1:A:747:LYS:O	1:A:827:ASN:OD1[4_445]	1.99	0.21
1:A:912:GLN:O	1:B:747:LYS:C[4_445]	2.00	0.20
1:A:913:GLY:N	1:B:748:LYS:CG[4_445]	2.00	0.20
1:A:913:GLY:CA	1:B:748:LYS:CG[4_445]	2.08	0.12
1:A:715:GLU:OE1	1:B:705:GLN:NE2[1_455]	2.10	0.10
1:A:912:GLN:O	1:B:748:LYS:CA[4_445]	2.10	0.10

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/352 (99%)	344 (98%)	4 (1%)	2 (1%)	21	13
1	B	350/352 (99%)	339 (97%)	5 (1%)	6 (2%)	7	2
All	All	700/704 (99%)	683 (98%)	9 (1%)	8 (1%)	10	4

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	748	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	749	PRO
1	B	663	SER
1	B	721	SER
1	A	662	LYS
1	A	748	LYS
1	B	747	LYS
1	B	750	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/309 (101%)	299 (96%)	12 (4%)	28	21
1	B	311/309 (101%)	290 (93%)	21 (7%)	14	7
All	All	622/618 (101%)	589 (95%)	33 (5%)	20	12

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	667	LYS
1	A	687	VAL
1	A	700	LYS
1	A	701	LEU
1	A	717	GLN
1	A	728	GLN
1	A	778	LEU
1	A	795	GLU
1	A	887	THR
1	A	940	LYS
1	A	962	ILE
1	A	963	SER
1	B	662	LYS
1	B	687	VAL
1	B	702	SER
1	B	703	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	705	GLN
1	B	713	LEU
1	B	722	GLN
1	B	730	LEU
1	B	739	LEU
1	B	746	MET
1	B	748	LYS
1	B	749	PRO
1	B	769	LEU
1	B	778	LEU
1	B	853	GLN
1	B	865	ARG
1	B	887	THR
1	B	940	LYS
1	B	943	LYS
1	B	962	ILE
1	B	980	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	660	HIS
1	A	718	GLN
1	A	767	ASN
1	A	855	HIS
1	A	961	ASN
1	A	998	ASN
1	B	660	HIS
1	B	759	GLN
1	B	868	ASN
1	B	980	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	7U9	A	1101	-	22,22,22	2.71	10 (45%)	25,32,32	3.54	15 (60%)
2	7U9	B	1101	-	22,22,22	2.43	9 (40%)	25,32,32	4.03	14 (56%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7U9	A	1101	-	-	0/10/21/21	0/3/3/3
2	7U9	B	1101	-	-	1/10/21/21	0/3/3/3

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1101	7U9	CAJ-NAT	6.31	1.58	1.46
2	A	1101	7U9	CAI-NAT	5.76	1.59	1.47
2	A	1101	7U9	CAG-CAQ	-4.99	1.31	1.39
2	A	1101	7U9	CAN-CAP	-4.86	1.37	1.50
2	A	1101	7U9	CAQ-CAR	-4.32	1.36	1.40
2	B	1101	7U9	CAN-CAP	-4.24	1.39	1.50
2	A	1101	7U9	CAO-CAM	-4.01	1.44	1.50
2	A	1101	7U9	CAJ-NAT	3.94	1.54	1.46
2	B	1101	7U9	CAO-CAR	-3.42	1.36	1.41
2	B	1101	7U9	CAO-CAM	-3.25	1.45	1.50
2	B	1101	7U9	CAQ-CAR	-2.95	1.37	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1101	7U9	CAJ-CAS	2.94	1.57	1.52
2	B	1101	7U9	CAI-NAT	2.76	1.52	1.47
2	B	1101	7U9	CAH-CAS	-2.68	1.49	1.53
2	A	1101	7U9	CAN-NAT	2.60	1.39	1.34
2	B	1101	7U9	CAR-NAK	-2.48	1.35	1.39
2	A	1101	7U9	CAR-NAK	-2.23	1.35	1.39
2	A	1101	7U9	CAH-CAS	-2.17	1.49	1.53
2	A	1101	7U9	CAO-CAR	-2.03	1.38	1.41

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1101	7U9	CAR-NAK-CAP	10.60	113.16	103.98
2	B	1101	7U9	CAI-NAT-CAJ	-8.31	102.68	112.38
2	B	1101	7U9	NAL-CAP-NAK	-7.64	105.05	113.22
2	A	1101	7U9	CAI-NAT-CAJ	-7.28	103.88	112.38
2	A	1101	7U9	CAR-NAK-CAP	7.18	110.19	103.98
2	A	1101	7U9	CAR-CAQ-NAL	6.75	108.97	105.13
2	B	1101	7U9	CAQ-CAR-NAK	-6.38	104.78	110.31
2	A	1101	7U9	CAQ-CAR-NAK	-5.01	105.97	110.31
2	B	1101	7U9	CAO-CAR-CAQ	5.00	123.89	120.25
2	A	1101	7U9	OAC-CAM-NAA	-4.91	115.52	122.62
2	B	1101	7U9	CAH-CAI-NAT	4.46	108.67	102.93
2	A	1101	7U9	CAP-CAN-NAT	4.25	125.73	119.04
2	B	1101	7U9	OAC-CAM-NAA	-4.15	116.61	122.62
2	A	1101	7U9	NAL-CAP-NAK	-4.07	108.86	113.22
2	A	1101	7U9	CAI-NAT-CAN	4.06	139.12	123.86
2	B	1101	7U9	CAR-CAQ-NAL	3.77	107.28	105.13
2	A	1101	7U9	CAF-CAE-CAG	-3.59	115.63	120.24
2	A	1101	7U9	CAO-CAR-CAQ	3.35	122.69	120.25
2	B	1101	7U9	CAP-CAN-NAT	3.27	124.19	119.04
2	B	1101	7U9	OAD-CAN-NAT	-3.23	117.56	122.67
2	B	1101	7U9	CAO-CAM-NAA	2.53	122.21	118.29
2	A	1101	7U9	CAH-CAI-NAT	-2.49	99.74	102.93
2	B	1101	7U9	CAQ-NAL-CAP	2.42	109.69	106.85
2	A	1101	7U9	OAD-CAN-NAT	-2.41	118.86	122.67
2	A	1101	7U9	CAJ-NAT-CAN	-2.34	116.85	123.77
2	A	1101	7U9	CAO-CAM-NAA	2.18	121.67	118.29
2	B	1101	7U9	CAG-CAQ-CAR	-2.08	120.61	122.57
2	B	1101	7U9	CAN-CAP-NAK	2.07	126.99	123.96
2	A	1101	7U9	CAG-CAQ-CAR	-2.03	120.66	122.57

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1101	7U9	OAC-CAM-CAO-CAR


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1101	7U9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	351/352 (99%)	0.42	29 (8%) 17 18	7, 18, 39, 76	3 (0%)
1	B	351/352 (99%)	0.91	48 (13%) 6 7	9, 21, 44, 146	3 (0%)
All	All	702/704 (99%)	0.66	77 (10%) 10 11	7, 19, 41, 146	6 (0%)

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	745	GLY	11.5
1	B	747	LYS	8.5
1	A	661	MET	8.3
1	B	1011	THR	8.3
1	B	746	MET	7.7
1	B	749	PRO	7.7
1	B	660	HIS	6.7
1	B	661	MET	6.3
1	A	660	HIS	5.9
1	B	748	LYS	5.7
1	A	1011	THR	5.3
1	B	962	ILE	5.0
1	B	662	LYS	4.5
1	B	980	ASN	4.5
1	B	750	PRO	4.5
1	A	747	LYS	4.4
1	B	981	ASP	4.4
1	B	726	ASP	4.2
1	B	943	LYS	4.0
1	B	744	PHE	3.9
1	A	728	GLN	3.8
1	B	730	LEU	3.6
1	B	723	GLY	3.5
1	B	731	ASP	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	966	GLY	3.4
1	A	722	GLN	3.4
1	B	705	GLN	3.3
1	B	978	GLY	3.3
1	A	689	TYR	3.3
1	B	979	VAL	3.2
1	A	694	GLN	3.2
1	A	727	SER	3.1
1	B	974	GLY	3.1
1	B	722	GLN	3.1
1	B	725	SER	3.1
1	A	691	ILE	3.0
1	B	752	LEU	3.0
1	B	956	PRO	2.9
1	A	729	ILE	2.9
1	A	806	ARG	2.9
1	A	807	ASP	2.8
1	B	827	ASN	2.7
1	A	743	ASP	2.7
1	B	961	ASN	2.7
1	B	779	ARG	2.7
1	A	726	ASP	2.6
1	A	744	PHE	2.6
1	B	975	ILE	2.6
1	B	959	SER	2.6
1	B	795	GLU	2.4
1	B	727	SER	2.4
1	A	723	GLY	2.4
1	A	781	GLY	2.4
1	A	662	LYS	2.4
1	B	960	ALA	2.4
1	A	687	VAL	2.4
1	B	964	LEU	2.3
1	B	853	GLN	2.3
1	B	695	LYS	2.3
1	A	725	SER	2.3
1	A	780	GLY	2.2
1	B	785	SER	2.2
1	A	980	ASN	2.2
1	B	694	GLN	2.2
1	A	719	ALA	2.2
1	A	720	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	756	ASP	2.2
1	B	944	GLY	2.2
1	B	664	LYS	2.2
1	B	806	ARG	2.1
1	A	718	GLN	2.1
1	A	846	GLN	2.1
1	A	809	GLU	2.1
1	B	684	LYS	2.1
1	A	751	LEU	2.1
1	B	720	VAL	2.0
1	B	854	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	7U9	A	1101	20/20	0.93	0.12	9,13,43,52	0
2	7U9	B	1101	20/20	0.94	0.08	9,14,29,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 7U9 A 1101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.