



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:19 AM UTC

PDB ID : 1WTD / pdb_00001wtd
Title : Crystal structure of type II restriction endonuclease, EcoO109I DNA-free form
Authors : Hashimoto, H.; Shimizu, T.; Imasaki, T.; Kato, M.; Shichijo, N.; Kita, K.; Sato, M.
Deposited on : 2004-11-22
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	10% 82% 14% • •
1	B	272	11% 80% 16% • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	B	308	-	X	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4230 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EcoO109IR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			2038	1308	344	379	7			
1	B	267	Total	C	N	O	S	0	0	0
			2087	1337	354	388	8			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	24	Total	O	0	0
			24	24		
3	B	21	Total	O	0	0
			21	21		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: EcoO109IR



• Molecule 1: EcoO109IR



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	175.55Å 175.55Å 44.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.40 20.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.8 (20.00-2.40) 98.8 (20.00-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.61 (at 2.41Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.255 , 0.297 0.264 , 0.302	Depositor DCC
R_{free} test set	1343 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	47.2	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 34.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4230	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	0/2083	0.97	3/2828 (0.1%)
1	B	0.84	0/2135	0.93	2/2894 (0.1%)
All	All	0.87	0/4218	0.95	5/5722 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	MET	N-CA-C	-10.87	80.56	111.00
1	B	99	GLU	CA-C-N	7.12	128.74	119.84
1	B	99	GLU	C-N-CA	7.12	128.74	119.84
1	A	1	MET	CB-CA-C	7.02	123.44	110.10
1	A	2	ASN	N-CA-C	-5.66	100.95	109.95

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2038	0	1909	28	0
1	B	2087	0	1963	28	0
2	A	18	0	24	2	0
2	B	42	0	55	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	24	0	0	0	0
3	B	21	0	0	0	0
All	All	4230	0	3951	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1:MET:N	1:B:208:ASN:OD1	2.20	0.74
1:A:61:ALA:HB1	2:B:308:GOL:H12	1.72	0.70
1:A:1:MET:N	1:A:208:ASN:HD21	1.91	0.69
1:A:65:MET:HE2	2:B:308:GOL:O1	1.95	0.65
1:A:65:MET:HE2	2:B:308:GOL:C1	2.29	0.63
1:A:61:ALA:HB1	2:B:308:GOL:C1	2.28	0.62
1:B:13:CYS:SG	1:B:83:ARG:NH1	2.74	0.61
1:B:184:ALA:O	1:B:185:ASN:CB	2.49	0.60
1:A:1:MET:H1	1:A:208:ASN:HD21	1.50	0.60
1:A:62:LYS:NZ	1:B:69:ASP:OD2	2.30	0.58
1:A:1:MET:H1	1:A:208:ASN:ND2	2.03	0.57
1:B:250:TYR:HA	1:B:269:MET:HE1	1.89	0.55
1:A:1:MET:N	1:A:208:ASN:ND2	2.56	0.53
1:B:63:HIS:CE1	2:B:310:GOL:H2	2.44	0.53
1:A:141:ASN:HD21	1:A:189:ILE:H	1.55	0.52
1:A:75:LEU:HD22	1:A:80:ILE:HG12	1.93	0.51
1:B:59:ILE:HG12	2:B:307:GOL:O1	2.11	0.51
1:B:249:LYS:O	1:B:269:MET:HE1	2.10	0.51
1:A:215:GLN:HE22	1:B:271:ASN:HB3	1.76	0.51
1:A:130:TRP:CZ3	1:A:160:VAL:HG11	2.45	0.50
1:B:208:ASN:O	1:B:209:ASN:ND2	2.45	0.50
1:B:61:ALA:HB3	2:B:308:GOL:H32	1.93	0.49
1:B:21:ARG:HD3	1:B:68:HIS:ND1	2.28	0.49
1:A:215:GLN:HE22	1:B:271:ASN:CB	2.25	0.49
1:A:61:ALA:HB1	2:B:308:GOL:H31	1.94	0.49
1:A:65:MET:CE	2:B:308:GOL:O1	2.60	0.49
1:A:1:MET:H1	1:A:208:ASN:CG	2.20	0.49
1:B:191:ILE:HD12	1:B:195:VAL:HG21	1.94	0.49
1:B:128:GLY:HA2	1:B:164:TYR:CE2	2.48	0.48
1:A:49:LEU:HD13	1:A:55:LEU:HA	1.96	0.47
1:B:3:LYS:HA	1:B:217:TRP:CE3	2.49	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:CYS:HG	2:A:304:GOL:HO3	1.62	0.46
1:B:90:LEU:HD11	1:B:112:VAL:HG23	1.97	0.45
1:A:271:ASN:CB	1:B:215:GLN:HE22	2.28	0.45
1:A:191:ILE:HD12	1:A:195:VAL:HG21	1.98	0.45
1:B:266:LEU:HD22	1:B:270:ILE:HD12	1.98	0.45
1:B:218:VAL:O	1:B:219:LEU:C	2.61	0.44
1:A:30:SER:HB2	1:A:246:VAL:HG22	1.99	0.43
1:B:62:LYS:HZ3	2:B:307:GOL:C1	2.31	0.43
1:B:174:TYR:CD1	1:B:197:VAL:HG11	2.53	0.43
1:A:61:ALA:O	1:A:65:MET:HB2	2.20	0.42
1:A:37:PRO:HB2	1:A:267:LEU:HA	2.02	0.41
1:B:23:SER:O	2:B:302:GOL:H2	2.20	0.41
1:B:61:ALA:CB	2:B:308:GOL:H32	2.50	0.41
1:A:40:THR:HG22	1:A:55:LEU:HD21	2.03	0.41
1:A:60:ILE:CD1	1:A:243:LYS:HA	2.51	0.41
1:B:99:GLU:CB	1:B:100:PRO:HD2	2.51	0.41
1:A:23:SER:O	2:A:306:GOL:H2	2.21	0.40
1:A:271:ASN:HB3	1:B:215:GLN:HE22	1.86	0.40
1:B:128:GLY:O	1:B:162:VAL:CG1	2.70	0.40
1:B:166:ASN:C	1:B:166:ASN:HD22	2.29	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/272 (96%)	236 (91%)	18 (7%)	6 (2%)	5	6
1	B	263/272 (97%)	245 (93%)	13 (5%)	5 (2%)	6	8
All	All	523/544 (96%)	481 (92%)	31 (6%)	11 (2%)	5	7

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	103	HIS
1	A	104	PRO
1	B	100	PRO
1	B	176	ILE
1	B	185	ASN
1	A	180	ILE
1	B	188	VAL
1	A	33	MET
1	A	211	GLU
1	A	118	GLY
1	B	134	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/237 (84%)	187 (94%)	13 (6%)	15	27
1	B	208/237 (88%)	195 (94%)	13 (6%)	16	29
All	All	408/474 (86%)	382 (94%)	26 (6%)	16	28

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	33	MET
1	A	75	LEU
1	A	109	ILE
1	A	112	VAL
1	A	113	ILE
1	A	115	ARG
1	A	134	LEU
1	A	151	TYR
1	A	209	ASN
1	A	234	LYS
1	A	267	LEU
1	A	272	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	53	ASP
1	B	66	THR
1	B	75	LEU
1	B	89	LYS
1	B	135	THR
1	B	166	ASN
1	B	168	HIS
1	B	202	GLU
1	B	207	LEU
1	B	209	ASN
1	B	219	LEU
1	B	266	LEU
1	B	267	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	22	GLN
1	A	141	ASN
1	A	157	ASN
1	A	168	HIS
1	A	186	HIS
1	A	196	HIS
1	A	209	ASN
1	A	215	GLN
1	A	235	ASN
1	A	256	ASN
1	B	22	GLN
1	B	114	GLN
1	B	133	GLN
1	B	149	ASN
1	B	166	ASN
1	B	209	ASN
1	B	215	GLN
1	B	235	ASN
1	B	262	GLN
1	B	271	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	304	-	5,5,5	0.50	0	5,5,5	0.65	0
2	GOL	B	301	-	5,5,5	0.45	0	5,5,5	0.40	0
2	GOL	A	306	-	5,5,5	0.69	0	5,5,5	1.25	1 (20%)
2	GOL	B	307	-	5,5,5	0.65	0	5,5,5	1.04	0
2	GOL	B	302	-	5,5,5	0.79	0	5,5,5	1.82	2 (40%)
2	GOL	B	308	-	5,5,5	1.29	1 (20%)	5,5,5	2.05	3 (60%)
2	GOL	B	310	-	5,5,5	0.30	0	5,5,5	0.38	0
2	GOL	B	305	-	5,5,5	0.51	0	5,5,5	1.80	1 (20%)
2	GOL	A	309	-	5,5,5	0.66	0	5,5,5	0.76	0
2	GOL	B	303	-	5,5,5	0.40	0	5,5,5	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	304	-	-	2/4/4/4	-
2	GOL	B	301	-	-	4/4/4/4	-
2	GOL	A	306	-	-	2/4/4/4	-
2	GOL	B	307	-	-	0/4/4/4	-
2	GOL	B	302	-	-	2/4/4/4	-
2	GOL	B	308	-	-	2/4/4/4	-
2	GOL	B	310	-	-	3/4/4/4	-
2	GOL	B	305	-	-	4/4/4/4	-
2	GOL	A	309	-	-	2/4/4/4	-
2	GOL	B	303	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	308	GOL	O2-C2	-2.40	1.36	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	305	GOL	O3-C3-C2	2.70	122.53	110.38
2	B	302	GOL	O3-C3-C2	2.55	121.86	110.38
2	B	308	GOL	C3-C2-C1	2.50	120.95	111.80
2	B	302	GOL	C3-C2-C1	2.47	120.86	111.80
2	B	308	GOL	O3-C3-C2	2.36	121.02	110.38
2	A	306	GOL	O3-C3-C2	2.24	120.46	110.38
2	B	308	GOL	O2-C2-C1	-2.09	100.55	109.18

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	304	GOL	O1-C1-C2-C3
2	B	301	GOL	O1-C1-C2-C3
2	B	302	GOL	C1-C2-C3-O3
2	B	302	GOL	O2-C2-C3-O3
2	B	303	GOL	O1-C1-C2-C3
2	B	305	GOL	O1-C1-C2-C3
2	B	305	GOL	C1-C2-C3-O3
2	B	308	GOL	C1-C2-C3-O3
2	B	308	GOL	O2-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	310	GOL	O1-C1-C2-C3
2	B	301	GOL	O1-C1-C2-O2
2	A	309	GOL	C1-C2-C3-O3
2	B	301	GOL	C1-C2-C3-O3
2	B	303	GOL	O1-C1-C2-O2
2	B	305	GOL	O2-C2-C3-O3
2	B	310	GOL	O1-C1-C2-O2
2	A	304	GOL	O1-C1-C2-O2
2	B	305	GOL	O1-C1-C2-O2
2	B	301	GOL	O2-C2-C3-O3
2	A	306	GOL	O1-C1-C2-O2
2	A	309	GOL	O2-C2-C3-O3
2	A	306	GOL	C1-C2-C3-O3
2	B	310	GOL	O2-C2-C3-O3

There are no ring outliers.

6 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	304	GOL	1	0
2	A	306	GOL	1	0
2	B	307	GOL	2	0
2	B	302	GOL	1	0
2	B	308	GOL	8	0
2	B	310	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/272 (97%)	0.85	28 (10%) 11 8	33, 49, 89, 96	0
1	B	267/272 (98%)	0.87	29 (10%) 10 8	30, 53, 91, 95	0
All	All	531/544 (97%)	0.86	57 (10%) 11 8	30, 51, 91, 96	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	177	LEU	6.2
1	A	133	GLN	4.6
1	A	179	GLY	4.5
1	A	185	ASN	4.2
1	B	136	MET	3.8
1	B	173	LYS	3.6
1	B	86	GLY	3.5
1	A	184	ALA	3.4
1	B	187	ASN	3.3
1	B	191	ILE	3.3
1	B	94	TYR	3.2
1	A	183	GLY	3.2
1	B	100	PRO	3.1
1	B	132	ILE	3.1
1	A	188	VAL	3.0
1	A	105	CYS	3.0
1	A	91	ASP	2.9
1	A	101	PHE	2.9
1	A	152	PRO	2.9
1	A	189	ILE	2.8
1	B	184	ALA	2.8
1	B	186	HIS	2.8
1	A	118	GLY	2.7
1	B	183	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	141	ASN	2.7
1	A	127	ALA	2.7
1	B	192	ARG	2.6
1	B	15	ALA	2.6
1	B	185	ASN	2.6
1	A	113	ILE	2.6
1	A	169	GLY	2.6
1	A	90	LEU	2.5
1	A	187	ASN	2.5
1	B	135	THR	2.5
1	B	161	GLY	2.5
1	B	134	LEU	2.5
1	B	156	ASP	2.4
1	B	176	ILE	2.4
1	A	191	ILE	2.4
1	B	189	ILE	2.4
1	B	118	GLY	2.3
1	A	148	ILE	2.3
1	B	102	ILE	2.3
1	B	154	VAL	2.2
1	B	131	THR	2.2
1	A	136	MET	2.2
1	A	120	ILE	2.2
1	A	92	LYS	2.2
1	A	182	THR	2.1
1	B	210	GLY	2.1
1	A	1	MET	2.1
1	A	186	HIS	2.1
1	B	196	HIS	2.1
1	B	95	ARG	2.1
1	B	85	PHE	2.1
1	A	137	ALA	2.1
1	A	171	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	B	303	6/6	0.61	0.17	82,83,83,83	0
2	GOL	A	304	6/6	0.69	0.18	76,78,78,79	0
2	GOL	B	301	6/6	0.72	0.18	73,74,75,75	0
2	GOL	A	306	6/6	0.72	0.28	47,50,53,54	0
2	GOL	B	305	6/6	0.75	0.22	46,50,51,51	0
2	GOL	B	302	6/6	0.82	0.19	38,40,41,44	0
2	GOL	B	307	6/6	0.83	0.18	37,41,42,44	0
2	GOL	B	308	6/6	0.84	0.22	43,45,47,49	0
2	GOL	A	309	6/6	0.86	0.17	48,53,54,54	0
2	GOL	B	310	6/6	0.86	0.14	63,65,66,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.