



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:47 AM UTC

PDB ID : 2WTN / pdb_00002wtm
Title : Ferulic Acid bound to Est1E from *Butyrivibrio proteoclasticus*
Authors : Goldstone, D.C.; Arcus, V.L.
Deposited on : 2009-09-17
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	83% 16% .
1	B	251	86% 12% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	1249	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 4191 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EST1E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	0	2	0
			1963	1255	325	370	13			
1	B	250	Total	C	N	O	S	0	2	0
			1957	1252	323	369	13			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 3-(4-HYDROXY-3-METHOXYPHENYL)-2-PROPENOIC ACID (CCD ID: FER) (formula: C₁₀H₁₀O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			14	10	4		
3	B	1	Total	C	O	0	0
			14	10	4		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	108	Total	O	0	0
			108	108		
5	B	94	Total	O	0	0
			94	94		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: EST1E



• Molecule 1: EST1E



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	144.28Å 48.18Å 97.03Å 90.00° 127.74° 90.00°	Depositor
Resolution (Å)	50.00 – 2.10 50.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.8 (50.00-2.10) 95.8 (50.00-2.10)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.86 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.164 , 0.225 0.175 , 0.226	Depositor DCC
R_{free} test set	1479 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	32.1	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4191	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, GOL, FER

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.41	9/2011 (0.4%)	1.18	6/2724 (0.2%)
1	B	1.41	3/2005 (0.1%)	1.26	7/2719 (0.3%)
All	All	1.41	12/4016 (0.3%)	1.22	13/5443 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	1	2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	134	ILE	CA-CB	7.14	1.57	1.54
1	A	199	ALA	CA-CB	-6.46	1.43	1.53
1	A	234	VAL	CA-CB	6.45	1.62	1.54
1	B	121	ILE	C-O	-6.40	1.17	1.23
1	A	114	ALA	CA-C	6.05	1.60	1.52
1	A	176	ILE	CA-CB	-5.84	1.47	1.54
1	A	213	LYS	C-O	-5.63	1.17	1.24
1	A	212	TYR	CA-C	5.55	1.59	1.52
1	A	102	ALA	CA-C	5.41	1.59	1.52
1	A	50	ASN	CG-ND2	-5.36	1.22	1.33
1	A	102	ALA	C-O	-5.07	1.17	1.23
1	B	136	GLU	CG-CD	5.03	1.64	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	20	ASN	N-CA-C	8.93	129.54	109.81
1	B	149	ASP	CA-C-N	-7.11	112.22	120.12
1	B	149	ASP	C-N-CA	-7.11	112.22	120.12
1	B	197	ASP	CA-C-N	6.75	130.23	120.38
1	B	197	ASP	C-N-CA	6.75	130.23	120.38
1	A	208	PHE	N-CA-C	6.74	118.71	111.36
1	A	107	GLY	CA-C-N	5.99	126.59	119.94
1	A	107	GLY	C-N-CA	5.99	126.59	119.94
1	A	229	HIS	N-CA-C	5.65	117.52	111.36
1	B	123	LYS	N-CA-C	5.57	117.43	111.36
1	B	121	ILE	CB-CA-C	-5.37	106.14	111.30
1	A	153	ILE	CA-C-N	-5.17	114.91	120.03
1	A	153	ILE	C-N-CA	-5.17	114.91	120.03

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	20	ASN	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	18	LYS	Peptide
1	B	19	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1963	0	1947	21	0
1	B	1957	0	1928	16	0
2	A	18	0	24	4	0
2	B	18	0	24	2	0
3	A	14	0	9	2	0
3	B	14	0	9	1	0
4	A	5	0	0	0	0
5	A	108	0	0	2	0
5	B	94	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4191	0	3941	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:200:VAL:HG12	5:A:2088:HOH:O	1.82	0.79
1:A:200:VAL:CG1	5:A:2088:HOH:O	2.36	0.74
1:B:1:MET:HE1	1:B:89:TYR:CZ	2.33	0.64
1:A:111:VAL:HG23	1:A:125:LEU:HD11	1.81	0.62
1:A:177[B]:ARG:NE	1:B:180:ASP:OD2	2.37	0.55
1:B:137:ILE:HG13	1:B:143:LEU:HD13	1.89	0.54
1:A:104:HIS:ND1	2:A:1249:GOL:H12	2.23	0.53
1:A:169:TYR:CD2	1:A:169:TYR:C	2.87	0.53
1:A:175:THR:HG22	2:B:1250:GOL:H2	1.93	0.51
1:A:41:HIS:NE2	2:A:1249:GOL:H11	2.27	0.49
1:B:123:LYS:NZ	1:B:245:GLN:O	2.46	0.49
1:A:8:ILE:HG23	1:A:68:ASP:HB2	1.95	0.49
1:A:32:GLY:HA3	2:A:1249:GOL:H2	1.95	0.48
1:A:54:VAL:HG12	1:A:55:ALA:O	2.14	0.47
1:B:105:SER:OG	3:B:1249:FER:C9	2.63	0.47
1:A:104:HIS:ND1	2:A:1249:GOL:C1	2.78	0.47
1:A:187:LYS:HB3	1:A:188:PRO:CD	2.44	0.47
1:A:105:SER:OG	3:A:1250:FER:O1	2.33	0.47
1:A:220:ILE:HG22	1:A:223:ASP:HB2	1.97	0.46
1:A:185:TYR:CZ	1:A:187:LYS:HB2	2.51	0.46
1:B:1:MET:HB2	1:B:1:MET:HE3	1.68	0.45
1:B:125:LEU:HG	1:B:127:PRO:HD3	1.98	0.45
1:B:187:LYS:HB3	1:B:188:PRO:CD	2.46	0.45
1:B:160:TRP:O	1:B:163:ARG:HD2	2.17	0.44
1:B:184:LYS:HG3	2:B:1250:GOL:H32	1.99	0.44
1:A:33:PHE:HB3	3:A:1250:FER:C8	2.48	0.43
1:A:78:LYS:HE2	1:B:6:ASP:OD2	2.19	0.43
1:B:63:GLY:HA2	1:B:69:GLY:O	2.20	0.42
1:B:19:ASN:CB	1:B:20:ASN:HA	2.50	0.42
1:B:185:TYR:CZ	1:B:187:LYS:HB2	2.56	0.41
1:A:100:TYR:CG	1:A:242:MET:HE1	2.56	0.41
1:A:243:LEU:O	1:A:244:GLU:C	2.63	0.40
1:A:248:LYS:OXT	1:A:248:LYS:CG	2.69	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:92:LYS:HE2	1:B:92:LYS:HB2	1.87	0.40
1:B:29:ILE:HD13	1:B:29:ILE:HG21	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/251 (99%)	237 (95%)	12 (5%)	0	100	100
1	B	250/251 (100%)	240 (96%)	8 (3%)	2 (1%)	16	12
All	All	499/502 (99%)	477 (96%)	20 (4%)	2 (0%)	30	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	19	ASN
1	B	20	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/212 (99%)	206 (98%)	4 (2%)	50	58
1	B	207/212 (98%)	201 (97%)	6 (3%)	37	42

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	417/424 (98%)	407 (98%)	10 (2%)	43 49

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	157	LEU
1	A	203	GLU
1	A	244	GLU
1	B	1	MET
1	B	9	LYS
1	B	23	LYS
1	B	133	MET
1	B	198	GLU
1	B	233	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	152	ASN
1	A	229	HIS
1	A	245	GLN
1	B	11	ASN
1	B	46	GLN
1	B	50	ASN
1	B	152	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	B	1251	-	5,5,5	0.58	0	5,5,5	0.40	0
2	GOL	A	1251	-	5,5,5	0.44	0	5,5,5	0.59	0
3	FER	A	1250	-	14,14,14	1.77	5 (35%)	18,18,18	1.42	2 (11%)
2	GOL	B	1250	-	5,5,5	0.40	0	5,5,5	0.64	0
2	GOL	B	1252	-	5,5,5	0.82	0	5,5,5	1.82	3 (60%)
2	GOL	A	1249	-	5,5,5	0.91	0	5,5,5	1.34	1 (20%)
4	PO4	A	1253	-	4,4,4	1.03	0	6,6,6	0.76	0
3	FER	B	1249	-	14,14,14	1.42	2 (14%)	18,18,18	1.86	4 (22%)
2	GOL	A	1252	-	5,5,5	0.43	0	5,5,5	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	1251	-	-	0/4/4/4	-
2	GOL	A	1251	-	-	2/4/4/4	-
3	FER	A	1250	-	-	0/7/7/7	0/1/1/1
2	GOL	B	1250	-	-	1/4/4/4	-
2	GOL	B	1252	-	-	0/4/4/4	-
2	GOL	A	1249	-	-	2/4/4/4	-
3	FER	B	1249	-	-	2/7/7/7	0/1/1/1
2	GOL	A	1252	-	-	2/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1250	FER	C8-C7	3.49	1.41	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1249	FER	C1-C7	-3.18	1.38	1.47
3	A	1250	FER	O2-C9	-3.10	1.22	1.30
3	A	1250	FER	C8-C9	-2.47	1.42	1.48
3	A	1250	FER	C1-C7	-2.35	1.40	1.47
3	A	1250	FER	O3-C10	2.15	1.48	1.42
3	B	1249	FER	C8-C7	2.05	1.38	1.33

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1249	FER	C10-O3-C3	-5.57	109.34	117.51
3	A	1250	FER	C10-O3-C3	-4.27	111.25	117.51
3	B	1249	FER	C1-C7-C8	-3.59	119.09	126.92
3	B	1249	FER	O3-C3-C4	3.40	119.64	114.55
2	A	1249	GOL	O1-C1-C2	2.75	122.75	110.38
3	A	1250	FER	O3-C3-C4	2.74	118.66	114.55
2	B	1252	GOL	O2-C2-C1	-2.46	99.00	109.18
3	B	1249	FER	O3-C3-C2	-2.30	120.12	124.08
2	B	1252	GOL	C3-C2-C1	2.26	120.08	111.80
2	B	1252	GOL	O3-C3-C2	2.01	119.44	110.38

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1252	GOL	O2-C2-C3-O3
3	B	1249	FER	C2-C1-C7-C8
3	B	1249	FER	C6-C1-C7-C8
2	A	1251	GOL	C1-C2-C3-O3
2	A	1252	GOL	C1-C2-C3-O3
2	A	1251	GOL	O2-C2-C3-O3
2	A	1249	GOL	O2-C2-C3-O3
2	B	1250	GOL	O2-C2-C3-O3
2	A	1249	GOL	C1-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1250	FER	2	0
2	B	1250	GOL	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1249	GOL	4	0
3	B	1249	FER	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	249/251 (99%)	-0.06	3 (1%) 76 78	15, 34, 51, 64	2 (0%)
1	B	250/251 (99%)	-0.15	3 (1%) 76 78	16, 31, 49, 66	2 (0%)
All	All	499/502 (99%)	-0.10	6 (1%) 76 78	15, 33, 51, 66	4 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	20	ASN	5.0
1	A	0	ALA	2.3
1	A	247	ALA	2.3
1	B	160	TRP	2.2
1	B	161	ASP	2.1
1	A	248	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	1252	6/6	0.89	0.12	41,52,54,58	0
2	GOL	B	1252	6/6	0.89	0.10	44,48,50,51	0
2	GOL	A	1251	6/6	0.90	0.11	33,42,43,44	0
3	FER	A	1250	14/14	0.90	0.10	30,47,49,52	0
2	GOL	B	1250	6/6	0.91	0.11	37,41,42,43	0
2	GOL	A	1249	6/6	0.92	0.09	30,42,44,44	0
2	GOL	B	1251	6/6	0.92	0.10	41,49,51,56	0
3	FER	B	1249	14/14	0.93	0.08	34,42,44,45	0
4	PO4	A	1253	5/5	0.94	0.08	79,79,80,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.