



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 07:17 PM UTC

PDB ID : 3WT2 / pdb_00003wt2
Title : Crystal structure of the b'-a' domain of thermophilic fungal protein disulfide isomerase (oxidized form)
Authors : Inagaki, K.; Satoh, T.; Itoh, S.G.; Okumura, H.; Kato, K.
Deposited on : 2014-04-02
Resolution : 3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	83% 13% ..
1	B	247	81% 16% ..
1	C	247	84% 13% ..

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 5637 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein disulfide-isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	242	1879	1214	299	363	3	0	0	0
1	B	242	1879	1214	299	363	3	0	0	0
1	C	242	1879	1214	299	363	3	0	0	0


There are 15 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	203	GLY	-	expression tag	UNP P55059
A	204	PRO	-	expression tag	UNP P55059
A	205	LEU	-	expression tag	UNP P55059
A	206	GLY	-	expression tag	UNP P55059
A	207	SER	-	expression tag	UNP P55059
B	203	GLY	-	expression tag	UNP P55059
B	204	PRO	-	expression tag	UNP P55059
B	205	LEU	-	expression tag	UNP P55059
B	206	GLY	-	expression tag	UNP P55059
B	207	SER	-	expression tag	UNP P55059
C	203	GLY	-	expression tag	UNP P55059
C	204	PRO	-	expression tag	UNP P55059
C	205	LEU	-	expression tag	UNP P55059
C	206	GLY	-	expression tag	UNP P55059
C	207	SER	-	expression tag	UNP P55059

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein disulfide-isomerase

Chain A:  83% 13% ..





- Molecule 1: Protein disulfide-isomerase

Chain B:  81% 16% ..



- Molecule 1: Protein disulfide-isomerase

Chain C:  84% 13% ..



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	96.31Å 96.31Å 69.49Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 3.30 20.00 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-3.30) 99.3 (20.00-3.30)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.55 (at 3.29Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.221 , 0.313 (Not available) , 0.303	Depositor DCC
R_{free} test set	516 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	77.7	Xtrriage
Anisotropy	0.113	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 31.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.000 for -h,-k,l 0.038 for h,-h-k,-l 0.000 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5637	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 66.15 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.3330e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.86	0/1922	1.05	2/2604 (0.1%)
1	B	0.82	1/1922 (0.1%)	1.04	3/2604 (0.1%)
1	C	0.84	0/1922	1.04	2/2604 (0.1%)
All	All	0.84	1/5766 (0.0%)	1.04	7/7812 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	415	LEU	CA-C	-5.05	1.46	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	403	VAL	CA-C-N	7.57	126.84	118.97
1	C	403	VAL	C-N-CA	7.57	126.84	118.97
1	B	403	VAL	CA-C-N	7.00	126.51	119.24
1	B	403	VAL	C-N-CA	7.00	126.51	119.24
1	A	403	VAL	CA-C-N	6.70	126.21	119.24
1	A	403	VAL	C-N-CA	6.70	126.21	119.24
1	B	218	TYR	N-CA-C	5.33	117.52	111.02

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1879	0	1869	16	0
1	B	1879	0	1869	21	0
1	C	1879	0	1869	19	0
All	All	5637	0	5607	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:388:LYS:HD2	1:C:369:LYS:HZ1	1.57	0.68
1:B:388:LYS:HD2	1:C:369:LYS:NZ	2.08	0.67
1:A:446:TYR:O	1:A:447:LYS:C	2.42	0.63
1:B:345:TYR:CE1	1:B:404:PRO:HG2	2.36	0.60
1:C:446:TYR:O	1:C:447:LYS:C	2.45	0.60
1:B:446:TYR:O	1:B:447:LYS:C	2.44	0.59
1:C:345:TYR:CE1	1:C:404:PRO:HG2	2.39	0.57
1:A:345:TYR:CE1	1:A:404:PRO:HG2	2.40	0.56
1:B:389:ASP:OD1	1:B:389:ASP:N	2.36	0.56
1:B:441:ALA:O	1:B:447:LYS:HA	2.09	0.51
1:A:441:ALA:O	1:A:447:LYS:HA	2.12	0.50
1:C:441:ALA:O	1:C:447:LYS:HA	2.12	0.50
1:B:233:ALA:HB3	1:B:239:ARG:HG3	1.94	0.50
1:C:381:LEU:CD2	1:C:437:ILE:HD11	2.42	0.50
1:A:389:ASP:OD1	1:A:389:ASP:N	2.37	0.49
1:A:233:ALA:HB3	1:A:239:ARG:HG3	1.94	0.49
1:C:233:ALA:HB3	1:C:239:ARG:HG3	1.95	0.49
1:C:369:LYS:O	1:C:369:LYS:HG3	2.03	0.48
1:A:381:LEU:CD2	1:A:437:ILE:HD11	2.43	0.48
1:B:381:LEU:CD2	1:B:437:ILE:HD11	2.43	0.47
1:B:410:PHE:HA	1:B:411:PRO:C	2.40	0.47
1:B:388:LYS:HD2	1:C:369:LYS:CE	2.43	0.47
1:B:304:THR:O	1:B:305:PHE:C	2.60	0.45
1:C:354:LYS:HE3	1:C:389:ASP:O	2.16	0.45
1:C:432:THR:HG23	1:C:435:ASP:H	1.81	0.45
1:B:354:LYS:HE3	1:B:389:ASP:O	2.16	0.45
1:B:346:ASN:OD1	1:B:350:LEU:HD12	2.17	0.45
1:A:247:LYS:HB3	1:A:248:PRO:HD3	1.98	0.45
1:B:388:LYS:HG3	1:C:369:LYS:HZ3	1.82	0.45
1:A:354:LYS:HE3	1:A:389:ASP:O	2.16	0.44
1:A:432:THR:HG23	1:A:435:ASP:H	1.82	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:323:SER:C	1:B:324:ILE:HG13	2.42	0.44
1:C:247:LYS:HB3	1:C:248:PRO:HD3	1.99	0.44
1:B:432:THR:HG23	1:B:435:ASP:H	1.83	0.44
1:B:247:LYS:HB3	1:B:248:PRO:HD3	2.00	0.44
1:A:346:ASN:OD1	1:A:350:LEU:HD12	2.17	0.43
1:C:346:ASN:OD1	1:C:350:LEU:HD12	2.18	0.43
1:A:304:THR:O	1:A:305:PHE:C	2.59	0.43
1:A:410:PHE:HA	1:A:411:PRO:C	2.43	0.42
1:A:446:TYR:O	1:A:448:ALA:N	2.53	0.42
1:C:304:THR:O	1:C:305:PHE:C	2.62	0.42
1:B:297:PHE:O	1:B:298:ASP:C	2.62	0.41
1:C:381:LEU:HD23	1:C:437:ILE:HD11	2.02	0.41
1:A:350:LEU:HA	1:A:421:LYS:HG2	2.02	0.41
1:B:350:LEU:HA	1:B:421:LYS:HG2	2.02	0.41
1:C:410:PHE:HA	1:C:411:PRO:C	2.45	0.41
1:B:319:LYS:HE3	1:C:237:GLU:OE1	2.20	0.41
1:A:305:PHE:C	1:A:305:PHE:CD2	2.99	0.40
1:B:381:LEU:HD23	1:B:437:ILE:HD11	2.03	0.40
1:C:305:PHE:C	1:C:305:PHE:CD2	2.99	0.40
1:A:344:ASN:OD1	1:A:344:ASN:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	240/247 (97%)	230 (96%)	8 (3%)	2 (1%)	16	45
1	B	240/247 (97%)	230 (96%)	8 (3%)	2 (1%)	16	45
1	C	240/247 (97%)	230 (96%)	8 (3%)	2 (1%)	16	45
All	All	720/741 (97%)	690 (96%)	24 (3%)	6 (1%)	16	45

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	447	LYS
1	B	447	LYS
1	C	447	LYS
1	A	351	ASP
1	B	351	ASP
1	C	351	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/199 (99%)	185 (94%)	12 (6%)	17	45
1	B	197/199 (99%)	185 (94%)	12 (6%)	17	45
1	C	197/199 (99%)	185 (94%)	12 (6%)	17	45
All	All	591/597 (99%)	555 (94%)	36 (6%)	17	45

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	209	LEU
1	A	254	ARG
1	A	256	VAL
1	A	261	THR
1	A	278	THR
1	A	332	LYS
1	A	333	GLN
1	A	365	CYS
1	A	369	LYS
1	A	389	ASP
1	A	421	LYS
1	A	434	GLU
1	B	209	LEU
1	B	254	ARG
1	B	256	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	261	THR
1	B	278	THR
1	B	332	LYS
1	B	333	GLN
1	B	365	CYS
1	B	369	LYS
1	B	389	ASP
1	B	421	LYS
1	B	434	GLU
1	C	209	LEU
1	C	254	ARG
1	C	256	VAL
1	C	261	THR
1	C	278	THR
1	C	332	LYS
1	C	333	GLN
1	C	365	CYS
1	C	369	LYS
1	C	389	ASP
1	C	421	LYS
1	C	434	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	273	ASN
1	A	292	ASN
1	B	273	ASN
1	B	292	ASN
1	B	443	ASN
1	C	273	ASN
1	C	292	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	242/247 (97%)	-0.62	0 100 100	44, 69, 110, 136	0
1	B	242/247 (97%)	-0.35	1 (0%) 88 79	50, 85, 120, 151	0
1	C	242/247 (97%)	-0.31	0 100 100	56, 89, 128, 159	0
All	All	726/741 (97%)	-0.43	1 (0%) 92 90	44, 82, 124, 159	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	278	THR	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.