



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:19 PM UTC

PDB ID : 6WT3 / pdb\_00006wt3  
Title : Structural basis for the binding of monoclonal antibody 5D2 to the tryptophan-rich lipid-binding loop in lipoprotein lipase  
Authors : Luz, J.G.; Birrane, G.; Young, S.G.; Meiyappan, M.; Ploug, M.  
Deposited on : 2020-05-01  
Resolution : 2.85 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	223	
2	L	214	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5D2 FAB HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	216	1685	1081	272	326	6	0	1	0

- Molecule 2 is a protein called 5D2 FAB LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	212	1627	1012	271	335	9	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	2	Total	O	0	0
			2	2		
3	L	5	Total	O	0	0
			5	5		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5D2 FAB HEAVY CHAIN



- Molecule 2: 5D2 FAB LIGHT CHAIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	138.79Å 37.07Å 107.96Å 90.00° 126.19° 90.00°	Depositor
Resolution (Å)	43.57 – 2.85 43.57 – 2.85	Depositor EDS
% Data completeness (in resolution range)	94.8 (43.57-2.85) 94.8 (43.57-2.85)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.51 (at 2.86Å)	Xtrriage
Refinement program	PHENIX (1.18_3855: ???)	Depositor
R, $R_{free}$	0.253 , 0.308 0.253 , 0.305	Depositor DCC
$R_{free}$ test set	498 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.0	Xtrriage
Anisotropy	0.270	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 6.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	3319	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	H	0.15	0/1738	0.39	0/2372
2	L	0.18	0/1666	0.37	0/2266
All	All	0.16	0/3404	0.38	0/4638

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1685	0	1636	21	1
2	L	1627	0	1554	22	2
3	H	2	0	0	0	0
3	L	5	0	0	0	0
All	All	3319	0	3190	42	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:183:SER:H	1:H:223:ASN:HD21	1.29	0.81

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:211:ASN:HD22	2:L:231:ASN:HD22	1.40	0.67
2:L:222:SER:OG	2:L:224:SER:O	2.14	0.64
2:L:24:ILE:HG12	2:L:49:SER:HB3	1.80	0.63
1:H:183:SER:H	1:H:223:ASN:ND2	1.98	0.58
1:H:79:TYR:HE1	1:H:89:LEU:HG	1.71	0.56
1:H:119:GLU:OE1	1:H:122:ASN:ND2	2.41	0.54
1:H:86:LYS:NZ	1:H:104:ARG:O	2.42	0.53
2:L:204:LYS:O	2:L:208:GLU:HG3	2.09	0.53
1:H:54:HIS:CD2	1:H:69[B]:TRP:HB3	2.44	0.52
1:H:235:LYS:NZ	2:L:144:GLU:OE1	2.40	0.52
1:H:57:LYS:HB2	1:H:67:ILE:HD11	1.93	0.51
2:L:166:ASN:HB2	2:L:218:THR:HG22	1.93	0.50
2:L:60:LYS:HD3	2:L:105:ALA:HB2	1.94	0.49
2:L:31:ALA:HA	2:L:123:THR:HA	1.93	0.49
1:H:178:THR:O	1:H:224:VAL:HA	2.12	0.49
2:L:196:MET:HE1	2:L:198:SER:HB2	1.95	0.48
2:L:23:GLN:N	2:L:114:SER:HG	2.11	0.48
1:H:199:SER:OG	1:H:200:ASP:N	2.43	0.48
2:L:140:PRO:HB3	2:L:230:PHE:CE1	2.50	0.47
1:H:215:TRP:CG	1:H:216:PRO:HA	2.50	0.47
1:H:170:LYS:HG2	1:H:171:GLY:N	2.28	0.47
1:H:146:PRO:HB3	1:H:172:TYR:HB3	1.98	0.45
1:H:174:PRO:O	1:H:175:GLU:HB2	2.16	0.45
1:H:86:LYS:HE3	1:H:109:ASP:OD2	2.16	0.45
2:L:23:GLN:HB2	2:L:116:PRO:HD2	1.98	0.44
2:L:24:ILE:HD13	2:L:51:VAL:HG23	2.00	0.44
1:H:174:PRO:HD2	1:H:226:HIS:NE2	2.32	0.44
2:L:165:ILE:HG13	2:L:218:THR:O	2.18	0.44
2:L:80:PRO:HB2	2:L:82:ARG:HG2	1.99	0.43
2:L:161:TYR:CG	2:L:162:PRO:HA	2.54	0.43
2:L:111:GLN:HG2	2:L:113:SER:H	1.83	0.43
2:L:34:SER:HB3	2:L:128:LYS:HE3	2.01	0.42
2:L:34:SER:OG	2:L:126:GLU:OE1	2.27	0.42
1:H:51:TYR:HB3	1:H:117:ARG:HG3	2.01	0.42
2:L:111:GLN:HE21	2:L:113:SER:N	2.18	0.42
2:L:184:TRP:CE2	2:L:196:MET:HG3	2.55	0.42
1:H:69[B]:TRP:CE2	1:H:78:LYS:HB3	2.56	0.41
2:L:180:VAL:HG12	2:L:181:LEU:N	2.37	0.40
1:H:67:ILE:HG21	1:H:100:MET:HE2	2.03	0.40
1:H:53:ILE:HG13	1:H:98:VAL:HG21	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:81:GLU:OE2	2:L:78:GLY:N[1_565]	2.12	0.08
2:L:91:SER:OG	2:L:131:ASP:OD2[4_556]	2.13	0.07

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	213/223 (96%)	193 (91%)	19 (9%)	1 (0%)	24	42
2	L	210/214 (98%)	190 (90%)	18 (9%)	2 (1%)	12	25
All	All	423/437 (97%)	383 (90%)	37 (9%)	3 (1%)	18	35

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	173	PHE
2	L	75	LEU
2	L	30	PRO

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	190/194 (98%)	177 (93%)	13 (7%)	14	30
2	L	187/188 (100%)	174 (93%)	13 (7%)	14	29

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	377/382 (99%)	351 (93%)	26 (7%)	14	30

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	86	LYS
1	H	94	SER
1	H	119	GLU
1	H	132	GLN
1	H	147	SER
1	H	175	GLU
1	H	178	THR
1	H	205	SER
1	H	213	SER
1	H	217	SER
1	H	219	THR
1	H	221	THR
1	H	222	CYS
2	L	36	SER
2	L	51	VAL
2	L	84	SER
2	L	90	THR
2	L	127	LEU
2	L	137	SER
2	L	143	SER
2	L	164	ASP
2	L	189	SER
2	L	197	SER
2	L	212	SER
2	L	222	SER
2	L	223	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	54	HIS
1	H	62	GLN
1	H	122	ASN
1	H	132	GLN
1	H	223	ASN
2	L	28	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	L	59	GLN
2	L	111	GLN
2	L	115	ASN
2	L	159	ASN
2	L	166	ASN
2	L	211	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	H	216/223 (96%)	0.94	18 (8%) 17 13	20, 37, 39, 41	1 (0%)
2	L	212/214 (99%)	1.09	21 (9%) 13 10	33, 36, 39, 40	0
All	All	428/437 (97%)	1.01	39 (9%) 15 11	20, 37, 39, 41	1 (0%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	198	GLN	3.9
2	L	74	ASN	3.7
2	L	29	SER	3.6
2	L	62	ARG	3.6
1	H	173	PHE	3.6
1	H	174	PRO	3.1
2	L	88	SER	3.1
2	L	30	PRO	3.1
2	L	201	THR	2.8
2	L	186	ASP	2.8
1	H	123	TYR	2.7
1	H	154	GLY	2.7
2	L	32	LEU	2.7
1	H	71	TYR	2.7
2	L	77	SER	2.6
1	H	92	ASP	2.6
1	H	205	SER	2.5
2	L	49	SER	2.4
1	H	131	GLY	2.4
2	L	102	GLU	2.4
1	H	203	THR	2.4
2	L	223	THR	2.4
2	L	179	GLY	2.3
2	L	78	GLY	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	H	73	GLY	2.3
1	H	120	ASP	2.3
2	L	100	GLU	2.2
1	H	79	TYR	2.2
1	H	189	GLY	2.2
1	H	199	SER	2.2
1	H	178	THR	2.1
2	L	86	SER	2.1
2	L	75	LEU	2.1
1	H	167	CYS	2.1
2	L	126	GLU	2.1
2	L	216	GLU	2.1
1	H	175	GLU	2.1
2	L	63	SER	2.0
2	L	173	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.