



wwPDB EM Validation Summary Report ⓘ

Mar 19, 2026 – 08:36 PM UTC

PDB ID : 7WTX / pdb_00007wtx
EMDB ID : EMD-32804
Title : Cryo-EM structure of a human pre-40S ribosomal subunit - State RRP12-B1
Authors : Cheng, J.; Lau, B.; Thoms, M.; Ameismeier, M.; Berninghausen, O.; Hurt, E.; Beckmann, R.
Deposited on : 2022-02-05
Resolution : 3.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : **FAILED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

i

ELECTRON MICROSCOPY

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Q-score	25397	14724 (2.60 - 3.60)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition

There are 35 unique types of molecules in this entry. The entry contains 85074 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 18S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	2	1605	Total	C	N	O	P	0	0
			34277	15298	6153	11222	1604		

- Molecule 2 is a protein called 40S ribosomal protein S5.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	F	189	Total	C	N	O	S	0	0
			1495	934	284	270	7		

- Molecule 3 is a protein called 40S ribosomal protein S12.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	M	123	Total	C	N	O	S	0	0
			953	598	169	177	9		

- Molecule 4 is a protein called 40S ribosomal protein S15.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	P	120	Total	C	N	O	S	0	0
			984	625	184	168	7		

- Molecule 5 is a protein called 40S ribosomal protein S16.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	Q	122	Total	C	N	O	S	0	0
			969	616	180	170	3		

- Molecule 6 is a protein called 40S ribosomal protein S17.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	R	112	Total	C	N	O	S	0	0
			911	571	169	168	3		

- Molecule 7 is a protein called 40S ribosomal protein S18.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	S	130	Total	C	N	O	S	0	0
			1083	686	214	182	1		

- Molecule 8 is a protein called 40S ribosomal protein S19.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	T	144	Total	C	N	O	S	0	0
			1122	703	217	199	3		

- Molecule 9 is a protein called 40S ribosomal protein S25.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	Z	72	Total	C	N	O	S	0	0
			574	368	104	101	1		

- Molecule 10 is a protein called 40S ribosomal protein S28.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	c	61	Total	C	N	O	S	0	0
			471	288	95	86	2		

- Molecule 11 is a protein called Ubiquitin-40S ribosomal protein S27a.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	f	61	Total	C	N	O	S	0	0
			498	316	94	81	7		

- Molecule 12 is a protein called 40S ribosomal protein SA.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	A	216	Total	C	N	O	S	0	0
			1705	1083	299	315	8		

- Molecule 13 is a protein called 40S ribosomal protein S3a.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	B	213	Total	C	N	O	S	0	0
			1729	1098	309	308	14		

- Molecule 14 is a protein called 40S ribosomal protein S2.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	C	218	Total	C	N	O	S	0	0
			1690	1094	289	297	10		

- Molecule 15 is a protein called 40S ribosomal protein S4, X isoform.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	E	262	Total	C	N	O	S	0	0
			2076	1324	386	358	8		

- Molecule 16 is a protein called 40S ribosomal protein S6.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	G	230	Total	C	N	O	S	0	0
			1862	1164	371	320	7		

- Molecule 17 is a protein called 40S ribosomal protein S7.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	H	186	Total	C	N	O	S	0	0
			1501	957	276	267	1		

- Molecule 18 is a protein called 40S ribosomal protein S8.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	I	205	Total	C	N	O	S	0	0
			1682	1056	331	290	5		

- Molecule 19 is a protein called 40S ribosomal protein S9.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	J	180	Total	C	N	O	S	0	0
			1499	955	300	242	2		

- Molecule 20 is a protein called 40S ribosomal protein S11.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	L	151	Total	C	N	O	S	0	0
			1229	782	230	211	6		

- Molecule 21 is a protein called 40S ribosomal protein S13.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	N	149	Total	C	N	O	S	0	0
			1202	770	228	203	1		

- Molecule 22 is a protein called 40S ribosomal protein S14.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	O	135	Total	C	N	O	S	0	0
			1009	618	198	187	6		

- Molecule 23 is a protein called 40S ribosomal protein S21.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	V	82	Total	C	N	O	S	0	0
			625	384	116	120	5		

- Molecule 24 is a protein called 40S ribosomal protein S15a.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	W	129	Total	C	N	O	S	0	0
			1034	659	193	176	6		

- Molecule 25 is a protein called 40S ribosomal protein S23.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	X	141	Total	C	N	O	S	0	0
			1098	693	219	183	3		

- Molecule 26 is a protein called 40S ribosomal protein S24.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	Y	124	Total	C	N	O	S	0	0
			1014	641	198	170	5		

- Molecule 27 is a protein called 40S ribosomal protein S27.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	b	82	Total	C	N	O	S	0	0
			640	402	118	113	7		

- Molecule 28 is a protein called 40S ribosomal protein S30.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	e	55	Total	C	N	O	S	0	0
			438	271	95	71	1		

- Molecule 29 is a protein called RNA-binding protein PNO1.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	x	178	Total	C	N	O	S	0	0
			1391	891	252	244	4		

- Molecule 30 is a protein called RNA-binding protein NOB1.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	y	325	Total	C	N	O	S	0	0
			2568	1622	473	463	10		

- Molecule 31 is a protein called Pre-rRNA-processing protein TSR1 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	u	629	Total	C	N	O	S	0	0
			5062	3249	902	887	24		

- Molecule 32 is a protein called Bystin.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	w	275	Total	C	N	O	S	0	0
			2248	1455	397	387	9		

- Molecule 33 is a protein called Protein LTV1 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
33	t	132	Total	C	N	O	S	0	0
			1131	701	212	214	4		

- Molecule 34 is a protein called RRP12-like protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	K	940	Total	C	N	O	S	0	0
			7302	4680	1266	1311	45		

- Molecule 35 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
35	f	1	Total 1	Zn 1	0
35	y	1	Total 1	Zn 1	0

MolProbity failed to run properly - this section is therefore empty.

3 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	27437	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION; Relion	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	44	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.498	Depositor
Minimum map value	-0.260	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.011	Depositor
Recommended contour level	0.02	Depositor
Map size ($\text{\AA}$)	381.24, 381.24, 381.24	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.059, 1.059, 1.059	Depositor

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


5 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-32804. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

5.1 Orthogonal projections [i](#)


5.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

5.2 Central slices [i](#)


5.2.1 Primary map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

5.3 Largest variance slices [i](#)


5.3.1 Primary map



X Index: 144



Y Index: 133




Z Index: 181


The images above show the largest variance slices of the map in three orthogonal directions.

5.4 Orthogonal standard-deviation projections (False-color) [i](#)


5.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

5.5 Orthogonal surface views [i](#)

5.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


5.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

6 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

6.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


6.2 Volume estimate [i](#)



The volume at the recommended contour level is 1418 nm^3 ; this corresponds to an approximate mass of 1281 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

6.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

7 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

7.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

7.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.10	-	-
Author-provided FSC curve	3.11	4.04	3.18
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


8 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-32804 and PDB model 7WTX. Per-residue inclusion information can be found in section ?? on page ??.

8.1 Map-model overlay [i](#)


This section was not generated.

8.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

8.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).

8.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 95% of all non-hydrogen atoms, are inside the map.

8.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9540	 0.4780
2	 0.9880	 0.5110
A	 0.9830	 0.5700
B	 0.9730	 0.5390
C	 0.9880	 0.5800
E	 0.9840	 0.5910
F	 0.9440	 0.4290
G	 0.9620	 0.5040
H	 0.9390	 0.4720
I	 0.9600	 0.5260
J	 0.9940	 0.6000
K	 0.7860	 0.1970
L	 0.9610	 0.5710
M	 0.7220	 0.2240
N	 0.9930	 0.5760
O	 0.9690	 0.5220
P	 0.9390	 0.4610
Q	 0.8940	 0.4160
R	 0.9120	 0.3240
S	 0.9110	 0.4010
T	 0.8860	 0.3800
V	 0.9800	 0.5730
W	 0.9910	 0.6160
X	 0.9980	 0.6220
Y	 0.9930	 0.5730
Z	 0.8410	 0.3270
b	 0.9860	 0.5550
c	 0.9490	 0.4070
e	 0.9480	 0.5300
f	 0.8990	 0.2570
t	 0.9520	 0.4070
u	 0.9780	 0.5400
w	 0.9410	 0.4050
x	 0.9880	 0.5620
y	 0.9430	 0.4510

