



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:29 PM UTC

PDB ID : 4WUP / pdb\_00004wup  
Title : Crystal structure of human carbonic anhydrase isozyme I with 4-[(2-Hydroxy ethyl)thio]benzenesulfonamide  
Authors : Smirnov, A.; Manakova, E.; Grazulis, S.  
Deposited on : 2014-11-03  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	 2% 88% 10%
1	B	260	 0% 90% 8%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbonic anhydrase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	256	2023	1278	352	390	3	0	4	0
1	B	257	2069	1307	359	400	3	0	7	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P00915
B	1	MET	-	initiating methionine	UNP P00915

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 4-[(2-hydroxyethyl)sulfanyl]benzenesulfonamide (CCD ID: 3UF) (formula: C<sub>8</sub>H<sub>11</sub>NO<sub>3</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	28	16	2	6	4	0	1
3	B	1	14	8	1	3	2	0	0

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula:  $C_2H_3O_2^-$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C O		
4	A	1	4	2 2	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	C O	0	0
			7	4 3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	172	Total	O	0	0
			172	172		
6	B	198	Total	O	0	0
			198	198		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Carbonic anhydrase 1



- Molecule 1: Carbonic anhydrase 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.33Å 71.28Å 120.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.36 – 1.75 60.36 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (60.36-1.75) 99.9 (60.36-1.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.32 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.185 , 0.232 0.182 , 0.230	Depositor DCC
$R_{free}$ test set	5608 reflections (10.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.8	Xtrriage
Anisotropy	0.547	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 31.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4517	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.30 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.8654e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, PEG, ZN, 3UF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.38	11/2081 (0.5%)	1.23	7/2831 (0.2%)
1	B	1.37	7/2130 (0.3%)	1.20	5/2895 (0.2%)
All	All	1.38	18/4211 (0.4%)	1.21	12/5726 (0.2%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	96	HIS	CE1-NE2	7.57	1.40	1.32
1	B	119	HIS	ND1-CE1	7.00	1.39	1.32
1	A	64	HIS	CE1-NE2	6.73	1.39	1.32
1	A	143	VAL	N-CA	6.27	1.53	1.46
1	B	107	HIS	ND1-CE1	6.23	1.38	1.32
1	B	122	HIS	ND1-CE1	5.73	1.38	1.32
1	B	94	HIS	C-O	5.65	1.30	1.23
1	A	96	HIS	CE1-NE2	5.59	1.38	1.32
1	B	119	HIS	CE1-NE2	5.57	1.38	1.32
1	A	59	ILE	C-O	5.44	1.29	1.24
1	A	53	PRO	C-O	-5.36	1.17	1.24
1	B	200	HIS	ND1-CE1	5.35	1.38	1.32
1	A	119	HIS	N-CA	5.33	1.52	1.46
1	A	107	HIS	CE1-NE2	5.33	1.37	1.32
1	A	231	SER	CA-C	5.28	1.59	1.52
1	A	243	HIS	CG-CD2	5.26	1.41	1.35
1	A	64	HIS	CG-CD2	5.17	1.41	1.35
1	A	166	ALA	N-CA	5.09	1.52	1.46

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	82	GLY	CA-C-N	7.67	128.96	120.45

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	82	GLY	C-N-CA	7.67	128.96	120.45
1	A	158	GLN	N-CA-C	6.67	119.16	111.02
1	A	154	ASN	CA-C-N	-6.52	113.09	119.87
1	A	154	ASN	C-N-CA	-6.52	113.09	119.87
1	B	180	ASP	CA-C-N	-5.30	114.57	120.45
1	B	180	ASP	C-N-CA	-5.30	114.57	120.45
1	B	251	LEU	N-CA-C	5.30	117.80	111.71
1	B	195	PRO	CA-C-N	-5.22	118.14	122.16
1	B	195	PRO	C-N-CA	-5.22	118.14	122.16
1	A	18	LYS	N-CA-C	-5.15	105.58	111.14
1	A	218	VAL	N-CA-C	-5.13	101.44	108.20

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2023	0	1922	6	0
1	B	2069	0	1971	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	28	0	22	2	0
3	B	14	0	11	1	0
4	A	4	0	3	0	0
5	B	7	0	10	0	0
6	A	172	0	0	0	0
6	B	198	0	0	0	0
All	All	4517	0	3939	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:190:ASP:HB2	1:B:213:LYS:HG2	1.84	0.60
1:B:101:ASN:ND2	1:B:220[B]:SER:OG	2.42	0.53
1:B:200:HIS:CE1	3:B:302:3UF:H5	2.45	0.52
1:A:58:GLU:OE1	1:A:173:ARG:HD3	2.12	0.50
1:B:45:LYS:O	1:B:82:GLY:HA2	2.12	0.48
1:A:102:GLU:HG3	1:A:103:HIS:ND1	2.28	0.48
1:B:201:PRO:HA	1:B:203:LEU:HG	1.98	0.46
1:A:127:LYS:HD3	1:A:139:ASP:OD2	2.18	0.42
1:B:26:ASN:HA	1:B:252:LYS:HD3	2.02	0.42
1:A:200:HIS:CE1	3:A:302[B]:3UF:H6	2.55	0.41
1:A:41:ASP:C	1:A:41:ASP:OD1	2.63	0.41
1:B:47:ILE:HD11	1:B:189:LEU:HB3	2.03	0.40
1:A:200:HIS:CE1	3:A:302[A]:3UF:H6	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/260 (99%)	252 (98%)	6 (2%)	0	100	100
1	B	261/260 (100%)	252 (97%)	9 (3%)	0	100	100
All	All	519/520 (100%)	504 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/226 (98%)	216 (98%)	5 (2%)	44	24
1	B	229/226 (101%)	224 (98%)	5 (2%)	45	26
All	All	450/452 (100%)	440 (98%)	10 (2%)	47	26

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LYS
1	A	48	SER
1	A	102	GLU
1	A	168	LYS
1	A	213	LYS
1	B	10	LYS
1	B	15[A]	GLN
1	B	15[B]	GLN
1	B	22	ILE
1	B	213	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	27	ASN
1	A	158	GLN
1	A	242	GLN
1	B	27	ASN
1	B	242	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	3UF	A	302[A]	2	14,14,14	2.61	4 (28%)	19,19,19	2.04	3 (15%)
5	PEG	B	303	-	6,6,6	0.85	0	5,5,5	0.77	0
3	3UF	A	302[B]	2	14,14,14	2.90	5 (35%)	19,19,19	2.20	6 (31%)
4	ACT	A	303	-	3,3,3	0.86	0	3,3,3	0.62	0
3	3UF	B	302	2	14,14,14	2.88	4 (28%)	19,19,19	2.80	8 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	3UF	A	302[B]	2	-	4/10/10/10	0/1/1/1
3	3UF	A	302[A]	2	-	6/10/10/10	0/1/1/1
3	3UF	B	302	2	-	5/10/10/10	0/1/1/1
5	PEG	B	303	-	-	3/4/4/4	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302[B]	3UF	O1-S2	6.29	1.54	1.43
3	A	302[A]	3UF	O1-S2	6.04	1.54	1.43
3	A	302[B]	3UF	S2-N3	6.03	1.72	1.60
3	B	302	3UF	O1-S2	5.50	1.53	1.43
3	A	302[A]	3UF	S2-N3	5.39	1.70	1.60

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	3UF	O4-S2	5.18	1.52	1.43
3	B	302	3UF	C8-S11	5.13	1.87	1.76
3	B	302	3UF	S2-N3	4.20	1.68	1.60
3	A	302[B]	3UF	O4-S2	3.66	1.50	1.43
3	A	302[B]	3UF	C8-S11	3.41	1.83	1.76
3	A	302[A]	3UF	C8-S11	2.98	1.82	1.76
3	A	302[A]	3UF	O4-S2	2.92	1.48	1.43
3	A	302[B]	3UF	C5-S2	2.17	1.80	1.77

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	3UF	O1-S2-O4	-6.45	108.88	118.80
3	A	302[B]	3UF	O1-S2-O4	-5.99	109.59	118.80
3	A	302[A]	3UF	O1-S2-O4	-5.60	110.18	118.80
3	A	302[A]	3UF	C5-S2-N3	5.40	115.92	108.40
3	A	302[B]	3UF	C5-S2-N3	5.21	115.66	108.40
3	B	302	3UF	O4-S2-C5	4.52	112.45	107.35
3	B	302	3UF	C10-C5-S2	-4.36	113.81	119.72
3	B	302	3UF	C7-C6-C5	-4.32	115.24	119.44
3	B	302	3UF	C12-S11-C8	4.00	111.45	104.03
3	B	302	3UF	C6-C7-C8	2.89	123.81	120.48
3	B	302	3UF	C10-C5-C6	2.85	124.19	120.47
3	B	302	3UF	C5-S2-N3	2.78	112.27	108.40
3	A	302[B]	3UF	O1-S2-C5	2.55	110.23	107.35
3	A	302[B]	3UF	C6-C5-S2	-2.52	116.30	119.72
3	A	302[A]	3UF	C6-C5-S2	-2.48	116.36	119.72
3	A	302[B]	3UF	C10-C5-S2	2.20	122.72	119.72
3	A	302[B]	3UF	C12-S11-C8	-2.20	99.93	104.03

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302[A]	3UF	S11-C12-C13-O14
3	A	302[B]	3UF	S11-C12-C13-O14
3	A	302[A]	3UF	C13-C12-S11-C8
3	B	302	3UF	C13-C12-S11-C8
5	B	303	PEG	O1-C1-C2-O2
5	B	303	PEG	O2-C3-C4-O4
5	B	303	PEG	C4-C3-O2-C2
3	B	302	3UF	C10-C5-S2-O4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	B	302	3UF	C7-C8-S11-C12
3	B	302	3UF	C6-C5-S2-O4
3	A	302[A]	3UF	C9-C8-S11-C12
3	A	302[A]	3UF	C10-C5-S2-O4
3	A	302[A]	3UF	C7-C8-S11-C12
3	B	302	3UF	C9-C8-S11-C12
3	A	302[B]	3UF	C10-C5-S2-O4
3	A	302[A]	3UF	C6-C5-S2-O4
3	A	302[B]	3UF	C7-C8-S11-C12
3	A	302[B]	3UF	C9-C8-S11-C12

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302[A]	3UF	1	0
3	A	302[B]	3UF	1	0
3	B	302	3UF	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	256/260 (98%)	0.01	6 (2%) 61 67	9, 24, 39, 63	4 (1%)
1	B	257/260 (98%)	-0.11	3 (1%) 76 82	8, 21, 35, 55	7 (2%)
All	All	513/520 (98%)	-0.05	9 (1%) 67 74	8, 22, 37, 63	11 (2%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	260[A]	PHE	6.4
1	A	40	HIS	3.0
1	A	42	THR	2.4
1	A	14	GLU	2.4
1	A	9	ASP	2.3
1	A	238	ALA	2.2
1	A	15	GLN	2.1
1	B	4	ASP	2.0
1	B	103	HIS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	PEG	B	303	7/7	0.71	0.21	44,54,60,60	0
3	3UF	A	302[B]	14/14	0.85	0.16	21,31,39,46	14
3	3UF	A	302[A]	14/14	0.85	0.16	16,25,36,40	14
4	ACT	A	303	4/4	0.87	0.11	32,33,34,35	0
3	3UF	B	302	14/14	0.95	0.13	15,28,57,66	0
2	ZN	B	301	1/1	1.00	0.01	13,13,13,13	0
2	ZN	A	301	1/1	1.00	0.01	15,15,15,15	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.