



Full wwPDB X-ray Structure Validation Report ⓘ

May 7, 2026 – 10:46 AM EDT

PDB ID : 2WVQ / pdb_00002wvq
Title : Structure of the HET-s N-terminal domain. Mutant D23A, P33H
Authors : Greenwald, J.; Buhtz, C.; Ritter, C.; Kwiatkowski, W.; Choe, S.; Saupe, S.J.; Riek, R.
Deposited on : 2009-10-19
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	
1	B	225	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3577 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SMALL S PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	0	5	0
			1705	1070	302	327	6			
1	B	207	Total	C	N	O	S	0	3	0
			1658	1039	296	317	6			

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	MET	-	expression tag	UNP Q03689
A	-2	ARG	-	expression tag	UNP Q03689
A	-1	GLY	-	expression tag	UNP Q03689
A	0	SER	-	expression tag	UNP Q03689
A	1	HIS	-	expression tag	UNP Q03689
A	2	HIS	-	expression tag	UNP Q03689
A	3	HIS	-	expression tag	UNP Q03689
A	4	HIS	-	expression tag	UNP Q03689
A	5	HIS	-	expression tag	UNP Q03689
A	6	HIS	-	expression tag	UNP Q03689
A	7	LEU	-	expression tag	UNP Q03689
A	8	VAL	-	expression tag	UNP Q03689
A	9	PRO	-	expression tag	UNP Q03689
A	10	ARG	-	expression tag	UNP Q03689
A	11	GLY	-	expression tag	UNP Q03689
A	12	SER	-	expression tag	UNP Q03689
A	23	ALA	ASP	engineered mutation	UNP Q03689
A	33	HIS	PRO	engineered mutation	UNP Q03689
B	-3	MET	-	expression tag	UNP Q03689
B	-2	ARG	-	expression tag	UNP Q03689
B	-1	GLY	-	expression tag	UNP Q03689
B	0	SER	-	expression tag	UNP Q03689
B	1	HIS	-	expression tag	UNP Q03689
B	2	HIS	-	expression tag	UNP Q03689
B	3	HIS	-	expression tag	UNP Q03689

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	4	HIS	-	expression tag	UNP Q03689
B	5	HIS	-	expression tag	UNP Q03689
B	6	HIS	-	expression tag	UNP Q03689
B	7	LEU	-	expression tag	UNP Q03689
B	8	VAL	-	expression tag	UNP Q03689
B	9	PRO	-	expression tag	UNP Q03689
B	10	ARG	-	expression tag	UNP Q03689
B	11	GLY	-	expression tag	UNP Q03689
B	12	SER	-	expression tag	UNP Q03689
B	23	ALA	ASP	engineered mutation	UNP Q03689
B	33	HIS	PRO	engineered mutation	UNP Q03689

- Molecule 2 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (CCD ID: DTT) (formula: $C_4H_{10}O_2S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	O	S	0	0
			8	4	2	2		

- Molecule 3 is (2R,3S)-1,4-DIMERCAPTOBUTANE-2,3-DIOL (CCD ID: DTU) (formula: $C_4H_{10}O_2S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	O	S	0	0
			8	4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	113	Total	O	0	0
			113	113		
4	B	85	Total	O	0	0
			85	85		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SMALL S PROTEIN



• Molecule 1: SMALL S PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.33Å 83.15Å 67.36Å 90.00° 104.55° 90.00°	Depositor
Resolution (Å)	65.23 – 2.00 65.20 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.1 (65.23-2.00) 96.1 (65.20-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.07 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.225 , 0.269 0.220 , 0.265	Depositor DCC
R_{free} test set	1697 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3577	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DTT, DTU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	2/1735 (0.1%)	1.14	4/2334 (0.2%)
1	B	0.95	3/1687 (0.2%)	1.15	5/2268 (0.2%)
All	All	0.95	5/3422 (0.1%)	1.14	9/4602 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	61	ILE	CA-CB	6.93	1.62	1.54
1	A	194	GLU	C-O	6.81	1.32	1.24
1	B	178	ALA	CA-CB	5.87	1.62	1.53
1	B	178	ALA	N-CA	5.57	1.53	1.46
1	B	48	ALA	CA-CB	5.13	1.61	1.53

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	28	VAL	N-CA-C	7.62	119.12	107.99
1	B	84	VAL	CB-CA-C	5.97	120.46	112.22
1	B	156	LEU	CA-CB-CG	5.67	136.13	116.30
1	A	84	VAL	CB-CA-C	5.50	119.81	112.22
1	A	132	LEU	CA-CB-CG	5.46	135.39	116.30
1	B	40	ARG	N-CA-C	5.08	117.71	111.82
1	A	84	VAL	CG1-CB-CG2	5.03	121.86	110.80
1	B	71	ALA	CA-C-N	5.02	124.93	119.76
1	B	71	ALA	C-N-CA	5.02	124.93	119.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1705	0	1705	31	0
1	B	1658	0	1659	30	0
2	B	8	0	9	1	0
3	B	8	0	9	0	0
4	A	113	0	0	6	0
4	B	85	0	0	2	1
All	All	3577	0	3382	59	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:GLU:OE2	1:A:198:THR:OG1	1.96	0.82
1:B:112:VAL:O	2:B:1222:DTT:H41	1.78	0.82
1:B:25:PHE:O	1:B:28:VAL:HG12	1.88	0.74
1:B:179:VAL:HG23	4:B:2072:HOH:O	1.91	0.70
1:B:160:VAL:CG1	1:B:207:ILE:HG21	2.22	0.69
1:A:75:LYS:O	1:A:79:LEU:HD13	1.93	0.68
1:A:70:ASP:CG	4:A:2045:HOH:O	2.35	0.68
1:A:39:GLU:HG3	1:A:118:MET:HG2	1.80	0.63
1:B:182:LYS:HE3	1:B:183:LEU:HD13	1.82	0.61
1:B:160:VAL:HG13	1:B:207:ILE:HG21	1.84	0.60
1:B:186:ILE:O	1:B:189:GLU:HG2	2.02	0.60
1:B:40:ARG:O	1:B:44[B]:ARG:HG3	2.04	0.58
1:A:103:LEU:HD23	1:B:79:LEU:HD13	1.83	0.58
1:A:201:LYS:HD2	1:A:219:ILE:HD11	1.84	0.57
1:A:17:LEU:HB3	1:A:84:VAL:HG13	1.86	0.56
1:B:66:ARG:HG2	1:B:77:VAL:HG21	1.86	0.56
1:A:194:GLU:HB2	4:A:2109:HOH:O	2.07	0.54
1:B:42:GLN:HG2	1:B:118:MET:CE	2.36	0.54
1:A:215:ALA:O	1:A:219:ILE:HG12	2.09	0.53
1:B:20:ASN:HB3	1:B:88:LEU:HD11	1.92	0.52
1:B:17:LEU:HB3	1:B:84:VAL:HG13	1.90	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:LEU:O	1:A:187:GLU:HG3	2.11	0.51
1:A:19:ASN:HD21	1:A:62:ASN:HD21	1.59	0.49
1:A:99:LYS:NZ	4:B:2025:HOH:O	2.41	0.49
1:A:128:ARG:HH21	1:A:190:GLU:HG2	1.77	0.49
1:B:219:ILE:HG13	1:B:220:ASP:N	2.28	0.49
1:A:79:LEU:HD12	1:B:103:LEU:HD23	1.94	0.48
1:A:114:GLU:HG3	1:A:116:LYS:HG2	1.94	0.48
1:A:165:ARG:NH1	4:A:2096:HOH:O	2.44	0.48
1:B:197:LEU:HD13	1:B:218:LYS:HB3	1.95	0.48
1:B:17:LEU:CB	1:B:84:VAL:CG1	2.91	0.48
1:B:17:LEU:HB2	1:B:84:VAL:CG1	2.43	0.48
1:B:87:ILE:O	1:B:90:LEU:HB3	2.14	0.47
1:A:26[A]:GLU:HG3	1:A:49:LYS:NZ	2.30	0.47
1:A:87:ILE:O	1:A:90:LEU:HB3	2.15	0.47
1:B:27:TYR:CE1	1:B:145:LYS:HB2	2.49	0.46
1:B:163:VAL:O	1:B:167:VAL:HG23	2.15	0.46
1:B:39:GLU:HG3	1:B:118:MET:HG2	1.97	0.45
1:A:199:ILE:HD13	4:A:2019:HOH:O	2.16	0.45
1:B:182:LYS:HE2	1:B:182:LYS:HB3	1.86	0.45
1:A:21:CYS:SG	1:A:88:LEU:HD13	2.57	0.45
1:B:13:ASN:ND2	1:B:16:GLY:H	2.15	0.45
1:A:86:GLU:HG2	4:A:2054:HOH:O	2.17	0.44
1:A:202:ASP:OD1	4:A:2111:HOH:O	2.21	0.44
1:B:83:ILE:CD1	1:B:169:GLU:HG3	2.48	0.43
1:A:25:PHE:HB2	1:A:49:LYS:HE3	2.00	0.43
1:B:42:GLN:HG2	1:B:118:MET:HE2	1.99	0.43
1:A:186:ILE:O	1:A:189:GLU:HG2	2.19	0.42
1:B:17:LEU:HB3	1:B:84:VAL:CG1	2.49	0.42
1:B:102:GLU:HG2	1:B:110:LEU:CD2	2.50	0.42
1:A:17:LEU:CB	1:A:84:VAL:HG13	2.50	0.42
1:A:53:SER:HB2	1:A:137:GLN:NE2	2.34	0.41
1:A:192:GLU:OE2	1:A:192:GLU:HA	2.21	0.41
1:A:34:PHE:HE1	1:A:150:LEU:O	2.02	0.41
1:B:64:ASP:OD1	1:B:66:ARG:NH1	2.54	0.41
1:A:14:VAL:HG22	1:A:81:LYS:HB2	2.03	0.41
1:A:121:ILE:HG23	1:A:122:GLY:N	2.35	0.41
1:B:115:ASP:O	1:B:118:MET:HB2	2.21	0.41
1:A:125:LEU:HD12	1:A:125:LEU:HA	1.98	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:2017:HOH:O	4:B:2074:HOH:O[2_657]	1.85	0.35

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/225 (95%)	208 (97%)	6 (3%)	0	100	100
1	B	206/225 (92%)	203 (98%)	3 (2%)	0	100	100
All	All	420/450 (93%)	411 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/189 (95%)	152 (84%)	28 (16%)	2	1
1	B	175/189 (93%)	146 (83%)	29 (17%)	2	1
All	All	355/378 (94%)	298 (84%)	57 (16%)	2	1

All (57) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ASN
1	A	28	VAL
1	A	32	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	36	ARG
1	A	40	ARG
1	A	45	LEU
1	A	59	VAL
1	A	84	VAL
1	A	88	LEU
1	A	89	LEU
1	A	99	LYS
1	A	104	VAL
1	A	114	GLU
1	A	125	LEU
1	A	129	LEU
1	A	132	LEU
1	A	133	VAL
1	A	140	THR
1	A	145	LYS
1	A	154	LYS
1	A	156	LEU
1	A	165	ARG
1	A	170	LEU
1	A	172	LYS
1	A	177	GLU
1	A	183	LEU
1	A	188	ILE
1	A	219	ILE
1	B	13	ASN
1	B	20	ASN
1	B	40	ARG
1	B	45	LEU
1	B	66	ARG
1	B	73	THR
1	B	74	ASP
1	B	79	LEU
1	B	84	VAL
1	B	88	LEU
1	B	89	LEU
1	B	103	LEU
1	B	104	VAL
1	B	106	ASP
1	B	119	LYS
1	B	125	LEU
1	B	129	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	132	LEU
1	B	133	VAL
1	B	146	THR
1	B	156	LEU
1	B	162	GLN
1	B	165	ARG
1	B	170	LEU
1	B	177	GLU
1	B	182	LYS
1	B	183	LEU
1	B	188	ILE
1	B	191	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	ASN
1	A	29	GLN
1	A	33	HIS
1	A	62	ASN
1	A	78	GLN
1	A	107	GLN
1	A	130	ASN
1	A	137	GLN
1	A	139	GLN
1	B	13	ASN
1	B	78	GLN
1	B	162	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	DTU	B	1223	2	7,7,7	0.70	0	4,8,8	0.80	0
2	DTT	B	1222	3	7,7,7	1.05	0	4,8,8	2.74	2 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DTU	B	1223	2	-	2/8/8/8	-
2	DTT	B	1222	3	-	3/8/8/8	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1222	DTT	C2-C1-S1	3.82	125.11	114.43
2	B	1222	DTT	O2-C2-C3	-3.42	102.39	109.57

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1222	DTT	S1-C1-C2-O2
2	B	1222	DTT	S1-C1-C2-C3
3	B	1223	DTU	S1-C1-C2-O2
3	B	1223	DTU	S1-C1-C2-C3
2	B	1222	DTT	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1222	DTT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	211/225 (93%)	1.06	39 (18%)	3 3	14, 37, 50, 60	5 (2%)
1	B	207/225 (92%)	1.06	29 (14%)	6 5	16, 36, 47, 63	3 (1%)
All	All	418/450 (92%)	1.06	68 (16%)	4 4	14, 36, 49, 63	8 (1%)

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	221	ALA	5.4
1	A	11	GLY	4.7
1	A	207	ILE	4.6
1	B	205	GLY	4.5
1	B	202[A]	ASP	4.4
1	A	205	GLY	4.3
1	A	195	ALA	4.3
1	A	146	THR	4.2
1	A	203	ALA	4.1
1	A	196	SER	4.1
1	A	191	VAL	4.1
1	A	186	ILE	3.9
1	B	219	ILE	3.9
1	B	27	TYR	3.8
1	B	207	ILE	3.8
1	B	10	ARG	3.8
1	A	143	ALA	3.8
1	A	199	ILE	3.6
1	B	146	THR	3.5
1	B	186	ILE	3.5
1	A	219	ILE	3.4
1	A	12	SER	3.4
1	B	188	ILE	3.4
1	A	190	GLU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	203	ALA	3.3
1	B	9	PRO	3.2
1	B	204	ALA	3.2
1	A	211	MET	3.2
1	A	142	LEU	3.1
1	B	199	ILE	3.1
1	B	206	GLY	3.1
1	B	221	ALA	3.1
1	A	197	LEU	3.1
1	B	198	THR	3.0
1	A	141	SER	3.0
1	A	194	GLU	3.0
1	B	220	ASP	2.8
1	A	188	ILE	2.8
1	A	200	LEU	2.7
1	B	195	ALA	2.6
1	A	108	GLN	2.6
1	A	144	LYS	2.5
1	A	192	GLU	2.5
1	A	183	LEU	2.4
1	B	11	GLY	2.4
1	A	204	ALA	2.4
1	A	220	ASP	2.4
1	B	145	LYS	2.4
1	A	209	ALA	2.3
1	B	137	GLN	2.3
1	A	145	LYS	2.3
1	B	121	ILE	2.3
1	A	175	PRO	2.2
1	B	182	LYS	2.2
1	B	170	LEU	2.2
1	A	27	TYR	2.2
1	B	196	SER	2.2
1	A	13	ASN	2.2
1	B	157	GLU	2.2
1	A	184	ALA	2.1
1	A	129	LEU	2.1
1	B	217	GLN	2.1
1	A	217	GLN	2.1
1	A	176	ILE	2.1
1	A	160	VAL	2.1
1	B	175	PRO	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	190	GLU	2.0
1	A	214	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	DTU	B	1223	8/8	0.81	0.17	59,62,64,67	0
2	DTT	B	1222	8/8	0.90	0.13	44,51,54,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.