



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:12 AM UTC

PDB ID : 4WVB / pdb_00004wvb
Title : Crystal structure of GH63 mannosylglycerate hydrolase from *Thermus thermophilus* HB8 in complex with glucose
Authors : Miyazaki, T.; Ichikawa, M.; Nishikawa, A.; Tonozuka, T.
Deposited on : 2014-11-05
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	5% 88% 6% 6%
1	B	420	7% 91% 8% .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7302 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	394	Total	C	N	O	S	0	0	0
			3246	2109	565	567	5			
1	B	417	Total	C	N	O	S	0	4	0
			3468	2250	608	605	5			

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	B	1	Total	Cl	0	0
			1	1		

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		
3	A	1	Total	C	O	0	0
			8	6	2		
3	B	1	Total	C	O	0	0
			8	6	2		
3	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	249	Total	O	0	0
			249	249		
5	B	293	Total	O	0	0
			293	293		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein



- Molecule 1: Uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	93.44Å 93.44Å 254.14Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.74 – 1.77 29.74 – 1.77	Depositor EDS
% Data completeness (in resolution range)	99.2 (29.74-1.77) 99.2 (29.74-1.77)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.65 (at 1.77Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.158 , 0.187 0.172 , 0.200	Depositor DCC
R_{free} test set	4028 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	19.0	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.016 for -h-k,k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7302	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, MPD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/3358	0.83	7/4576 (0.2%)
1	B	0.76	0/3599	0.85	1/4903 (0.0%)
All	All	0.71	0/6957	0.84	8/9479 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	VAL	N-CA-C	6.44	117.96	110.62
1	A	310	THR	CA-C-N	-6.12	113.98	120.03
1	A	310	THR	C-N-CA	-6.12	113.98	120.03
1	A	348	ARG	N-CA-C	5.71	117.21	109.11
1	B	107	VAL	N-CA-C	5.47	116.20	110.62
1	A	34	TRP	N-CA-CB	5.17	118.26	110.30
1	A	304	LEU	CA-C-N	-5.02	114.44	119.56
1	A	304	LEU	C-N-CA	-5.02	114.44	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3246	0	3111	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3468	0	3350	24	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	16	0	28	5	0
3	B	16	0	28	7	0
4	A	12	0	12	1	0
5	A	249	0	0	1	0
5	B	293	0	0	6	0
All	All	7302	0	6529	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:334:LEU:H	1:B:359:ASN:HD21	1.34	0.73
1:B:146:ARG:NE	3:B:502:MPD:H4	2.06	0.70
1:A:222:ARG:HG3	1:B:257:GLU:OE1	1.93	0.69
1:B:146:ARG:HE	3:B:502:MPD:H4	1.61	0.65
1:A:66:ASP:HB3	5:B:818:HOH:O	1.97	0.64
1:A:334:LEU:H	1:A:359:ASN:HD21	1.44	0.64
1:B:14:ARG:HD2	1:B:25:HIS:CE1	2.33	0.64
1:B:345:GLU:OE2	1:B:348:ARG:NH1	2.30	0.61
1:A:345:GLU:OE2	1:A:348:ARG:HD3	2.02	0.59
1:B:332:TYR:OH	1:B:388:ARG:NH1	2.36	0.58
1:B:221:PRO:HA	3:B:502:MPD:H53	1.85	0.58
1:A:146:ARG:NE	3:A:503:MPD:H4	2.19	0.56
1:A:146:ARG:HE	3:A:503:MPD:C4	2.18	0.56
1:B:219:TYR:HD2	3:B:502:MPD:HM3	1.72	0.54
1:A:13:GLN:HG2	5:A:848:HOH:O	2.07	0.54
1:A:146:ARG:HE	3:A:503:MPD:H4	1.74	0.53
1:B:139:HIS:HE1	5:B:706:HOH:O	1.91	0.53
1:A:65:PRO:HA	1:B:65:PRO:HA	1.91	0.52
3:B:501:MPD:O2	3:B:501:MPD:H52	2.09	0.52
1:B:139:HIS:HD2	1:B:245:ASN:OD1	1.92	0.51
1:B:321:GLN:NE2	5:B:602:HOH:O	2.43	0.51
1:B:33[A]:LEU:HD13	1:B:69:LEU:HD13	1.93	0.51
1:B:146:ARG:HE	3:B:502:MPD:C4	2.23	0.51
1:B:168:PRO:HG2	1:B:337:VAL:HG11	1.95	0.49
3:A:502:MPD:O2	3:A:502:MPD:H52	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:349:TYR:OH	4:A:504:GLC:H1	2.15	0.47
1:B:56:LEU:HD11	1:B:127:ARG:HD2	1.97	0.47
1:B:113:TYR:CE1	1:B:117:LYS:HG3	2.49	0.47
1:B:298[A]:LYS:NZ	5:B:865:HOH:O	2.42	0.47
1:B:329:LYS:NZ	5:B:603:HOH:O	2.47	0.46
1:A:221:PRO:HA	3:A:503:MPD:H53	1.96	0.46
1:A:172:LYS:HB3	1:A:173:PRO:HD3	1.97	0.46
3:B:501:MPD:O2	3:B:501:MPD:C5	2.64	0.46
1:A:15:ASN:HD21	1:A:25:HIS:HD2	1.63	0.45
1:B:129:ARG:NH2	5:B:604:HOH:O	2.48	0.45
1:A:113:TYR:CE1	1:A:117:LYS:HG3	2.54	0.42
1:B:60:ASP:HB3	1:B:88:TRP:CZ2	2.54	0.42
1:A:337:VAL:HG12	1:A:352:GLY:HA2	2.03	0.41
1:A:414:LEU:O	1:A:418:TRP:HB2	2.20	0.41
1:B:414:LEU:O	1:B:418:TRP:HB2	2.20	0.41
1:B:164:MET:HG2	1:B:350:TRP:CZ3	2.57	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/420 (92%)	378 (97%)	10 (3%)	0	100	100
1	B	417/420 (99%)	406 (97%)	11 (3%)	0	100	100
All	All	805/840 (96%)	784 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/349 (93%)	323 (100%)	1 (0%)	86	80
1	B	350/349 (100%)	348 (99%)	2 (1%)	78	70
All	All	674/698 (97%)	671 (100%)	3 (0%)	84	78

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	410	TRP
1	B	410	TRP
1	B	418	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	ASN
1	A	25	HIS
1	A	313	GLN
1	A	321	GLN
1	A	359	ASN
1	B	15	ASN
1	B	25	HIS
1	B	77	GLN
1	B	139	HIS
1	B	313	GLN
1	B	321	GLN
1	B	359	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GLC	A	504	-	12,12,12	0.84	0	17,17,17	0.90	2 (11%)
3	MPD	B	501	-	7,7,7	0.33	0	9,10,10	1.01	1 (11%)
3	MPD	B	502	-	7,7,7	0.31	0	9,10,10	0.98	0
3	MPD	A	502	-	7,7,7	0.28	0	9,10,10	0.78	0
3	MPD	A	503	-	7,7,7	0.40	0	9,10,10	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	A	504	-	-	0/2/22/22	0/1/1/1
3	MPD	B	501	-	-	3/5/5/5	-
3	MPD	B	502	-	-	3/5/5/5	-
3	MPD	A	502	-	-	4/5/5/5	-
3	MPD	A	503	-	-	3/5/5/5	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	A	504	GLC	O5-C5-C4	2.16	113.59	109.70
3	B	501	MPD	C5-C4-C3	2.16	121.70	111.67
4	A	504	GLC	O4-C4-C5	2.13	114.58	109.32

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	MPD	C2-C3-C4-C5
3	B	501	MPD	C2-C3-C4-C5
3	A	503	MPD	O2-C2-C3-C4
3	A	503	MPD	C1-C2-C3-C4
3	A	503	MPD	CM-C2-C3-C4
3	A	502	MPD	O2-C2-C3-C4
3	A	502	MPD	C2-C3-C4-O4
3	B	501	MPD	C2-C3-C4-O4
3	B	502	MPD	C2-C3-C4-O4
3	A	502	MPD	C1-C2-C3-C4
3	B	501	MPD	C1-C2-C3-C4
3	B	502	MPD	C1-C2-C3-C4
3	B	502	MPD	CM-C2-C3-C4

There are no ring outliers.

5 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	504	GLC	1	0
3	B	501	MPD	2	0
3	B	502	MPD	5	0
3	A	502	MPD	1	0
3	A	503	MPD	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	394/420 (93%)	0.37	22 (5%) 30 36	15, 23, 46, 75	0
1	B	417/420 (99%)	0.24	30 (7%) 21 26	9, 19, 47, 80	4 (0%)
All	All	811/840 (96%)	0.30	52 (6%) 25 30	9, 21, 47, 80	4 (0%)

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	85	PRO	7.3
1	A	85	PRO	4.3
1	A	179	VAL	4.1
1	B	92	ALA	4.1
1	B	2	ALA	4.0
1	B	88	TRP	4.0
1	B	89	GLY	3.9
1	A	94	ALA	3.9
1	B	84	GLY	3.8
1	A	212	TYR	3.8
1	A	349	TYR	3.8
1	B	94	ALA	3.5
1	B	83	PRO	3.4
1	A	92	ALA	3.3
1	A	89	GLY	3.3
1	B	182	LEU	3.3
1	A	84	GLY	3.2
1	B	179	VAL	3.2
1	B	212	TYR	3.1
1	B	226	ARG	3.1
1	B	28	TYR	3.1
1	B	181	ASN	3.0
1	A	204	TYR	2.9
1	B	329	LYS	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	195	PRO	2.9
1	B	340	THR	2.8
1	A	28	TYR	2.8
1	B	202	GLU	2.8
1	B	25	HIS	2.7
1	A	202	GLU	2.7
1	A	88	TRP	2.6
1	B	91	GLU	2.6
1	B	90	ARG	2.6
1	B	222	ARG	2.6
1	B	149	TYR	2.6
1	B	198	ARG	2.5
1	B	184	PRO	2.5
1	A	90	ARG	2.5
1	A	345	GLU	2.5
1	B	208	LEU	2.5
1	A	222	ARG	2.4
1	A	7	LYS	2.3
1	A	83	PRO	2.3
1	B	217	LEU	2.3
1	A	64	GLY	2.2
1	A	340	THR	2.2
1	B	200	ARG	2.2
1	A	208	LEU	2.1
1	B	219	TYR	2.1
1	B	196	GLU	2.1
1	A	205	ASP	2.0
1	A	178	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MPD	A	502	8/8	0.85	0.14	35,40,45,52	0
3	MPD	B	501	8/8	0.86	0.12	22,25,31,36	0
3	MPD	A	503	8/8	0.88	0.12	29,34,36,36	0
4	GLC	A	504	12/12	0.88	0.11	25,30,33,36	0
3	MPD	B	502	8/8	0.91	0.10	25,28,30,30	0
2	CL	A	501	1/1	0.99	0.05	19,19,19,19	1
2	CL	B	500	1/1	1.00	0.01	17,17,17,17	1

6.5 Other polymers [i](#)

There are no such residues in this entry.