



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 01:46 PM UTC

PDB ID : 5WVC / pdb_00005wvc
Title : Structure of the CARD-CARD disk
Authors : Lin, S.C.; Lo, Y.C.; Su, T.W.
Deposited on : 2016-12-24
Resolution : 2.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	95	0% 91% 9%
1	C	95	0% 87% 13%
1	E	95	0% 85% 14%
2	B	151	0% 55% 13% 32%
2	D	151	2% 61% 7% 32%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	151	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IOD	B	401	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4796 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Apoptotic protease-activating factor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	95	Total	C	N	O	S	0	0	0
			762	479	131	146	6			
1	A	95	Total	C	N	O	S	0	0	0
			762	479	131	146	6			
1	E	94	Total	C	N	O	S	0	0	0
			756	476	130	144	6			

- Molecule 2 is a protein called Caspase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	103	Total	C	N	O	S	0	0	0
			835	512	162	156	5			
2	B	103	Total	C	N	O	S	0	0	0
			835	512	162	156	5			
2	F	103	Total	C	N	O	S	0	0	0
			835	512	162	156	5			

There are 69 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	178	MET	-	expression tag	UNP A8K7U6
D	179	GLY	-	expression tag	UNP A8K7U6
D	180	SER	-	expression tag	UNP A8K7U6
D	181	SER	-	expression tag	UNP A8K7U6
D	182	HIS	-	expression tag	UNP A8K7U6
D	183	HIS	-	expression tag	UNP A8K7U6
D	184	HIS	-	expression tag	UNP A8K7U6
D	185	HIS	-	expression tag	UNP A8K7U6
D	186	HIS	-	expression tag	UNP A8K7U6
D	187	HIS	-	expression tag	UNP A8K7U6
D	188	SER	-	expression tag	UNP A8K7U6
D	189	SER	-	expression tag	UNP A8K7U6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	190	GLY	-	expression tag	UNP A8K7U6
D	191	LEU	-	expression tag	UNP A8K7U6
D	192	VAL	-	expression tag	UNP A8K7U6
D	193	PRO	-	expression tag	UNP A8K7U6
D	194	ARG	-	expression tag	UNP A8K7U6
D	195	GLY	-	expression tag	UNP A8K7U6
D	196	SER	-	expression tag	UNP A8K7U6
D	197	HIS	-	expression tag	UNP A8K7U6
D	198	MET	-	expression tag	UNP A8K7U6
D	199	ALA	-	expression tag	UNP A8K7U6
D	200	SER	-	expression tag	UNP A8K7U6
B	178	MET	-	expression tag	UNP A8K7U6
B	179	GLY	-	expression tag	UNP A8K7U6
B	180	SER	-	expression tag	UNP A8K7U6
B	181	SER	-	expression tag	UNP A8K7U6
B	182	HIS	-	expression tag	UNP A8K7U6
B	183	HIS	-	expression tag	UNP A8K7U6
B	184	HIS	-	expression tag	UNP A8K7U6
B	185	HIS	-	expression tag	UNP A8K7U6
B	186	HIS	-	expression tag	UNP A8K7U6
B	187	HIS	-	expression tag	UNP A8K7U6
B	188	SER	-	expression tag	UNP A8K7U6
B	189	SER	-	expression tag	UNP A8K7U6
B	190	GLY	-	expression tag	UNP A8K7U6
B	191	LEU	-	expression tag	UNP A8K7U6
B	192	VAL	-	expression tag	UNP A8K7U6
B	193	PRO	-	expression tag	UNP A8K7U6
B	194	ARG	-	expression tag	UNP A8K7U6
B	195	GLY	-	expression tag	UNP A8K7U6
B	196	SER	-	expression tag	UNP A8K7U6
B	197	HIS	-	expression tag	UNP A8K7U6
B	198	MET	-	expression tag	UNP A8K7U6
B	199	ALA	-	expression tag	UNP A8K7U6
B	200	SER	-	expression tag	UNP A8K7U6
F	178	MET	-	expression tag	UNP A8K7U6
F	179	GLY	-	expression tag	UNP A8K7U6
F	180	SER	-	expression tag	UNP A8K7U6
F	181	SER	-	expression tag	UNP A8K7U6
F	182	HIS	-	expression tag	UNP A8K7U6
F	183	HIS	-	expression tag	UNP A8K7U6
F	184	HIS	-	expression tag	UNP A8K7U6
F	185	HIS	-	expression tag	UNP A8K7U6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	186	HIS	-	expression tag	UNP A8K7U6
F	187	HIS	-	expression tag	UNP A8K7U6
F	188	SER	-	expression tag	UNP A8K7U6
F	189	SER	-	expression tag	UNP A8K7U6
F	190	GLY	-	expression tag	UNP A8K7U6
F	191	LEU	-	expression tag	UNP A8K7U6
F	192	VAL	-	expression tag	UNP A8K7U6
F	193	PRO	-	expression tag	UNP A8K7U6
F	194	ARG	-	expression tag	UNP A8K7U6
F	195	GLY	-	expression tag	UNP A8K7U6
F	196	SER	-	expression tag	UNP A8K7U6
F	197	HIS	-	expression tag	UNP A8K7U6
F	198	MET	-	expression tag	UNP A8K7U6
F	199	ALA	-	expression tag	UNP A8K7U6
F	200	SER	-	expression tag	UNP A8K7U6


- Molecule 3 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	2	Total I 2 2	0	0
3	D	4	Total I 4 4	0	0
3	B	2	Total I 2 2	0	0
3	E	1	Total I 1 1	0	0
3	F	2	Total I 2 2	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Apoptotic protease-activating factor 1




- Molecule 1: Apoptotic protease-activating factor 1



- Molecule 1: Apoptotic protease-activating factor 1



- Molecule 2: Caspase



- Molecule 2: Caspase



LEU
GLU
ASN
LEU
THR
PRO
VAL
VAL
LEU
ARG
PRO
GLU
ILE
ARG
PRO
GLU
VAL
LEU
ARG
PRO
GLU
THR
PRO
ARG
PRO

● Molecule 2: Caspase



MET
GLY
SER
SER
HIS
HIS
HIS
HIS
HIS
HIS
SER
SER
GLY
LEU
VAL
PRO
ARG
GLY
SER
HIS
MET
A199
S200
M201
R206
V217
D223
Q224
L225
L229
E233
R236
M239
I243
R252
D253
R256
R265
L272
S275
K300
P301
THR
LEU
GLU
ASN

LEU
THR
PRO
VAL
LEU
ARG
PRO
GLU
ILE
ARG
LYS
PRO
GLU
VAL
LEU
ARG
PRO
GLU
THR
PRO
ARG
PRO

4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	190.11Å 190.11Å 68.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.85 – 2.99 24.85 – 2.99	Depositor EDS
% Data completeness (in resolution range)	97.6 (24.85-2.99) 97.4 (24.85-2.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.99Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.197 , 0.228 0.202 , 0.231	Depositor DCC
R_{free} test set	2006 reflections (6.93%)	wwPDB-VP
Wilson B-factor (Å ²)	66.7	Xtriage
Anisotropy	0.546	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 26.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.065 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4796	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IOD, CSO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/773	0.74	0/1038
1	C	0.28	0/773	0.70	0/1038
1	E	0.28	0/767	0.72	0/1030
2	B	0.27	0/835	0.74	0/1118
2	D	0.27	0/835	0.67	0/1118
2	F	0.27	0/835	0.71	0/1118
All	All	0.27	0/4818	0.71	0/6460

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	762	0	768	5	0
1	C	762	0	768	7	0
1	E	756	0	763	5	0
2	B	835	0	852	13	0
2	D	835	0	851	7	0
2	F	835	0	851	13	0
3	B	2	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	2	0	0	0	0
3	D	4	0	0	1	0
3	E	1	0	0	0	0
3	F	2	0	0	0	0
All	All	4796	0	4853	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 49 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MET:HE1	1:A:9:LEU:HD12	1.71	0.72
1:E:4:LYS:HA	1:E:7:ASN:HB2	1.71	0.72
2:B:300:LYS:HG2	2:B:301:PRO:HA	1.72	0.71
2:D:213:ARG:HD3	2:D:259:ILE:HD11	1.73	0.71
1:C:26:MET:HE2	1:C:35:LEU:HD21	1.77	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	93/95 (98%)	89 (96%)	4 (4%)	0	100	100
1	C	93/95 (98%)	93 (100%)	0	0	100	100
1	E	92/95 (97%)	83 (90%)	8 (9%)	1 (1%)	11	43
2	B	100/151 (66%)	97 (97%)	3 (3%)	0	100	100
2	D	100/151 (66%)	99 (99%)	1 (1%)	0	100	100
2	F	100/151 (66%)	97 (97%)	3 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	578/738 (78%)	558 (96%)	19 (3%)	1 (0%)	43 76

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	3	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	84/84 (100%)	84 (100%)	0	100 100
1	C	84/84 (100%)	83 (99%)	1 (1%)	63 82
1	E	83/84 (99%)	81 (98%)	2 (2%)	43 73
2	B	90/135 (67%)	90 (100%)	0	100 100
2	D	90/135 (67%)	90 (100%)	0	100 100
2	F	90/135 (67%)	88 (98%)	2 (2%)	45 74
All	All	521/657 (79%)	516 (99%)	5 (1%)	68 84

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	94	VAL
1	E	87	LEU
1	E	94	VAL
2	F	233	GLU
2	F	300	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	88	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	11	GLN
2	F	294	GLN
2	D	292	ASN
1	C	73	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CSO	F	212	2	3,6,7	0.63	0	1,6,8	0.10	0
2	CSO	D	212	2	3,6,7	0.63	0	1,6,8	0.33	0
2	CSO	B	212	2	3,6,7	0.64	0	1,6,8	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSO	F	212	2	-	0/1/5/7	-
2	CSO	D	212	2	-	0/1/5/7	-
2	CSO	B	212	2	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	95/95 (100%)	-0.01	1 (1%) 78 57	54, 80, 109, 124	0
1	C	95/95 (100%)	-0.21	1 (1%) 78 57	45, 60, 80, 122	0
1	E	94/95 (98%)	0.13	1 (1%) 78 57	60, 86, 123, 152	0
2	B	102/151 (67%)	-0.10	1 (0%) 79 59	44, 66, 95, 157	0
2	D	102/151 (67%)	0.04	3 (2%) 53 31	49, 65, 91, 152	0
2	F	102/151 (67%)	0.05	3 (2%) 53 31	56, 82, 106, 130	0
All	All	590/738 (79%)	-0.02	10 (1%) 69 45	44, 72, 110, 157	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	199	ALA	3.9
1	E	94	VAL	3.5
2	D	199	ALA	3.4
2	D	301	PRO	3.3
2	F	301	PRO	3.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CSO	B	212	7/8	0.87	0.12	58,62,82,118	0
2	CSO	F	212	7/8	0.90	0.12	53,55,80,133	0
2	CSO	D	212	7/8	0.91	0.11	53,56,64,102	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	IOD	C	102	1/1	0.74	0.23	186,186,186,186	1
3	IOD	B	402	1/1	0.85	0.17	178,178,178,178	1
3	IOD	D	404	1/1	0.92	0.08	150,150,150,150	1
3	IOD	D	403	1/1	0.94	0.20	123,123,123,123	1
3	IOD	E	101	1/1	0.94	0.07	135,135,135,135	1
3	IOD	C	101	1/1	0.95	0.11	93,93,93,93	1
3	IOD	D	402	1/1	0.96	0.10	105,105,105,105	1
3	IOD	B	401	1/1	0.97	0.05	78,78,78,78	1
3	IOD	F	1202	1/1	0.97	0.04	83,83,83,83	1
3	IOD	D	401	1/1	0.98	0.03	63,63,63,63	1
3	IOD	F	1201	1/1	0.99	0.02	63,63,63,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.