



wwPDB EM Validation Summary Report ⓘ

Mar 29, 2026 – 10:15 AM UTC

PDB ID : 2WWB / pdb_00002wwb
EMDB ID : EMD-1652
Title : CRYO-EM STRUCTURE OF THE MAMMALIAN SEC61 COMPLEX
BOUND TO THE ACTIVELY TRANSLATING WHEAT GERM 80S RI-
BOSOME
Authors : Becker, T.; Mandon, E.; Bhushan, S.; Jarasch, A.; Armache, J.P.; Funes,
S.; Jossinet, F.; Gumbart, J.; Mielke, T.; Berninghausen, O.; Schulten, K.;
Westhof, E.; Gilmore, R.; Beckmann, R.
Deposited on : 2009-10-22
Resolution : 6.48 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 6.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	496 (5.98 - 6.95)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	
2	B	68	
3	C	96	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	63	
5	E	34	
6	F	25	
7	G	14	
8	H	362	
9	I	184	
10	J	189	
11	K	142	
12	L	127	
13	M	113	
14	N	120	
15	O	51	

2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 14313 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN TRANSPORT PROTEIN SEC61 SUBUNIT ALPHA ISOFORM 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	476	Total	C	N	O	S	0	0
			3675	2409	591	650	25		

- Molecule 2 is a protein called PROTEIN TRANSPORT PROTEIN SEC61 SUBUNIT GAMMA.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	68	Total	C	N	O	S	0	0
			543	355	94	89	5		

- Molecule 3 is a protein called PROTEIN TRANSPORT PROTEIN SEC61 SUBUNIT BETA.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	36	Total	C	N	O	S	0	0
			281	188	44	47	2		

- Molecule 4 is a RNA chain called 5.8S RRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	63	Total	C	N	O	P	0	0
			1347	603	245	436	63		

- Molecule 5 is a RNA chain called 25S RRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	34	Total	C	N	O	P	0	0
			740	332	148	226	34		

- Molecule 6 is a RNA chain called 25S RRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	25	Total	C	N	O	P	0	0
			536	239	99	173	25		

- Molecule 7 is a RNA chain called 25S RRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	14	Total	C	N	O	P	0	0
			296	133	54	96	13		

- Molecule 8 is a protein called 60S RIBOSOMAL PROTEIN L4-B.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	269	Total	C	N	O	S	0	0
			2039	1281	391	363	4		

- Molecule 9 is a protein called 60S RIBOSOMAL PROTEIN L17-A.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	I	153	Total	C	N	O	S	0	0
			1212	756	236	219	1		

- Molecule 10 is a protein called 60S RIBOSOMAL PROTEIN L19.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	J	53	Total	C	N	O	S	0	0
			410	254	83	72	1		

- Molecule 11 is a protein called 60S RIBOSOMAL PROTEIN L25.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	K	83	Total	C	N	O	S	0	0
			663	424	111	126	2		

- Molecule 12 is a protein called 60S RIBOSOMAL PROTEIN L26-A.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	L	127	Total	C	N	O	S	0	0
			1002	630	193	178	1		

- Molecule 13 is a protein called 60S RIBOSOMAL PROTEIN L31-A.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	M	84	Total	C	N	O	S	0	0
			706	447	140	118	1		

- Molecule 14 is a protein called 60S RIBOSOMAL PROTEIN L35.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	N	69	Total	C	N	O	S	0	0
			547	345	101	99	2		


- Molecule 15 is a protein called 60S RIBOSOMAL PROTEIN L39.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	O	37	Total	C	N	O	S	0	0
			316	200	66	48	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTEIN TRANSPORT PROTEIN SEC61 SUBUNIT ALPHA ISOFORM 1




- Molecule 2: PROTEIN TRANSPORT PROTEIN SEC61 SUBUNIT GAMMA




- Molecule 3: PROTEIN TRANSPORT PROTEIN SEC61 SUBUNIT BETA






• Molecule 4: 5.8S rRNA



• Molecule 5: 25S rRNA






• Molecule 6: 25S rRNA




• Molecule 7: 25S rRNA




• Molecule 8: 60S RIBOSOMAL PROTEIN L4-B




• Molecule 13: 60S RIBOSOMAL PROTEIN L31-A



• Molecule 14: 60S RIBOSOMAL PROTEIN L35



• Molecule 15: 60S RIBOSOMAL PROTEIN L39



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	221445	Depositor
Resolution determination method	Not provided	
CTF correction method	DEFOCUS GROUP VOLUMES	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	25	Depositor
Minimum defocus (nm)	1250	Depositor
Maximum defocus (nm)	4700	Depositor
Magnification	38000	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	1.706	Depositor
Minimum map value	-0.671	Depositor
Average map value	0.005	Depositor
Map value standard deviation	0.100	Depositor
Recommended contour level	0.2	Depositor
Map size ($\text{\AA}$)	455.4, 455.4, 455.4	wwPDB
Map dimensions	368, 368, 368	wwPDB
Map angles ($^\circ$)	90, 90, 90	wwPDB
Pixel spacing ($\text{\AA}$)	1.2375, 1.2375, 1.2375	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.46	8/3755 (0.2%)	1.70	49/5089 (1.0%)
2	B	1.19	0/553	1.72	9/738 (1.2%)
3	C	1.10	0/289	1.62	3/391 (0.8%)
4	D	0.92	0/1508	1.30	12/2348 (0.5%)
5	E	0.91	0/833	1.16	3/1298 (0.2%)
6	F	0.91	0/599	1.07	1/932 (0.1%)
7	G	0.95	0/330	1.21	3/513 (0.6%)
8	H	1.27	0/2079	2.03	82/2817 (2.9%)
9	I	1.30	0/1235	2.00	50/1662 (3.0%)
10	J	1.28	0/412	1.46	0/551
11	K	1.15	0/670	1.66	9/903 (1.0%)
12	L	1.28	0/1013	1.81	26/1351 (1.9%)
13	M	1.37	0/719	1.79	19/959 (2.0%)
14	N	1.32	1/549 (0.2%)	2.05	24/733 (3.3%)
15	O	1.34	0/321	1.86	6/426 (1.4%)
All	All	1.60	9/14865 (0.1%)	1.69	296/20711 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
3	C	0	1
4	D	0	10
5	E	0	6
6	F	0	3
7	G	0	1
8	H	0	12
9	I	0	6
11	K	0	3
12	L	0	5

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
13	M	0	2
14	N	0	2
15	O	0	7
All	All	0	63

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	104	ASP	N-CA	126.72	3.09	1.46
1	A	5	PHE	CG-CD2	20.78	1.82	1.38
1	A	5	PHE	CG-CD1	20.48	1.81	1.38
1	A	5	PHE	CD2-CE2	14.34	1.81	1.38
1	A	5	PHE	CD1-CE1	13.99	1.80	1.38

The worst 5 of 296 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	532	A	P-O3'-C3'	13.96	141.14	120.20
1	A	103	GLY	CA-C-N	12.76	145.91	121.54
1	A	103	GLY	C-N-CA	12.76	145.91	121.54
9	I	56	ARG	CA-C-N	12.75	144.64	121.70
9	I	56	ARG	C-N-CA	12.75	144.64	121.70

There are no chirality outliers.

5 of 63 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	112	PHE	Sidechain
1	A	270	ALA	Peptide
1	A	271	ARG	Sidechain
1	A	336	TYR	Peptide
1	A	404	HIS	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3675	0	3798	45	0
2	B	543	0	577	1	0
3	C	281	0	294	0	0
4	D	1347	0	678	2	0
5	E	740	0	369	1	0
6	F	536	0	272	1	0
7	G	296	0	154	0	0
8	H	2039	0	2106	16	0
9	I	1212	0	1231	9	0
10	J	410	0	452	0	0
11	K	663	0	699	5	0
12	L	1002	0	1093	2	0
13	M	706	0	741	3	0
14	N	547	0	613	8	0
15	O	316	0	349	2	0
All	All	14313	0	13426	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:PHE:CD2	1:A:5:PHE:CE2	1.81	1.68
1:A:5:PHE:CD1	1:A:5:PHE:CE1	1.80	1.66
1:A:5:PHE:CE2	1:A:5:PHE:CZ	1.79	1.64
1:A:5:PHE:CD1	1:A:5:PHE:CG	1.81	1.61
1:A:5:PHE:CE1	1:A:5:PHE:CZ	1.80	1.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/476 (100%)	402 (85%)	46 (10%)	26 (6%)	1	14
2	B	66/68 (97%)	60 (91%)	2 (3%)	4 (6%)	1	12
3	C	34/96 (35%)	27 (79%)	4 (12%)	3 (9%)	0	8
8	H	267/362 (74%)	181 (68%)	38 (14%)	48 (18%)	0	2
9	I	151/184 (82%)	119 (79%)	15 (10%)	17 (11%)	0	5
10	J	51/189 (27%)	46 (90%)	4 (8%)	1 (2%)	6	31
11	K	81/142 (57%)	73 (90%)	5 (6%)	3 (4%)	2	19
12	L	125/127 (98%)	101 (81%)	9 (7%)	15 (12%)	0	4
13	M	82/113 (73%)	65 (79%)	8 (10%)	9 (11%)	0	5
14	N	67/120 (56%)	59 (88%)	3 (4%)	5 (8%)	1	10
15	O	35/51 (69%)	26 (74%)	0	9 (26%)	0	1
All	All	1433/1928 (74%)	1159 (81%)	134 (9%)	140 (10%)	1	7

5 of 140 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	56	SER
1	A	104	ASP
1	A	108	ASP
1	A	110	ALA
1	A	142	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	398/398 (100%)	389 (98%)	9 (2%)	44	64
2	B	59/59 (100%)	56 (95%)	3 (5%)	21	42
3	C	32/74 (43%)	31 (97%)	1 (3%)	35	56
8	H	209/288 (73%)	186 (89%)	23 (11%)	6	20
9	I	124/146 (85%)	105 (85%)	19 (15%)	3	12

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
10	J	44/154 (29%)	41 (93%)	3 (7%)	14	36
11	K	74/118 (63%)	63 (85%)	11 (15%)	3	12
12	L	110/110 (100%)	96 (87%)	14 (13%)	4	16
13	M	74/97 (76%)	67 (90%)	7 (10%)	8	25
14	N	62/105 (59%)	56 (90%)	6 (10%)	8	24
15	O	33/46 (72%)	28 (85%)	5 (15%)	3	12
All	All	1219/1595 (76%)	1118 (92%)	101 (8%)	12	30

5 of 101 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
10	J	53	LYS
12	L	52	ARG
15	O	23	LEU
11	K	67	ILE
11	K	117	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
12	L	98	ASN
13	M	15	ASN
1	A	360	HIS
1	A	343	HIS
14	N	20	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	D	62/63 (98%)	28 (45%)	10 (16%)
5	E	33/34 (97%)	9 (27%)	3 (9%)
6	F	24/25 (96%)	3 (12%)	0
7	G	14/14 (100%)	6 (42%)	2 (14%)
All	All	133/136 (97%)	46 (34%)	15 (11%)

5 of 46 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
4	D	48	A
4	D	49	G
4	D	51	G
4	D	52	A
4	D	56	G

5 of 15 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
4	D	90	U
7	G	1917	A
4	D	93	U
7	G	1926	G
5	E	558	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-1652. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 184



Y Index: 184




Z Index: 184


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 214



Y Index: 191




Z Index: 168


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.2. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 3348 nm³; this corresponds to an approximate mass of 3024 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.154 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-1652 and PDB model 2WWB. Per-residue inclusion information can be found in section [3](#) on page [7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.2 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.2).

9.4 Atom inclusion [i](#)



At the recommended contour level, 79% of all backbone atoms, 70% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.2) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6970	 0.0920
A	 0.5780	 0.0860
B	 0.6690	 0.1140
C	 0.3970	 0.0410
D	 0.8190	 0.0880
E	 0.9350	 0.1460
F	 0.8770	 0.1340
G	 0.9430	 0.1730
H	 0.6630	 0.0690
I	 0.6890	 0.0790
J	 0.6670	 0.1180
K	 0.6500	 0.0970
L	 0.8560	 0.0970
M	 0.7070	 0.0760
N	 0.7430	 0.0920
O	 0.5690	 0.0980

