



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 09:58 AM UTC

PDB ID : 4WW2 / pdb_00004ww2
Title : Crystal structure of human TCR Alpha Chain-TRAV21-TRAJ8, Beta Chain-TRBV7-8, Antigen-presenting glycoprotein CD1d, and Beta-2-microglobulin
Authors : Le Nours, J.; Praveena, T.; Pellicci, D.; Gherardin, N.A.; Lim, R.T.; Besra, G.; Keshipeddy, S.; Richardson, S.K.; Howell, A.R.; Gras, S.; Godfrey, D.I.; Rossjohn, J.; Uldrich, A.P.
Deposited on : 2014-11-10
Resolution : 2.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	6% 86% 12%
2	B	243	2% 92% 8%
3	C	284	4% 81% 15%
4	F	99	38% 93%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6540 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TCR Alpha Chain-TRAV21-TRAJ8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	0	1	0
			1501	943	249	300	9			

- Molecule 2 is a protein called TCR Beta Chain-TRBV7-8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	243	Total	C	N	O	S	4	3	0
			1936	1219	339	372	6			

- Molecule 3 is a protein called Antigen-presenting glycoprotein CD1d.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	275	Total	C	N	O	S	0	0	0
			2162	1384	378	393	7			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	SER	-	expression tag	UNP P15813
C	1	PRO	-	expression tag	UNP P15813
C	2	GLY	-	expression tag	UNP P15813
C	278	HIS	-	expression tag	UNP P15813
C	279	HIS	-	expression tag	UNP P15813
C	280	HIS	-	expression tag	UNP P15813
C	281	HIS	-	expression tag	UNP P15813
C	282	HIS	-	expression tag	UNP P15813
C	283	HIS	-	expression tag	UNP P15813

- Molecule 4 is a protein called Beta-2-microglobulin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	96	Total	C	N	O	S	0	0	0
			685	440	116	127	2			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is (15Z)-N-[(2S,3S,4R)-1-(alpha-D-galactopyranosyloxy)-3,4-dihydroxyoctadecan-2-yl]tetracos-15-enamide (CCD ID: JLS) (formula: $C_{48}H_{93}NO_9$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	C	1	Total	C	N	O	0	0
			58	48	1	9		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	50	Total	O	0	0
			50	50		
7	B	65	Total	O	0	0
			65	65		
7	C	48	Total	O	0	0
			48	48		
7	F	7	Total	O	0	0
			7	7		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TCR Alpha Chain-TRAV21-TRAJ8




- Molecule 2: TCR Beta Chain-TRBV7-8



- Molecule 3: Antigen-presenting glycoprotein CD1d



- Molecule 4: Beta-2-microglobulin



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	209.99Å 46.30Å 124.65Å 90.00° 122.28° 90.00°	Depositor
Resolution (Å)	44.80 – 2.48 44.80 – 2.48	Depositor EDS
% Data completeness (in resolution range)	99.3 (44.80-2.48) 99.4 (44.80-2.48)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 2.48Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.196 , 0.244 (Not available) , 0.275	Depositor DCC
R_{free} test set	1816 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	41.6	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 78.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6540	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: JLS, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	0/1532	1.16	7/2086 (0.3%)
2	B	0.73	0/1990	1.12	4/2708 (0.1%)
3	C	0.72	0/2228	1.16	5/3042 (0.2%)
4	F	0.74	0/708	1.13	2/973 (0.2%)
All	All	0.75	0/6458	1.14	18/8809 (0.2%)

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	20	ASN	CA-CB-CG	6.78	119.38	112.60
2	B	189	GLU	CA-C-N	6.13	128.62	122.00
2	B	189	GLU	C-N-CA	6.13	128.62	122.00
4	F	32	PRO	CA-C-N	5.97	128.28	120.28
4	F	32	PRO	C-N-CA	5.97	128.28	120.28
3	C	272	ASP	CA-CB-CG	5.53	118.13	112.60
2	B	73	SER	CA-C-N	5.42	128.08	120.28
2	B	73	SER	C-N-CA	5.42	128.08	120.28
1	A	108	ASN	CA-C-N	5.41	128.93	120.82
1	A	108	ASN	C-N-CA	5.41	128.93	120.82
3	C	66	LEU	CA-C-N	5.35	127.71	120.38
3	C	66	LEU	C-N-CA	5.35	127.71	120.38
1	A	140	ASP	CA-CB-CG	5.29	117.89	112.60
1	A	178	MET	CA-C-N	5.26	127.59	120.38
1	A	178	MET	C-N-CA	5.26	127.59	120.38
3	C	234	ASP	CA-CB-CG	5.26	117.86	112.60
1	A	182	ASP	CA-C-N	5.19	128.48	121.05
1	A	182	ASP	C-N-CA	5.19	128.48	121.05

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1501	0	1418	11	0
2	B	1936	0	1839	8	0
3	C	2162	0	2056	19	0
4	F	685	0	531	2	0
5	C	28	0	26	0	0
6	C	58	0	93	2	0
7	A	50	0	0	0	0
7	B	65	0	0	0	0
7	C	48	0	0	0	0
7	F	7	0	0	0	0
All	All	6540	0	5963	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:96:LEU:HD21	3:C:118:PHE:HD1	1.55	0.71
3:C:235:ILE:HG12	3:C:245:LEU:HD12	1.74	0.70
3:C:217:TRP:HB3	3:C:264:LYS:HB2	1.74	0.70
3:C:154:THR:HG23	6:C:303:JLS:H2	1.84	0.59
3:C:187:PRO:HB3	3:C:211:PHE:HB3	1.86	0.57
3:C:66:LEU:HD21	3:C:165:THR:HG21	1.86	0.56
3:C:206:CYS:HG	3:C:261:CYS:HG	0.57	0.56
1:A:56:ILE:HD11	1:A:78:ALA:HB3	1.87	0.56
2:B:163:ASP:OD1	2:B:186:PRO:HG2	2.06	0.56
1:A:40:GLN:HG2	1:A:55:LEU:HD12	1.88	0.55
3:C:102:CYS:SG	3:C:166:CYS:SG	3.06	0.53
3:C:29:LEU:HD11	3:C:31:TRP:NE1	2.24	0.52
4:F:54:LEU:HA	4:F:64:LEU:CD2	2.40	0.51
1:A:39:LEU:C	1:A:39:LEU:HD23	2.35	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:18:PHE:HB2	3:C:94:LEU:HB2	1.93	0.51
2:B:68:LYS:HB3	2:B:71:LEU:HD12	1.93	0.51
4:F:54:LEU:HA	4:F:64:LEU:HD21	1.93	0.50
3:C:29:LEU:C	3:C:29:LEU:HD12	2.37	0.50
2:B:227:LEU:HD22	2:B:240:PRO:HD2	1.94	0.49
2:B:21:LEU:HD22	2:B:119:THR:HG21	1.95	0.49
3:C:129:THR:O	3:C:155:ARG:HD2	2.15	0.47
1:A:19:LEU:C	1:A:19:LEU:HD12	2.40	0.47
2:B:12:VAL:HB	2:B:162:PRO:HG3	1.97	0.46
2:B:16:GLY:O	2:B:93:ARG:HA	2.16	0.45
1:A:21:LEU:HD22	1:A:120:THR:HG21	1.99	0.45
1:A:56:ILE:HD11	1:A:78:ALA:CB	2.46	0.45
3:C:204:LEU:O	3:C:248:THR:HA	2.15	0.45
3:C:14:GLN:HB3	3:C:98:VAL:HB	1.99	0.44
3:C:21:SER:O	3:C:82:LYS:HE3	2.17	0.44
1:A:43:ARG:HB2	1:A:53:LEU:HD11	1.99	0.43
1:A:46:PRO:HA	1:A:47:GLY:HA2	1.68	0.43
3:C:12:CYS:HB3	6:C:303:JLS:H26	2.01	0.43
1:A:13:VAL:HG22	1:A:14:PRO:HD2	2.01	0.42
3:C:204:LEU:HB3	3:C:220:TRP:CZ2	2.54	0.42
2:B:186:PRO:HG3	2:B:200:LEU:HD13	2.02	0.41
3:C:179:SER:HA	3:C:182:LYS:HE3	2.03	0.41
2:B:233:TRP:CZ2	2:B:240:PRO:HD3	2.56	0.41
1:A:126:PRO:HG3	1:A:175:VAL:HG13	2.02	0.41
3:C:25:ARG:HD3	3:C:27:ASP:OD1	2.21	0.41
1:A:39:LEU:HD13	1:A:87:SER:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	202/207 (98%)	195 (96%)	7 (4%)	0	100	100
2	B	244/243 (100%)	234 (96%)	10 (4%)	0	100	100
3	C	273/284 (96%)	265 (97%)	8 (3%)	0	100	100
4	F	94/99 (95%)	91 (97%)	3 (3%)	0	100	100
All	All	813/833 (98%)	785 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/183 (89%)	160 (98%)	3 (2%)	51	74
2	B	208/209 (100%)	204 (98%)	4 (2%)	50	73
3	C	228/249 (92%)	217 (95%)	11 (5%)	23	43
4	F	56/94 (60%)	55 (98%)	1 (2%)	51	74
All	All	655/735 (89%)	636 (97%)	19 (3%)	37	62

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	MET
1	A	57	GLN
1	A	179	ARG
2	B	15	ARG
2	B	51	GLU
2	B	185	GLN
2	B	203	ARG
3	C	20	ASN
3	C	45	ASP
3	C	48	ARG
3	C	50	LEU
3	C	61	GLN
3	C	95	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	104	VAL
3	C	150	GLN
3	C	192	SER
3	C	204	LEU
3	C	207	HIS
4	F	80	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	ASN
2	B	95	GLN
2	B	113	GLN
3	C	14	GLN
3	C	184	GLN
4	F	2	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	C	301	3	14,14,15	0.27	0	17,19,21	0.45	0
5	NAG	C	302	3	14,14,15	0.31	0	17,19,21	0.50	0
6	JLS	C	303	-	58,58,58	0.41	0	64,67,67	0.71	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	301	3	-	1/6/23/26	0/1/1/1
5	NAG	C	302	3	-	1/6/23/26	0/1/1/1
6	JLS	C	303	-	-	22/56/76/76	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	303	JLS	C1A-O5A-C5M	2.04	117.71	113.72

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	303	JLS	CAA-CAB-CAC-CAD
6	C	303	JLS	C9-C10-C11-C12
6	C	303	JLS	CAJ-CAK-CAL-CAM
6	C	303	JLS	CAH-CAI-CAJ-CAK
6	C	303	JLS	CAR-CAS-CAT-CAU
6	C	303	JLS	C11-C10-C9-C8
6	C	303	JLS	CAF-CAG-CAH-CAI
6	C	303	JLS	CAK-CAL-CAM-CAN
5	C	301	NAG	O5-C5-C6-O6
6	C	303	JLS	CAT-CAU-CAV-CAW
6	C	303	JLS	C13-C14-C15-C16
5	C	302	NAG	O5-C5-C6-O6
6	C	303	JLS	CAL-CAM-CAN-CAO
6	C	303	JLS	C11-C12-C13-C14
6	C	303	JLS	CAU-CAV-CAW-CAX
6	C	303	JLS	CAG-CAH-CAI-CAJ
6	C	303	JLS	CAP-CAQ-CAR-CAS

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	C	303	JLS	O1A-C1-C2-C3
6	C	303	JLS	CAQ-CAR-CAS-CAT
6	C	303	JLS	C12-C13-C14-C15
6	C	303	JLS	C14-C15-C16-C17
6	C	303	JLS	CAM-CAN-CAO-CAP
6	C	303	JLS	CAB-CAC-CAD-CAE
6	C	303	JLS	C6-C7-C8-C9

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	303	JLS	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/207 (98%)	0.36	12 (5%) 28 25	19, 48, 94, 116	2 (0%)
2	B	243/243 (100%)	-0.03	6 (2%) 58 55	18, 41, 68, 98	4 (1%)
3	C	275/284 (96%)	0.43	10 (3%) 46 42	36, 57, 90, 110	1 (0%)
4	F	96/99 (96%)	2.50	38 (39%) 1 0	20, 92, 120, 128	6 (6%)
All	All	817/833 (98%)	0.52	66 (8%) 18 15	18, 52, 100, 128	13 (1%)

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	F	17	ASN	25.8
4	F	16	GLU	19.1
4	F	73	THR	17.8
4	F	18	GLY	17.7
4	F	42	ASN	12.6
4	F	41	LYS	7.9
4	F	97	ARG	4.1
1	A	176	LEU	3.8
2	B	1	GLY	3.7
2	B	0	MET	3.6
4	F	78	TYR	3.6
4	F	35	ILE	3.6
3	C	277	TRP	3.5
4	F	34	ASP	3.4
4	F	77	GLU	3.3
4	F	80	CYS	3.3
4	F	19	LYS	3.3
1	A	216	SER	3.2
4	F	72	PRO	3.2
1	A	215	PRO	3.1
3	C	202	LEU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	2	ALA	3.1
4	F	43	GLY	3.0
2	B	195	ASP	3.0
4	F	83	ASN	3.0
3	C	193	ARG	2.9
3	C	253	ALA	2.9
4	F	36	GLU	2.9
4	F	76	ASP	2.9
4	F	82	VAL	2.9
4	F	65	LEU	2.9
1	A	181	MET	2.9
2	B	193	LEU	2.8
4	F	90	PRO	2.7
4	F	24	ASN	2.7
4	F	67	TYR	2.6
4	F	74	GLU	2.5
4	F	49	VAL	2.5
4	F	38	ASP	2.5
4	F	23	LEU	2.5
4	F	85	VAL	2.5
1	A	209	PRO	2.4
4	F	15	ALA	2.4
1	A	213	PHE	2.4
1	A	164	LYS	2.4
4	F	20	SER	2.4
4	F	46	ILE	2.4
4	F	66	TYR	2.4
3	C	3	VAL	2.3
1	A	144	SER	2.2
3	C	78	THR	2.2
4	F	71	THR	2.2
1	A	145	ASP	2.2
2	B	254	ASP	2.2
1	A	162	GLN	2.2
3	C	194	GLY	2.1
4	F	64	LEU	2.1
3	C	200	GLY	2.1
4	F	68	THR	2.1
1	A	210	GLU	2.1
4	F	48	LYS	2.1
3	C	239	ALA	2.1
4	F	37	VAL	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	F	45	ARG	2.0
1	A	194	LYS	2.0
3	C	136	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	C	301	14/15	0.73	0.16	114,120,123,124	0
5	NAG	C	302	14/15	0.77	0.19	81,86,90,93	0
6	JLS	C	303	58/58	0.95	0.10	36,50,61,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around JLS C 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.