



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:34 PM UTC

PDB ID : 4WW4 / pdb_00004ww4
Title : Double-heterohexameric rings of full-length Rvb1(ADP)/Rvb2(ADP)
Authors : Lakomek, K.; Hopfner, K.-P.
Deposited on : 2014-11-10
Resolution : 2.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1159 (2.96-2.92)
Clashscore	190562	1184 (2.96-2.92)
Ramachandran outliers	187476	1131 (2.96-2.92)
Sidechain outliers	187428	1131 (2.96-2.92)
RSRZ outliers	180081	1159 (2.96-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	462	
2	B	513	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6603 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RuvB-like 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	416	Total	C	N	O	S	0	0	0
			3190	1999	573	604	14			

- Molecule 2 is a protein called RuvB-like 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	435	Total	C	N	O	S	0	0	0
			3345	2092	592	647	14			

There are 25 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-24	MET	-	initiating methionine	UNP G0RYC2
B	-23	GLY	-	expression tag	UNP G0RYC2
B	-22	SER	-	expression tag	UNP G0RYC2
B	-21	SER	-	expression tag	UNP G0RYC2
B	-20	HIS	-	expression tag	UNP G0RYC2
B	-19	HIS	-	expression tag	UNP G0RYC2
B	-18	HIS	-	expression tag	UNP G0RYC2
B	-17	HIS	-	expression tag	UNP G0RYC2
B	-16	HIS	-	expression tag	UNP G0RYC2
B	-15	HIS	-	expression tag	UNP G0RYC2
B	-14	HIS	-	expression tag	UNP G0RYC2
B	-13	HIS	-	expression tag	UNP G0RYC2
B	-12	SER	-	expression tag	UNP G0RYC2
B	-11	SER	-	expression tag	UNP G0RYC2
B	-10	GLY	-	expression tag	UNP G0RYC2
B	-9	LEU	-	expression tag	UNP G0RYC2
B	-8	GLU	-	expression tag	UNP G0RYC2
B	-7	VAL	-	expression tag	UNP G0RYC2
B	-6	LEU	-	expression tag	UNP G0RYC2
B	-5	PHE	-	expression tag	UNP G0RYC2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLN	-	expression tag	UNP G0RYC2
B	-3	GLY	-	expression tag	UNP G0RYC2
B	-2	PRO	-	expression tag	UNP G0RYC2
B	-1	GLY	-	expression tag	UNP G0RYC2
B	0	SER	-	expression tag	UNP G0RYC2

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	O	0	0
			4	4		
4	B	10	Total	O	0	0
			10	10		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RuvB-like 1



• Molecule 2: RuvB-like 2



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	206.85Å 206.85Å 137.44Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	64.16 – 2.94 64.16 – 2.94	Depositor EDS
% Data completeness (in resolution range)	97.2 (64.16-2.94) 97.2 (64.16-2.94)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 2.96Å)	Xtriage
Refinement program	BUSTER-TNT	Depositor
R, R_{free}	0.198 , 0.223 0.214 , 0.244	Depositor DCC
R_{free} test set	1193 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	77.4	Xtriage
Anisotropy	0.270	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 92.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6603	wwPDB-VP
Average B, all atoms (Å ²)	99.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/3227	1.39	8/4353 (0.2%)
2	B	0.73	0/3383	1.34	4/4551 (0.1%)
All	All	0.75	0/6610	1.37	12/8904 (0.1%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	ALA	CA-C-N	7.77	133.68	121.56
1	A	218	ALA	C-N-CA	7.77	133.68	121.56
1	A	218	ALA	N-CA-C	-6.79	104.96	113.18
1	A	309	ASP	CA-CB-CG	5.59	118.19	112.60
2	B	17	GLY	CA-C-N	5.48	129.03	120.82
2	B	17	GLY	C-N-CA	5.48	129.03	120.82
2	B	405	ILE	CA-C-N	5.19	127.54	120.54
2	B	405	ILE	C-N-CA	5.19	127.54	120.54
1	A	245	ASP	CA-CB-CG	5.18	117.78	112.60
1	A	207	SER	CA-C-N	5.13	127.87	120.38
1	A	207	SER	C-N-CA	5.13	127.87	120.38
1	A	264	LEU	N-CA-C	-5.09	99.96	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3190	0	3272	22	0
2	B	3345	0	3430	35	0
3	A	27	0	12	0	0
3	B	27	0	12	0	0
4	A	4	0	0	0	0
4	B	10	0	0	0	0
All	All	6603	0	6726	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:246:ILE:HG22	2:B:275:ILE:HG12	1.79	0.65
1:A:387:LEU:HB3	1:A:426:VAL:HB	1.87	0.57
1:A:122:GLY:HA2	1:A:240:THR:HA	1.87	0.57
1:A:78:THR:HG22	1:A:301:PHE:CE2	2.40	0.56
2:B:294:VAL:HG22	2:B:322:ILE:HB	1.86	0.56
1:A:266:LYS:HG3	1:A:269:MET:H	1.73	0.54
2:B:163:LYS:HG2	2:B:168:GLU:HA	1.90	0.53
1:A:267:PRO:HB3	2:B:116:THR:HG22	1.90	0.53
1:A:340:ARG:HH22	2:B:338:ASP:H	1.57	0.52
2:B:72:ALA:HB3	2:B:353:VAL:HG12	1.91	0.52
2:B:80:SER:HB3	2:B:398:LEU:HB2	1.92	0.51
2:B:60:LEU:HA	2:B:63:ILE:HD12	1.93	0.50
1:A:88:LEU:HB3	1:A:92:ILE:HD13	1.93	0.50
2:B:9:VAL:HG23	2:B:249:ILE:HD11	1.95	0.49
2:B:300:VAL:HG12	2:B:303:LEU:HD12	1.96	0.48
2:B:333:ARG:HD3	2:B:340:LYS:HE3	1.94	0.48
2:B:59:VAL:HA	2:B:62:MET:HE3	1.96	0.48
1:A:71:GLY:O	1:A:333:ASN:HA	2.14	0.47
2:B:247:ASP:HA	2:B:275:ILE:HD13	1.96	0.47
2:B:160:LEU:HD13	2:B:181:MET:HE1	1.97	0.47
1:A:126:ARG:HE	1:A:236:VAL:HG12	1.80	0.46
2:B:58:VAL:HG12	2:B:62:MET:HE2	1.96	0.45
1:A:36:GLN:HG2	1:A:41:VAL:HG22	1.97	0.45
1:A:68:LEU:HD23	1:A:361:ILE:HG12	1.99	0.45
2:B:128:GLY:HA2	2:B:242:SER:HA	1.98	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:263:GLN:HG3	2:B:319:LEU:HD12	1.99	0.44
2:B:299:GLU:O	2:B:302:MET:HB2	2.18	0.44
2:B:74:LEU:HD23	2:B:325:MET:HG3	1.99	0.44
2:B:381:LEU:HD13	2:B:386:LEU:HG	2.00	0.44
2:B:268:ARG:HB2	2:B:271:ILE:HD12	2.00	0.44
1:A:356:LEU:HA	1:A:359:LEU:HD12	2.00	0.43
2:B:84:THR:HG22	2:B:296:PHE:CE2	2.53	0.43
1:A:78:THR:HG22	1:A:301:PHE:HE2	1.81	0.43
2:B:33:ASP:HB3	2:B:36:THR:O	2.18	0.43
1:A:102:ILE:HG12	1:A:113:LEU:HD13	2.00	0.42
1:A:445:ALA:HB2	2:B:342:PRO:HB3	2.01	0.42
1:A:303:ASP:HA	1:A:331:ALA:HB3	2.01	0.42
2:B:125:LYS:HE2	2:B:244:HIS:CE1	2.54	0.42
2:B:250:ASN:HB2	2:B:275:ILE:HD11	2.01	0.42
2:B:25:SER:HB2	2:B:185:ARG:HH21	1.85	0.42
2:B:300:VAL:HG11	2:B:325:MET:SD	2.60	0.42
2:B:176:LYS:H	2:B:176:LYS:HG3	1.70	0.41
2:B:9:VAL:HG23	2:B:249:ILE:CD1	2.49	0.41
2:B:364:ASP:O	2:B:368:GLN:HG2	2.21	0.41
2:B:105:ALA:HB1	2:B:303:LEU:HG	2.03	0.41
1:A:102:ILE:HG22	1:A:109:LYS:HG2	2.02	0.41
2:B:296:PHE:HA	2:B:324:ILE:O	2.21	0.41
1:A:3:GLN:HB3	1:A:248:ASN:ND2	2.35	0.40
1:A:441:ALA:HB1	2:B:342:PRO:HG2	2.04	0.40
1:A:123:LEU:HD22	1:A:293:ALA:HB1	2.03	0.40
2:B:392:ILE:HG21	2:B:404:LEU:HD12	2.03	0.40
1:A:25:LEU:HD12	1:A:87:GLU:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/462 (87%)	381 (95%)	20 (5%)	1 (0%)	43	65
2	B	427/513 (83%)	416 (97%)	11 (3%)	0	100	100
All	All	829/975 (85%)	797 (96%)	31 (4%)	1 (0%)	48	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	161	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/377 (90%)	301 (88%)	40 (12%)	5	15
2	B	363/423 (86%)	323 (89%)	40 (11%)	6	17
All	All	704/800 (88%)	624 (89%)	80 (11%)	5	16

All (80) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	ILE
1	A	61	LYS
1	A	88	LEU
1	A	92	ILE
1	A	105	THR
1	A	112	VAL
1	A	135	GLU
1	A	158	LEU
1	A	160	ILE
1	A	162	LEU
1	A	164	SER
1	A	174	ASP
1	A	183	LYS
1	A	194	ILE
1	A	195	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	198	THR
1	A	216	LEU
1	A	219	GLU
1	A	222	VAL
1	A	258	ILE
1	A	264	LEU
1	A	265	MET
1	A	269	MET
1	A	279	GLU
1	A	300	LEU
1	A	324	ILE
1	A	330	LEU
1	A	343	ASP
1	A	356	LEU
1	A	360	LEU
1	A	374	ARG
1	A	381	GLN
1	A	387	LEU
1	A	404	LEU
1	A	408	LEU
1	A	420	VAL
1	A	423	ARG
1	A	425	GLN
1	A	442	ARG
1	A	457	LEU
2	B	20	LEU
2	B	38	GLU
2	B	51	LYS
2	B	65	GLN
2	B	74	LEU
2	B	86	ILE
2	B	88	MET
2	B	92	GLN
2	B	161	THR
2	B	176	LYS
2	B	184	GLU
2	B	192	ILE
2	B	196	LYS
2	B	198	SER
2	B	200	LYS
2	B	201	ILE
2	B	233	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	234	ARG
2	B	245	GLU
2	B	259	LEU
2	B	270	GLU
2	B	275	ILE
2	B	278	LYS
2	B	290	ILE
2	B	300	VAL
2	B	302	MET
2	B	317	SER
2	B	333	ARG
2	B	338	ASP
2	B	345	LEU
2	B	368	GLN
2	B	377	GLU
2	B	378	GLU
2	B	386	LEU
2	B	389	LEU
2	B	395	GLU
2	B	399	ARG
2	B	406	THR
2	B	415	ARG
2	B	422	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	HIS
1	A	242	HIS
1	A	263	GLN
1	A	357	GLN
1	A	429	GLN
2	B	24	HIS
2	B	49	GLN
2	B	146	GLN
2	B	244	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ADP	A	501	-	28,29,29	0.36	0	43,45,45	0.59	0
3	ADP	B	501	-	28,29,29	0.40	0	43,45,45	0.67	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	501	-	-	5/16/32/32	0/3/3/3
3	ADP	B	501	-	-	7/16/32/32	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	ADP	O2B-PB-O3A	2.47	112.93	104.64

There are no chirality outliers.


All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	ADP	C5'-O5'-PA-O2A
3	A	501	ADP	C5'-O5'-PA-O3A
3	B	501	ADP	PA-O3A-PB-O2B
3	B	501	ADP	PA-O3A-PB-O3B
3	B	501	ADP	C5'-O5'-PA-O1A
3	B	501	ADP	C5'-O5'-PA-O2A
3	B	501	ADP	C5'-O5'-PA-O3A
3	A	501	ADP	C5'-O5'-PA-O1A
3	A	501	ADP	C4'-C5'-O5'-PA
3	B	501	ADP	C4'-C5'-O5'-PA
3	B	501	ADP	PA-O3A-PB-O1B
3	A	501	ADP	PB-O3A-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	416/462 (90%)	0.15	14 (3%) 48 40	57, 101, 152, 174	0
2	B	435/513 (84%)	-0.06	3 (0%) 84 79	55, 94, 128, 180	0
All	All	851/975 (87%)	0.04	17 (1%) 65 56	55, 97, 143, 180	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	264	LEU	4.0
1	A	103	TYR	3.2
1	A	110	THR	3.0
1	A	194	ILE	3.0
1	A	344	ASP	2.9
1	A	156	SER	2.9
1	A	102	ILE	2.7
1	A	188	VAL	2.7
2	B	149	ARG	2.6
2	B	169	ALA	2.3
1	A	39	GLY	2.3
1	A	159	LEU	2.3
1	A	157	THR	2.3
1	A	158	LEU	2.2
2	B	207	SER	2.1
1	A	249	ALA	2.1
1	A	258	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ADP	B	501	27/27	0.95	0.07	67,75,84,85	0
3	ADP	A	501	27/27	0.96	0.08	73,82,88,89	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around ADP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.