



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 11:10 AM UTC

PDB ID : 1WYZ / pdb_00001wyz
Title : X-Ray structure of the putative methyltransferase from Bacteroides thetaio-
taomicron VPI-5482 at the resolution 2.5 Å. Northeast Structural Genomics
Consortium target Btr28
Authors : Kuzin, A.P.; Chen, Y.; Forouhar, F.; Vorobiev, S.M.; Acton, T.; Ma, L.-
C.; Xiao, R.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural
Genomics Consortium (NESG)
Deposited on : 2005-02-21
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	242	10% 44% 41% 7% 8%
1	B	242	4% 52% 37% 5% 6%
1	C	242	4% 48% 40% • 8%
1	D	242	6% 52% 37% • 6%

Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7245 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called putative S-adenosylmethionine-dependent methyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	223	Total	C	N	O	S	Se	0	0	0
			1765	1134	301	323	4	3			
1	B	227	Total	C	N	O	S	Se	0	0	0
			1796	1154	305	330	4	3			
1	C	223	Total	C	N	O	S	Se	0	0	0
			1762	1134	297	324	4	3			
1	D	227	Total	C	N	O	S	Se	0	0	0
			1796	1154	305	330	4	3			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP Q8A031
A	86	MSE	MET	modified residue	UNP Q8A031
A	129	MSE	MET	modified residue	UNP Q8A031
A	181	MSE	MET	modified residue	UNP Q8A031
A	235	LEU	-	expression tag	UNP Q8A031
A	236	GLU	-	expression tag	UNP Q8A031
A	237	HIS	-	expression tag	UNP Q8A031
A	238	HIS	-	expression tag	UNP Q8A031
A	239	HIS	-	expression tag	UNP Q8A031
A	240	HIS	-	expression tag	UNP Q8A031
A	241	HIS	-	expression tag	UNP Q8A031
A	242	HIS	-	expression tag	UNP Q8A031
B	1	MSE	MET	modified residue	UNP Q8A031
B	86	MSE	MET	modified residue	UNP Q8A031
B	129	MSE	MET	modified residue	UNP Q8A031
B	181	MSE	MET	modified residue	UNP Q8A031
B	235	LEU	-	expression tag	UNP Q8A031
B	236	GLU	-	expression tag	UNP Q8A031
B	237	HIS	-	expression tag	UNP Q8A031
B	238	HIS	-	expression tag	UNP Q8A031
B	239	HIS	-	expression tag	UNP Q8A031

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	240	HIS	-	expression tag	UNP Q8A031
B	241	HIS	-	expression tag	UNP Q8A031
B	242	HIS	-	expression tag	UNP Q8A031
C	1	MSE	MET	modified residue	UNP Q8A031
C	86	MSE	MET	modified residue	UNP Q8A031
C	129	MSE	MET	modified residue	UNP Q8A031
C	181	MSE	MET	modified residue	UNP Q8A031
C	235	LEU	-	expression tag	UNP Q8A031
C	236	GLU	-	expression tag	UNP Q8A031
C	237	HIS	-	expression tag	UNP Q8A031
C	238	HIS	-	expression tag	UNP Q8A031
C	239	HIS	-	expression tag	UNP Q8A031
C	240	HIS	-	expression tag	UNP Q8A031
C	241	HIS	-	expression tag	UNP Q8A031
C	242	HIS	-	expression tag	UNP Q8A031
D	1	MSE	MET	modified residue	UNP Q8A031
D	86	MSE	MET	modified residue	UNP Q8A031
D	129	MSE	MET	modified residue	UNP Q8A031
D	181	MSE	MET	modified residue	UNP Q8A031
D	235	LEU	-	expression tag	UNP Q8A031
D	236	GLU	-	expression tag	UNP Q8A031
D	237	HIS	-	expression tag	UNP Q8A031
D	238	HIS	-	expression tag	UNP Q8A031
D	239	HIS	-	expression tag	UNP Q8A031
D	240	HIS	-	expression tag	UNP Q8A031
D	241	HIS	-	expression tag	UNP Q8A031
D	242	HIS	-	expression tag	UNP Q8A031


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	34	Total O 34 34	0	0
2	B	38	Total O 38 38	0	0
2	C	28	Total O 28 28	0	0
2	D	26	Total O 26 26	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: putative S-adenosylmethionine-dependent methyltransferase




- Molecule 1: putative S-adenosylmethionine-dependent methyltransferase




- Molecule 1: putative S-adenosylmethionine-dependent methyltransferase





- Molecule 1: putative S-adenosylmethionine-dependent methyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.78Å 101.54Å 108.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.74 – 2.50 28.74 – 2.50	Depositor EDS
% Data completeness (in resolution range)	88.3 (28.74-2.50) 96.0 (28.74-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.89 (at 2.48Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.226 , 0.293 0.231 , 0.295	Depositor DCC
R_{free} test set	3077 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.511	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 67.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7245	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/1798	1.06	11/2428 (0.5%)
1	B	0.49	1/1830 (0.1%)	1.01	7/2473 (0.3%)
1	C	0.48	0/1794	1.02	7/2423 (0.3%)
1	D	0.45	0/1830	0.97	6/2473 (0.2%)
All	All	0.47	1/7252 (0.0%)	1.01	31/9797 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	199	ALA	CA-C	5.06	1.54	1.52

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	205	GLY	N-CA-C	-11.72	95.22	115.61
1	A	8	LEU	N-CA-C	8.36	120.85	109.24
1	A	226	ILE	CA-C-N	8.03	128.08	119.89
1	A	226	ILE	C-N-CA	8.03	128.08	119.89
1	A	150	GLU	N-CA-C	-8.01	102.41	112.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1765	0	1808	123	0
1	B	1796	0	1842	100	0
1	C	1762	0	1808	91	0
1	D	1796	0	1842	88	0
2	A	34	0	0	1	0
2	B	38	0	0	3	0
2	C	28	0	0	1	0
2	D	26	0	0	1	0
All	All	7245	0	7300	375	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 375 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:44:ARG:HG3	1:C:56:ILE:CD1	1.54	1.37
1:C:44:ARG:CG	1:C:56:ILE:HD11	1.64	1.28
1:A:36:ILE:HD13	1:A:62:TYR:HB2	1.37	1.05
1:A:73:ILE:HB	1:A:107:ILE:HD12	1.44	1.00
1:B:36:ILE:HD13	1:B:62:TYR:HB2	1.44	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/242 (90%)	197 (91%)	17 (8%)	3 (1%)	9	17
1	B	223/242 (92%)	201 (90%)	18 (8%)	4 (2%)	6	12
1	C	217/242 (90%)	196 (90%)	19 (9%)	2 (1%)	14	27
1	D	223/242 (92%)	201 (90%)	20 (9%)	2 (1%)	14	27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	880/968 (91%)	795 (90%)	74 (8%)	11 (1%)	9	18

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	59	LEU
1	B	82	GLY
1	C	101	GLY
1	C	200	ASN
1	A	56	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/207 (95%)	186 (95%)	10 (5%)	21	43
1	B	200/207 (97%)	186 (93%)	14 (7%)	14	29
1	C	196/207 (95%)	182 (93%)	14 (7%)	13	29
1	D	200/207 (97%)	186 (93%)	14 (7%)	14	29
All	All	792/828 (96%)	740 (93%)	52 (7%)	15	32

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	57	ASP
1	C	170	LEU
1	D	212	THR
1	C	64	LEU
1	C	89	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	160	GLN
1	D	34	HIS
1	D	200	ASN
1	C	192	GLN
1	D	109	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	220/242 (90%)	0.52	24 (10%) 10 9	21, 44, 99, 126	0
1	B	224/242 (92%)	0.24	10 (4%) 38 33	15, 41, 84, 153	0
1	C	220/242 (90%)	0.47	10 (4%) 38 33	16, 45, 85, 123	0
1	D	224/242 (92%)	0.35	15 (6%) 24 21	16, 42, 82, 116	0
All	All	888/968 (91%)	0.39	59 (6%) 24 21	15, 43, 88, 153	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	56	ILE	5.6
1	D	64	LEU	5.2
1	A	59	LEU	5.0
1	A	225	LYS	4.9
1	B	68	THR	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.