



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 08:01 PM UTC

PDB ID : 2WYY / pdb_00002wyy
EMDB ID : EMD-1663
Title : CRYOEM MODEL OF THE VESICULAR STOMATITIS VIRUS
Authors : Ge, P.; Tsao, J.; Green, T.J.; Luo, M.; Zhou, Z.H.
Deposited on : 2009-11-20
Resolution : 10.60 Å(reported)
Based on initial model : 2WYY

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : **NOT EXECUTED**
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 10.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	229148	23984
Ramachandran outliers	224038	23583
Sidechain outliers	223484	23102
RNA backbone	8273	3508

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	98% 64% 28% 5%
1	C	422	98% 64% 28% 5%
1	D	422	98% 65% 27% 5%
1	F	422	98% 65% 27% 5%
1	H	422	98% 65% 27% 6%
1	I	422	98% 65% 27% 6%
1	J	422	98% 65% 27% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	422	98% 66% 27% 5%
1	L	422	98% 65% 27% 6%
1	M	422	98% 65% 27% 6%
2	R	45	100% 38% 53% 9%
2	S	45	100% 33% 56% 9%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 34620 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NUCLEOPROTEIN.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	414	Total	C	N	O	S	0	0
			3282	2093	551	620	18		
1	C	414	Total	C	N	O	S	0	0
			3282	2093	551	620	18		
1	D	414	Total	C	N	O	S	0	0
			3282	2093	551	620	18		
1	F	414	Total	C	N	O	S	0	0
			3282	2093	551	620	18		
1	H	414	Total	C	N	O	S	0	0
			3282	2093	551	620	18		
1	I	414	Total	C	N	O	S	0	0
			3282	2093	551	620	18		
1	J	414	Total	C	N	O	S	0	0
			3282	2093	551	620	18		
1	K	414	Total	C	N	O	S	0	0
			3282	2093	551	620	18		
1	L	414	Total	C	N	O	S	0	0
			3282	2093	551	620	18		
1	M	414	Total	C	N	O	S	0	0
			3282	2093	551	620	18		


- Molecule 2 is a RNA chain called POLY-URIDINE.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	R	45	Total	C	N	O	P	0	0
			900	405	90	360	45		
2	S	45	Total	C	N	O	P	0	0
			900	405	90	360	45		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NUCLEOPROTEIN




• Molecule 1: NUCLEOPROTEIN




D421
K422

- Molecule 1: NUCLEOPROTEIN

MET
S2




• Molecule 1: NUCLEOPROTEIN



• Molecule 1: NUCLEOPROTEIN




- Molecule 1: NUCLEOPROTEIN





• Molecule 1: NUCLEOPROTEIN

Chain J:




• Molecule 1: NUCLEOPROTEIN

Chain K:







• Molecule 1: NUCLEOPROTEIN



• Molecule 2: POLY-URIDINE



• Molecule 2: POLY-URIDINE



4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=Not provided°, rise=Not provided Å, axial sym=Not provided	Depositor
Number of segments used	Not provided	
Resolution determination method	Not provided	
CTF correction method	Not provided	
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	20	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	98000	Depositor
Image detector	GENERIC TVIPS	Depositor
Maximum map value	5.997	Depositor
Minimum map value	-6.598	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.999	Depositor
Recommended contour level	1	Depositor
Map size (Å)	490.24, 490.24, 490.24	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90, 90, 90	wwPDB
Pixel spacing (Å)	1.532, 1.532, 1.532	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	1/3357 (0.0%)	0.94	4/4543 (0.1%)
1	C	0.70	1/3357 (0.0%)	0.94	4/4543 (0.1%)
1	D	0.65	1/3357 (0.0%)	0.92	4/4543 (0.1%)
1	F	0.70	1/3357 (0.0%)	0.93	4/4543 (0.1%)
1	H	0.70	1/3357 (0.0%)	0.94	4/4543 (0.1%)
1	I	0.70	1/3357 (0.0%)	0.94	4/4543 (0.1%)
1	J	0.70	1/3357 (0.0%)	0.94	4/4543 (0.1%)
1	K	0.65	1/3357 (0.0%)	0.92	4/4543 (0.1%)
1	L	0.70	1/3357 (0.0%)	0.93	4/4543 (0.1%)
1	M	0.70	1/3357 (0.0%)	0.94	4/4543 (0.1%)
2	R	0.70	0/989	0.97	4/1526 (0.3%)
2	S	0.70	0/989	0.97	4/1526 (0.3%)
All	All	0.69	10/35548 (0.0%)	0.94	48/48482 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
1	D	0	1
1	F	0	1
1	H	0	1
1	I	0	1
1	J	0	1
1	K	0	1
1	L	0	1
1	M	0	1
All	All	0	10

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	64	VAL	C-N	-6.18	1.25	1.33
1	I	64	VAL	C-N	-6.14	1.25	1.33
1	D	64	VAL	C-N	-5.95	1.25	1.33
1	H	64	VAL	C-N	-5.92	1.25	1.33
1	K	64	VAL	C-N	-5.92	1.25	1.33

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	263	LEU	CA-C-N	7.07	127.38	119.32
1	M	263	LEU	C-N-CA	7.07	127.38	119.32
1	H	263	LEU	CA-C-N	7.05	127.35	119.32
1	H	263	LEU	C-N-CA	7.05	127.35	119.32
1	D	263	LEU	CA-C-N	6.86	127.14	119.32

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	106	ASP	Peptide
1	C	106	ASP	Peptide
1	D	106	ASP	Peptide
1	F	106	ASP	Peptide
1	H	106	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3282	0	3249	112	0
1	C	3282	0	3249	115	0
1	D	3282	0	3249	114	0
1	F	3282	0	3249	113	0
1	H	3282	0	3249	107	0
1	I	3282	0	3249	114	0
1	J	3282	0	3249	103	0
1	K	3282	0	3249	112	0
1	L	3282	0	3249	114	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	M	3282	0	3249	115	0
2	R	900	0	451	70	0
2	S	900	0	450	70	0
All	All	34620	0	33391	1101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 1101 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:184:VAL:HG21	1:M:166:MET:CE	1.39	1.51
1:L:166:MET:CE	1:M:184:VAL:HG21	1.40	1.51
1:I:184:VAL:HG21	1:J:166:MET:CE	1.40	1.51
1:D:166:MET:CE	1:F:184:VAL:HG21	1.40	1.50
1:F:166:MET:CE	1:H:184:VAL:HG21	1.40	1.50

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	410/422 (97%)	341 (83%)	45 (11%)	24 (6%)	1	13
1	C	410/422 (97%)	341 (83%)	45 (11%)	24 (6%)	1	13
1	D	410/422 (97%)	341 (83%)	45 (11%)	24 (6%)	1	13
1	F	410/422 (97%)	341 (83%)	45 (11%)	24 (6%)	1	13
1	H	410/422 (97%)	340 (83%)	46 (11%)	24 (6%)	1	13
1	I	410/422 (97%)	341 (83%)	45 (11%)	24 (6%)	1	13
1	J	410/422 (97%)	341 (83%)	45 (11%)	24 (6%)	1	13

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	K	410/422 (97%)	341 (83%)	45 (11%)	24 (6%)	1	13
1	L	410/422 (97%)	341 (83%)	45 (11%)	24 (6%)	1	13
1	M	410/422 (97%)	340 (83%)	46 (11%)	24 (6%)	1	13
All	All	4100/4220 (97%)	3408 (83%)	452 (11%)	240 (6%)	2	13

5 of 240 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	24	PRO
1	A	98	ALA
1	A	128	SER
1	A	150	PRO
1	A	168	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/363 (98%)	310 (87%)	47 (13%)	4	15
1	C	357/363 (98%)	309 (87%)	48 (13%)	4	14
1	D	357/363 (98%)	310 (87%)	47 (13%)	4	15
1	F	357/363 (98%)	308 (86%)	49 (14%)	3	14
1	H	357/363 (98%)	310 (87%)	47 (13%)	4	15
1	I	357/363 (98%)	310 (87%)	47 (13%)	4	15
1	J	357/363 (98%)	309 (87%)	48 (13%)	4	14
1	K	357/363 (98%)	310 (87%)	47 (13%)	4	15
1	L	357/363 (98%)	308 (86%)	49 (14%)	3	14
1	M	357/363 (98%)	310 (87%)	47 (13%)	4	15
All	All	3570/3630 (98%)	3094 (87%)	476 (13%)	6	14

5 of 476 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	354	LYS
1	M	149	MET
1	J	25	VAL
1	M	107	LEU
1	M	407	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 94 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	385	GLN
1	K	187	ASN
1	J	148	GLN
1	J	347	GLN
1	K	385	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	R	44/45 (97%)	34 (77%)	1 (2%)
2	S	44/45 (97%)	34 (77%)	1 (2%)
All	All	88/90 (97%)	68 (77%)	2 (2%)

5 of 68 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	R	19	U
2	R	21	U
2	R	23	U
2	R	24	U
2	R	25	U

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	R	24	U
2	S	24	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-1663. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 212



Y Index: 213




Z Index: 317


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 1.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_1663_msk_1.map [i](#)



X




Y




Z


6.6.2 emd_1663_msk_2.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 10476 nm³; this corresponds to an approximate mass of 9463 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.094 $\text{\AA}^{-1}$

8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-1663 and PDB model 2WYY. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay [i](#)


This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (1).

9.4 Atom inclusion [i](#)



At the recommended contour level, 0% of all backbone atoms, 0% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (1) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.0000	0.0000
A	0.0000	0.0000
C	0.0000	-0.0000
D	0.0000	0.0000
F	0.0000	0.0000
H	0.0000	0.0000
I	0.0000	0.0010
J	0.0000	0.0030
K	0.0000	0.0000
L	0.0000	-0.0000
M	0.0000	0.0000
R	0.0000	0.0000
S	0.0000	0.0010

1.0

0.0

<0.0