



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:31 AM UTC

PDB ID : 3WY4 / pdb_00003wy4
Title : Crystal structure of alpha-glucosidase mutant E271Q in complex with maltose
Authors : Shen, X.; Gai, Z.; Kato, K.; Yao, M.
Deposited on : 2014-08-18
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	538	88% 11% .
1	B	538	9% 62% 12% .. 22%
2	C	2	50% 50%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-glucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	535	Total	C	N	O	S	0	0	0
			4292	2730	735	809	18			
1	B	421	Total	C	N	O	S	0	0	0
			3398	2172	578	634	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	271	GLN	GLU	engineered mutation	UNP H3K096
B	271	GLN	GLU	engineered mutation	UNP H3K096

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	293	Total	O	0	0
			293	293		
5	B	159	Total	O	0	0
			159	159		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-glucosidase



- Molecule 1: Alpha-glucosidase



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	111.46Å 181.06Å 51.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.46 – 2.50 47.46 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.46-2.50) 99.6 (47.46-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.202 , 0.241 0.210 , 0.242	Depositor DCC
R_{free} test set	1865 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	37.1	Xtriage
Anisotropy	0.594	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8190	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GLC, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/4423	0.74	4/6033 (0.1%)
1	B	0.29	0/3496	0.85	16/4762 (0.3%)
All	All	0.28	0/7919	0.79	20/10795 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	441	ASP	CA-C-N	8.64	130.64	119.84
1	B	441	ASP	C-N-CA	8.64	130.64	119.84
1	B	302	MET	CA-C-N	-7.80	111.93	119.89
1	B	302	MET	C-N-CA	-7.80	111.93	119.89
1	B	219	VAL	CA-C-N	6.58	128.06	119.84
1	B	219	VAL	C-N-CA	6.58	128.06	119.84
1	B	297	PHE	CA-C-N	6.37	133.17	121.70
1	B	297	PHE	C-N-CA	6.37	133.17	121.70
1	A	177	HIS	CA-C-N	5.85	125.32	119.24
1	A	177	HIS	C-N-CA	5.85	125.32	119.24
1	B	304	HIS	CA-C-N	5.67	131.90	121.70
1	B	304	HIS	C-N-CA	5.67	131.90	121.70
1	B	301	ASN	CA-C-N	5.58	131.74	121.70
1	B	301	ASN	C-N-CA	5.58	131.74	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	17	TYR	CA-C-N	5.55	125.91	119.47
1	A	17	TYR	C-N-CA	5.55	125.91	119.47
1	B	276	ASN	CA-C-N	5.49	124.68	118.97
1	B	276	ASN	C-N-CA	5.49	124.68	118.97
1	B	472	VAL	N-CA-C	-5.33	107.94	113.10
1	B	514	ALA	CB-CA-C	-5.22	110.54	116.54

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	58	SER	Peptide
1	B	59	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4292	0	4053	41	0
1	B	3398	0	3223	62	0
2	C	23	0	21	0	0
3	A	1	0	0	0	0
4	A	18	0	24	3	0
4	B	6	0	8	2	0
5	A	293	0	0	11	0
5	B	159	0	0	11	0
All	All	8190	0	7329	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:44:SER:OG	5:B:800:HOH:O	1.81	0.96

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:480:THR:OG1	5:B:729:HOH:O	1.95	0.84
1:A:514:ALA:O	5:A:873:HOH:O	1.96	0.83
1:B:139:PRO:O	5:B:795:HOH:O	1.96	0.82
1:B:297:PHE:HA	1:B:298:ASP:HB2	1.61	0.80
1:A:396:GLU:OE1	5:A:934:HOH:O	2.03	0.77
1:B:305:SER:HB3	1:B:308:TYR:HB3	1.67	0.77
1:A:209:HIS:HB3	4:A:601:GOL:H32	1.66	0.76
1:B:499:GLN:OE1	5:B:731:HOH:O	2.05	0.74
1:B:297:PHE:HA	1:B:298:ASP:CB	2.16	0.74
1:B:179:GLU:OE1	5:B:777:HOH:O	2.06	0.73
1:A:311:GLU:OE1	5:A:827:HOH:O	2.07	0.72
1:A:315:ARG:NH1	5:A:962:HOH:O	2.19	0.71
1:A:289:ASP:OD2	5:A:888:HOH:O	2.09	0.70
1:A:328:ALA:O	5:A:802:HOH:O	2.07	0.70
1:A:25:ARG:NH2	1:A:33:ASN:OD1	2.25	0.70
1:B:19:ARG:HD3	1:B:372:GLU:HB3	1.73	0.69
1:B:447:ASN:HA	1:B:450:ARG:HD2	1.75	0.69
1:B:298:ASP:O	1:B:300:LEU:N	2.28	0.66
1:A:172:ASP:OD1	5:A:710:HOH:O	2.13	0.66
1:B:472:VAL:HG21	1:B:478:GLY:HA3	1.79	0.64
1:B:301:ASN:N	1:B:301:ASN:OD1	2.31	0.63
1:B:301:ASN:HB2	1:B:302:MET:HA	1.82	0.62
1:B:200:ARG:NH2	1:B:271:GLN:OE1	2.33	0.62
1:A:105:HIS:O	5:A:703:HOH:O	2.16	0.61
1:A:441:ASP:O	1:A:447:ASN:ND2	2.32	0.61
1:A:438:GLN:HG3	1:A:444:ALA:HB1	1.83	0.61
1:A:304:HIS:NE2	5:A:733:HOH:O	2.32	0.60
1:B:165:ASN:ND2	4:B:601:GOL:O3	2.34	0.60
1:B:360:SER:O	1:B:481:ARG:NH1	2.37	0.58
1:B:165:ASN:HD22	1:B:206:PHE:HD1	1.52	0.57
1:B:120:ARG:NH2	5:B:845:HOH:O	2.38	0.56
1:A:138:THR:HG21	1:B:142:ASN:HB2	1.88	0.56
1:B:61:LEU:HG	1:B:62:ASP:H	1.70	0.56
1:B:304:HIS:HA	1:B:305:SER:HB2	1.88	0.56
1:B:59:PRO:HG2	5:B:840:HOH:O	2.05	0.55
1:B:372:GLU:O	5:B:848:HOH:O	2.18	0.55
1:A:153:THR:OG1	1:A:164:HIS:NE2	2.37	0.55
1:A:225:LYS:HE3	1:A:231:GLU:HG3	1.87	0.54
1:B:469:LEU:HD23	1:B:479:PHE:HB3	1.89	0.53
1:A:19:ARG:NH2	1:A:76:MET:HE1	2.24	0.52
1:A:438:GLN:HG2	1:A:446:LEU:HB3	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:ASN:ND2	5:A:904:HOH:O	2.40	0.51
1:B:301:ASN:CB	1:B:302:MET:HA	2.40	0.51
1:B:329:THR:HG21	1:B:367:LEU:HD12	1.93	0.51
1:B:61:LEU:O	1:B:62:ASP:HB2	2.11	0.50
1:A:310:ARG:NH1	1:A:476:LEU:O	2.43	0.50
1:B:61:LEU:HG	1:B:63:PHE:H	1.77	0.50
1:B:80:LEU:HG	1:B:84:LYS:HE3	1.94	0.49
1:B:58:SER:O	1:B:60:MET:HB2	2.12	0.49
1:A:165:ASN:ND2	1:A:172:ASP:OD2	2.41	0.49
1:A:129:VAL:HB	1:A:163:LEU:HB3	1.94	0.48
1:B:206:PHE:O	4:B:601:GOL:H12	2.14	0.48
1:B:442:PRO:HA	1:B:444:ALA:H	1.78	0.48
1:A:204:VAL:HG21	1:A:291:LEU:HD11	1.96	0.48
1:B:476:LEU:HD21	1:B:527:LEU:HD13	1.95	0.48
1:A:63:PHE:CZ	1:A:386:GLN:HB3	2.49	0.48
1:A:206:PHE:O	4:A:601:GOL:H12	2.14	0.48
1:B:329:THR:OG1	1:B:368:TYR:N	2.43	0.47
1:B:129:VAL:HB	1:B:163:LEU:HB3	1.96	0.47
1:B:347:PRO:HB3	1:B:350:TYR:HB3	1.96	0.47
1:B:209:HIS:HD2	5:B:833:HOH:O	1.97	0.47
1:A:408:PRO:HG3	1:A:430:HIS:HB3	1.96	0.47
1:B:297:PHE:CZ	1:B:327:TRP:HA	2.50	0.46
1:A:466:ASP:O	1:A:481:ARG:HA	2.15	0.46
1:B:442:PRO:CA	1:B:444:ALA:H	2.28	0.46
1:A:328:ALA:HB2	1:A:366:CYS:HB2	1.97	0.46
1:B:40:ASP:OD2	1:B:89:LYS:NZ	2.44	0.46
1:B:219:VAL:HG22	1:B:239:ARG:HG3	1.97	0.45
1:A:400:ARG:O	1:A:404:ARG:HD3	2.17	0.44
1:A:66:ASP:OD1	1:A:66:ASP:N	2.50	0.44
1:A:17:TYR:CE2	1:A:404:ARG:HG2	2.52	0.44
1:B:347:PRO:HB2	1:B:348:HIS:H	1.57	0.44
1:B:153:THR:OG1	1:B:164:HIS:NE2	2.44	0.44
1:A:469:LEU:HD23	1:A:479:PHE:HB3	2.00	0.43
1:A:503:LEU:HD11	1:A:520:ARG:HD2	2.00	0.43
1:B:81:GLU:OE1	1:B:81:GLU:N	2.42	0.43
1:B:297:PHE:CE1	1:B:327:TRP:HA	2.53	0.43
1:A:219:VAL:HG22	1:A:239:ARG:HG3	1.99	0.43
1:B:347:PRO:C	1:B:349:ALA:H	2.25	0.43
1:A:112:TRP:CD1	1:A:124:LYS:HE2	2.54	0.43
1:B:4:ASN:N	5:B:719:HOH:O	2.51	0.43
1:B:302:MET:HE2	1:B:302:MET:HB2	1.93	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:260:MET:HE2	1:B:268:THR:HG23	2.01	0.42
1:B:508:ALA:N	1:B:536:GLN:O	2.49	0.42
1:B:175:PHE:HB3	1:B:208:PHE:CZ	2.54	0.42
1:A:51:TRP:CD1	1:A:51:TRP:C	2.97	0.42
1:B:235:TYR:O	1:B:240:HIS:NE2	2.41	0.42
1:B:326:CYS:SG	5:B:759:HOH:O	2.62	0.42
1:A:441:ASP:HB3	1:A:444:ALA:HB2	2.01	0.42
1:A:288:GLY:HA2	5:A:777:HOH:O	2.18	0.42
1:A:220:PRO:HB2	1:A:223:GLU:HG3	2.02	0.41
1:B:166:PHE:CZ	1:B:203:THR:HG21	2.55	0.41
1:A:172:ASP:HB3	4:A:601:GOL:H2	2.02	0.41
1:B:57:THR:O	1:B:58:SER:OG	2.33	0.41
1:B:466:ASP:O	1:B:481:ARG:HA	2.21	0.41
1:B:441:ASP:HA	1:B:442:PRO:HD2	1.67	0.40
1:A:438:GLN:HG3	1:A:444:ALA:CB	2.50	0.40
1:B:325:PRO:HG2	1:B:363:GLY:HA3	2.03	0.40
1:B:140:PRO:HG2	1:B:152:TRP:CE2	2.56	0.40
1:B:206:PHE:HA	1:B:244:LEU:HB2	2.03	0.40
1:B:453:LEU:HD23	1:B:453:LEU:HA	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	533/538 (99%)	516 (97%)	17 (3%)	0	100	100
1	B	409/538 (76%)	380 (93%)	16 (4%)	13 (3%)	3	4
All	All	942/1076 (88%)	896 (95%)	33 (4%)	13 (1%)	9	17

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	58	SER
1	B	62	ASP
1	B	220	PRO
1	B	298	ASP
1	B	299	LEU
1	B	302	MET
1	B	304	HIS
1	B	347	PRO
1	B	297	PHE
1	B	305	SER
1	B	348	HIS
1	B	442	PRO
1	B	321	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	458/461 (99%)	452 (99%)	6 (1%)	61	82
1	B	365/461 (79%)	357 (98%)	8 (2%)	45	73
All	All	823/922 (89%)	809 (98%)	14 (2%)	53	78

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ARG
1	A	100	ASP
1	A	221	LYS
1	A	241	VAL
1	A	244	LEU
1	A	319	LEU
1	B	60	MET
1	B	61	LEU
1	B	62	ASP
1	B	63	PHE
1	B	244	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	300	LEU
1	B	301	ASN
1	B	304	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	160	GLN
1	A	438	GLN
1	B	105	HIS
1	B	165	ASN
1	B	170	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	C	1	2	12,12,12	0.52	0	17,17,17	0.55	0
2	GLC	C	2	2	11,11,12	0.49	0	15,15,17	1.71	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	GLC	C1-O5-C5	5.43	119.47	112.19


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	602	-	5,5,5	0.38	0	5,5,5	0.25	0
4	GOL	A	601	-	5,5,5	0.38	0	5,5,5	0.23	0
4	GOL	B	601	-	5,5,5	0.37	0	5,5,5	0.27	0
4	GOL	A	603	-	5,5,5	0.38	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	602	-	-	2/4/4/4	-
4	GOL	A	601	-	-	2/4/4/4	-
4	GOL	B	601	-	-	2/4/4/4	-
4	GOL	A	603	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	601	GOL	O1-C1-C2-C3
4	B	601	GOL	O1-C1-C2-C3
4	A	602	GOL	O1-C1-C2-C3
4	A	603	GOL	O1-C1-C2-C3
4	A	601	GOL	O1-C1-C2-O2
4	A	603	GOL	O1-C1-C2-O2
4	B	601	GOL	O1-C1-C2-O2
4	A	602	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	601	GOL	3	0
4	B	601	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	535/538 (99%)	0.06	5 (0%) 81 78	26, 34, 52, 68	0
1	B	421/538 (78%)	0.69	49 (11%) 9 8	28, 45, 86, 123	0
All	All	956/1076 (88%)	0.34	54 (5%) 30 26	26, 38, 71, 123	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	63	PHE	5.9
1	B	347	PRO	5.4
1	B	303	PRO	5.0
1	B	30	GLY	4.8
1	B	369	GLN	4.4
1	B	442	PRO	4.4
1	B	61	LEU	4.4
1	B	62	ASP	4.4
1	B	328	ALA	4.1
1	B	348	HIS	4.0
1	B	232	ALA	3.9
1	B	330	SER	3.7
1	B	305	SER	3.6
1	B	297	PHE	3.4
1	B	346	ASP	3.4
1	B	444	ALA	3.3
1	B	302	MET	3.1
1	B	295	TYR	3.1
1	B	304	HIS	2.9
1	B	22	LEU	2.9
1	B	300	LEU	2.9
1	B	319	LEU	2.9
1	B	79	THR	2.8
1	A	221	LYS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	80	LEU	2.7
1	B	34	GLY	2.6
1	B	446	LEU	2.6
1	B	221	LYS	2.6
1	B	443	ASN	2.6
1	B	296	THR	2.5
1	B	60	MET	2.5
1	B	275	ASP	2.5
1	B	441	ASP	2.5
1	B	320	ALA	2.5
1	B	58	SER	2.5
1	B	306	ALA	2.4
1	B	17	TYR	2.4
1	B	368	TYR	2.4
1	A	348	HIS	2.4
1	B	308	TYR	2.4
1	A	514	ALA	2.4
1	B	322	ASP	2.3
1	B	231	GLU	2.3
1	B	298	ASP	2.3
1	B	329	THR	2.2
1	B	4	ASN	2.2
1	B	372	GLU	2.2
1	B	20	SER	2.1
1	B	299	LEU	2.1
1	B	219	VAL	2.1
1	B	496	GLY	2.1
1	A	75	PRO	2.1
1	A	522	GLY	2.1
1	B	21	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	C	1	12/12	0.88	0.12	32,36,40,40	0
2	GLC	C	2	11/12	0.91	0.10	30,32,35,36	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	601	6/6	0.70	0.26	35,35,37,40	0
4	GOL	B	601	6/6	0.74	0.31	42,45,46,46	0
4	GOL	A	603	6/6	0.79	0.23	39,40,41,42	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	602	6/6	0.86	0.16	42,43,44,47	0
3	MG	A	600	1/1	0.93	0.10	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.