



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 05:49 PM EDT

PDB ID : 3WYH / pdb_00003wyh
Title : Structure of disulfide bond deletion mutant of ostrich egg white lysozyme
Authors : Kawaguchi, Y.; Yoneda, K.; Araki, T.
Deposited on : 2014-08-28
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	185	 61% 37%
1	B	185	 59% 38%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	TAM	A	201	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3200 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lysozyme g.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	185	Total	C	N	O	S	0	0	0
			1443	892	274	272	5			
1	B	185	Total	C	N	O	S	0	0	0
			1443	892	274	272	5			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	4	SER	CYS	engineered mutation	UNP P00719
A	18	SER	CYS	engineered mutation	UNP P00719
A	29	SER	CYS	engineered mutation	UNP P00719
A	60	SER	CYS	engineered mutation	UNP P00719
B	4	SER	CYS	engineered mutation	UNP P00719
B	18	SER	CYS	engineered mutation	UNP P00719
B	29	SER	CYS	engineered mutation	UNP P00719
B	60	SER	CYS	engineered mutation	UNP P00719

- Molecule 2 is TRIS(HYDROXYETHYL)AMINOMETHANE (CCD ID: TAM) (formula: $C_7H_{17}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			11	7	1	3		
2	B	1	Total	C	N	O	0	0
			11	7	1	3		

- Molecule 3 is HEXAETHYLENE GLYCOL (CCD ID: P6G) (formula: $C_{12}H_{26}O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			19	12	7		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	150	Total 150	O 150	0	0
4	B	123	Total 123	O 123	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Lysozyme g



• Molecule 1: Lysozyme g



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	38.19Å 64.93Å 131.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.10 – 1.77 29.10 – 1.77	Depositor EDS
% Data completeness (in resolution range)	89.9 (29.10-1.77) 99.8 (29.10-1.77)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.34 (at 1.77Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.186 , 0.224 0.191 , 0.196	Depositor DCC
R_{free} test set	1654 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	15.6	Xtriage
Anisotropy	0.269	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3200	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TAM, P6G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.17	64/1469 (4.4%)	1.07	1/1970 (0.1%)
1	B	2.21	70/1469 (4.8%)	1.11	1/1970 (0.1%)
All	All	2.19	134/2938 (4.6%)	1.09	2/3940 (0.1%)

All (134) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	185	TYR	C-OXT	-18.52	0.86	1.23
1	B	33	ALA	CA-CB	-9.00	1.39	1.53
1	B	96	VAL	C-O	-8.96	1.14	1.24
1	A	45	MET	C-O	-8.59	1.14	1.24
1	A	168	ASP	C-O	-8.52	1.14	1.24
1	A	73	GLU	C-O	-8.46	1.14	1.24
1	A	185	TYR	C-OXT	-8.46	1.06	1.23
1	A	116	GLY	C-O	-8.25	1.13	1.23
1	A	58	LYS	C-O	-7.77	1.15	1.24
1	A	172	ASP	C-O	-7.63	1.15	1.24
1	B	13	THR	C-O	-7.53	1.14	1.23
1	B	7	ASP	C-O	-7.43	1.15	1.24
1	B	149	ALA	CA-CB	-7.39	1.42	1.53
1	B	138	GLN	C-O	-7.35	1.15	1.24
1	B	122	SER	C-O	-7.30	1.15	1.24
1	A	157	TYR	C-O	-7.24	1.16	1.24
1	B	181	LYS	C-O	-7.22	1.15	1.24
1	B	69	ILE	C-O	-7.19	1.15	1.24
1	A	160	MET	C-O	-7.16	1.15	1.24
1	B	178	GLN	C-O	-7.02	1.16	1.24
1	B	57	GLN	C-O	-6.97	1.16	1.24
1	A	44	SER	C-O	-6.95	1.16	1.24
1	A	41	ASP	C-O	-6.85	1.14	1.24
1	B	136	LYS	C-O	-6.84	1.16	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	82	ASN	C-O	-6.71	1.15	1.23
1	B	115	GLN	C-O	-6.70	1.16	1.24
1	B	120	LEU	C-O	-6.57	1.16	1.24
1	B	157	TYR	C-O	-6.56	1.16	1.24
1	B	179	TYR	C-O	-6.51	1.16	1.24
1	B	2	THR	C-O	-6.46	1.16	1.24
1	B	26	LEU	C-O	-6.43	1.16	1.24
1	B	35	ARG	C-O	-6.42	1.16	1.24
1	A	61	VAL	C-O	-6.41	1.17	1.24
1	B	177	ALA	C-O	-6.41	1.16	1.24
1	B	12	ASP	C-O	-6.41	1.15	1.23
1	A	118	GLU	C-O	-6.41	1.16	1.24
1	A	20	SER	C-O	-6.39	1.15	1.24
1	B	45	MET	C-O	-6.36	1.16	1.24
1	A	18	SER	C-O	-6.35	1.16	1.24
1	A	19	LYS	C-O	-6.35	1.16	1.24
1	B	36	LYS	C-O	-6.31	1.16	1.24
1	A	33	ALA	CA-CB	-6.27	1.43	1.53
1	A	151	PRO	C-O	-6.25	1.16	1.24
1	A	131	PHE	C-O	-6.22	1.16	1.24
1	A	127	ILE	CA-CB	-6.21	1.46	1.54
1	B	81	ARG	C-O	-6.20	1.17	1.24
1	A	34	SER	C-O	-6.20	1.16	1.24
1	A	165	THR	C-O	-6.17	1.16	1.23
1	B	175	ALA	C-O	-6.17	1.17	1.24
1	B	76	ALA	CA-CB	-6.13	1.43	1.53
1	A	36	LYS	C-O	-6.09	1.17	1.24
1	A	81	ARG	C-O	-6.07	1.17	1.24
1	A	111	ARG	C-O	-6.05	1.17	1.24
1	A	144	ILE	C-O	-6.03	1.17	1.24
1	A	17	SER	C-O	-6.02	1.15	1.23
1	B	183	HIS	C-O	-5.97	1.16	1.23
1	B	20	SER	C-O	-5.96	1.16	1.24
1	A	180	TYR	C-O	-5.95	1.16	1.24
1	A	49	LYS	C-O	-5.95	1.17	1.24
1	A	33	ALA	C-O	-5.93	1.17	1.24
1	B	176	ARG	C-O	-5.92	1.17	1.24
1	A	119	ILE	C-O	-5.92	1.17	1.24
1	B	114	MET	C-O	-5.89	1.17	1.24
1	B	44	SER	C-O	-5.84	1.17	1.24
1	A	178	GLN	C-O	-5.83	1.17	1.24
1	A	179	TYR	C-O	-5.83	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	131	PHE	C-O	-5.81	1.17	1.24
1	B	144	ILE	C-O	-5.80	1.17	1.24
1	B	43	GLN	C-O	-5.79	1.17	1.24
1	A	60	SER	C-O	-5.76	1.16	1.23
1	A	126	ALA	CA-CB	-5.72	1.44	1.53
1	B	180	TYR	C-O	-5.70	1.17	1.24
1	B	6	GLY	C-O	-5.70	1.16	1.23
1	A	170	ALA	C-O	-5.67	1.17	1.24
1	A	174	VAL	C-O	-5.67	1.17	1.24
1	B	147	TYR	C-O	-5.65	1.17	1.24
1	B	34	SER	C-O	-5.65	1.17	1.24
1	B	62	ASP	C-O	-5.65	1.17	1.24
1	B	145	SER	C-O	-5.65	1.17	1.24
1	B	53	LYS	C-O	-5.59	1.17	1.24
1	B	113	LEU	C-O	-5.59	1.17	1.24
1	B	123	MET	C-O	-5.58	1.17	1.24
1	B	11	VAL	C-O	-5.58	1.18	1.24
1	B	8	VAL	C-O	-5.54	1.16	1.24
1	B	150	GLY	C-O	-5.51	1.16	1.24
1	B	72	ARG	C-O	-5.51	1.17	1.24
1	A	112	HIS	C-O	-5.51	1.17	1.24
1	A	154	VAL	C-O	-5.51	1.17	1.23
1	A	150	GLY	C-O	-5.48	1.16	1.24
1	A	71	SER	C-O	-5.47	1.17	1.24
1	B	24	GLU	C-O	-5.47	1.16	1.24
1	A	26	LEU	C-O	-5.45	1.17	1.23
1	A	52	ILE	C-O	-5.42	1.18	1.24
1	A	97	ASP	C-O	-5.41	1.17	1.24
1	B	109	GLY	C-O	-5.41	1.17	1.23
1	A	92	GLY	C-O	-5.40	1.17	1.23
1	B	40	ARG	C-O	-5.39	1.17	1.24
1	A	76	ALA	CA-CB	-5.39	1.44	1.53
1	B	15	GLY	C-O	-5.39	1.15	1.24
1	A	129	LYS	C-O	-5.38	1.17	1.24
1	A	98	ARG	C-O	-5.37	1.17	1.24
1	A	142	GLY	C-O	-5.36	1.17	1.23
1	B	141	LYS	C-O	-5.36	1.17	1.24
1	A	140	LEU	C-O	-5.35	1.18	1.24
1	A	35	ARG	C-O	-5.34	1.17	1.24
1	A	139	GLN	C-O	-5.34	1.17	1.24
1	B	84	TRP	C-O	-5.33	1.17	1.23
1	B	174	VAL	C-O	-5.28	1.18	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	160	MET	C-O	-5.24	1.17	1.24
1	B	135	THR	C-O	-5.23	1.16	1.23
1	B	116	GLY	C-O	-5.23	1.17	1.23
1	B	33	ALA	C-O	-5.21	1.18	1.24
1	A	130	LYS	C-O	-5.18	1.18	1.24
1	A	29	SER	C-O	-5.18	1.16	1.23
1	B	91	PHE	C-O	-5.17	1.17	1.24
1	A	123	MET	C-O	-5.16	1.18	1.24
1	B	100	SER	C-O	-5.15	1.17	1.24
1	B	133	ARG	C-O	-5.15	1.17	1.24
1	A	47	ARG	C-O	-5.14	1.17	1.24
1	A	169	TYR	C-O	-5.12	1.18	1.24
1	A	24	GLU	C-O	-5.11	1.17	1.24
1	B	8	VAL	CB-CG2	-5.11	1.35	1.52
1	B	59	LEU	C-O	-5.11	1.16	1.23
1	A	100	SER	C-O	-5.11	1.17	1.24
1	B	21	ALA	C-O	-5.11	1.17	1.24
1	B	119	ILE	C-O	-5.10	1.18	1.24
1	B	29	SER	C-O	-5.09	1.16	1.23
1	A	181	LYS	CE-NZ	-5.08	1.34	1.49
1	A	77	GLY	C-O	-5.08	1.16	1.24
1	A	5	TYR	C-O	-5.07	1.16	1.24
1	B	76	ALA	C-O	-5.06	1.17	1.23
1	A	3	GLY	C-O	-5.04	1.17	1.23
1	B	108	ASN	C-O	-5.03	1.16	1.23
1	A	16	ALA	C-O	-5.01	1.17	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	92	GLY	CA-C-O	-5.54	117.47	122.57
1	A	6	GLY	CA-C-O	-5.02	118.96	122.22

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1443	0	1435	13	0
1	B	1443	0	1435	11	0
2	A	11	0	17	7	0
2	B	11	0	17	3	0
3	A	19	0	26	6	0
4	A	150	0	0	2	0
4	B	123	0	0	3	0
All	All	3200	0	2930	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:GLN:HG2	4:A:324:HOH:O	1.68	0.93
1:A:148:ASN:HA	2:A:201:TAM:H41	1.61	0.82
1:B:101:HIS:O	1:B:103:PRO:HD3	1.93	0.69
1:A:147:TYR:CZ	3:A:202:P6G:H182	2.29	0.68
1:A:101:HIS:HE1	4:A:422:HOH:O	1.79	0.65
1:A:148:ASN:HA	2:A:201:TAM:C4	2.28	0.62
2:A:201:TAM:H11	3:A:202:P6G:H181	1.82	0.62
1:A:123:MET:HE1	3:A:202:P6G:H22	1.88	0.55
2:A:201:TAM:O4	2:A:201:TAM:H22	2.08	0.54
1:B:1:ARG:HB3	4:B:403:HOH:O	2.10	0.52
2:A:201:TAM:O5	2:A:201:TAM:H12	2.09	0.52
1:B:101:HIS:NE2	2:B:201:TAM:H12	2.28	0.49
1:B:134:TRP:HB3	1:B:138:GLN:HB2	1.94	0.49
1:B:101:HIS:CE1	2:B:201:TAM:HO4	2.31	0.49
1:A:147:TYR:CE2	3:A:202:P6G:H182	2.48	0.49
1:B:123:MET:HA	1:B:123:MET:HE2	1.96	0.48
1:B:43:GLN:HG2	4:B:344:HOH:O	2.15	0.46
1:A:22:LYS:HB3	1:A:23:PRO:HD3	1.98	0.45
1:B:43:GLN:HG3	1:B:44:SER:N	2.30	0.45
1:A:148:ASN:O	3:A:202:P6G:H112	2.17	0.44
1:B:22:LYS:HB2	1:B:23:PRO:HD3	2.00	0.43
1:A:147:TYR:O	2:A:201:TAM:H41	2.18	0.43
1:A:151:PRO:HD2	3:A:202:P6G:H32	2.01	0.42
1:A:10:ARG:HB2	1:A:137:GLU:HG2	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:100:SER:OG	2:B:201:TAM:H11	2.21	0.40
1:A:147:TYR:CD2	2:A:201:TAM:H42	2.57	0.40
1:B:166:HIS:HE1	4:B:422:HOH:O	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	183/185 (99%)	181 (99%)	2 (1%)	0	100	100
1	B	183/185 (99%)	179 (98%)	4 (2%)	0	100	100
All	All	366/370 (99%)	360 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	148/148 (100%)	147 (99%)	1 (1%)	76	66
1	B	148/148 (100%)	147 (99%)	1 (1%)	76	66
All	All	296/296 (100%)	294 (99%)	2 (1%)	76	66

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LYS
1	B	129	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	HIS
1	B	43	GLN
1	B	166	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	P6G	A	202	-	18,18,18	2.04	5 (27%)	17,17,17	4.01	6 (35%)
2	TAM	A	201	-	10,10,10	1.16	1 (10%)	12,12,12	2.90	5 (41%)
2	TAM	B	201	-	10,10,10	1.17	2 (20%)	12,12,12	2.79	5 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	P6G	A	202	-	-	9/16/16/16	-
2	TAM	A	201	-	-	3/12/12/12	-
2	TAM	B	201	-	-	4/12/12/12	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	202	P6G	O13-C14	4.61	1.62	1.42
3	A	202	P6G	O13-C12	3.81	1.58	1.42
3	A	202	P6G	O19-C18	3.38	1.59	1.42
3	A	202	P6G	O4-C3	3.25	1.56	1.42
3	A	202	P6G	O7-C6	2.66	1.53	1.42
2	B	201	TAM	C2-C5	2.29	1.56	1.52
2	A	201	TAM	C1-C4	-2.13	1.48	1.52
2	B	201	TAM	C2-C	2.09	1.56	1.53

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	202	P6G	C8-O7-C6	11.80	164.92	113.26
3	A	202	P6G	O7-C8-C9	8.88	150.81	110.35
2	A	201	TAM	C6-C3-C	-6.68	108.09	115.97
2	B	201	TAM	C5-C2-C	5.37	122.30	115.97
2	B	201	TAM	C6-C3-C	-5.22	109.81	115.97
2	A	201	TAM	C3-C-C1	-4.32	102.88	110.50
3	A	202	P6G	O10-C9-C8	-4.25	91.00	110.35
2	A	201	TAM	C4-C1-C	-4.09	111.14	115.97
2	B	201	TAM	C3-C-C2	3.58	116.81	110.50
2	B	201	TAM	C4-C1-C	-3.45	111.90	115.97
3	A	202	P6G	C11-O10-C9	3.28	127.63	113.26
3	A	202	P6G	O16-C15-C14	2.92	123.67	110.35
2	A	201	TAM	C1-C-N	2.65	114.87	108.22
2	A	201	TAM	C3-C-C2	2.47	114.87	110.50
3	A	202	P6G	O1-C2-C3	-2.10	99.44	111.82
2	B	201	TAM	O5-C5-C2	2.05	117.09	111.33

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	201	TAM	N-C-C2-C5
2	A	201	TAM	C-C3-C6-O6
2	B	201	TAM	C2-C-C1-C4
2	B	201	TAM	C3-C-C1-C4
2	B	201	TAM	N-C-C1-C4
2	B	201	TAM	C-C2-C5-O5
3	A	202	P6G	O7-C8-C9-O10
3	A	202	P6G	O16-C17-C18-O19
3	A	202	P6G	O4-C5-C6-O7
3	A	202	P6G	C5-C6-O7-C8
3	A	202	P6G	O13-C14-C15-O16
3	A	202	P6G	O10-C11-C12-O13
3	A	202	P6G	C9-C8-O7-C6
2	A	201	TAM	C3-C-C2-C5
3	A	202	P6G	C6-C5-O4-C3
3	A	202	P6G	C2-C3-O4-C5

There are no ring outliers.

3 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	202	P6G	6	0
2	A	201	TAM	7	0
2	B	201	TAM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	185/185 (100%)	-0.13	0 100 100	9, 13, 20, 26	0
1	B	185/185 (100%)	0.04	2 (1%) 78 84	9, 15, 25, 36	0
All	All	370/370 (100%)	-0.05	2 (0%) 87 91	9, 14, 24, 36	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	ARG	2.4
1	B	133	ARG	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	P6G	A	202	19/19	0.72	0.21	31,34,37,41	0
2	TAM	B	201	11/11	0.77	0.14	20,33,35,38	0
2	TAM	A	201	11/11	0.80	0.15	19,22,25,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.