



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:40 PM UTC

PDB ID : 6WYR / pdb_00006wyr
Title : Crystal structure of anti-Muscle Specific Kinase (MuSK) Fab, MuSK1A
Authors : Vieni, C.; Ekiert, D.
Deposited on : 2020-05-13
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	214	6% 95% 5%
1	M	214	10% 93% 6%
2	H	232	4% 90% 6%
2	I	232	4% 91% 6%
3	A	3	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	B	2	 100%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 14243 atoms, of which 6747 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MuSK1A light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	L	213	Total	C	H	N	O	S	0	11	0
			3281	1039	1612	277	346	7			
1	M	212	Total	C	H	N	O	S	0	6	0
			3209	1018	1574	270	339	8			

- Molecule 2 is a protein called MuSK1A heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	H	222	Total	C	H	N	O	S	0	11	0
			3401	1080	1690	287	336	8			
2	I	226	Total	C	H	N	O	S	0	17	0
			3464	1094	1716	295	349	10			

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	3	Total	C	H	N	O	0	0	0
			73	22	34	2	15			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	2	Total	C	H	N	O	0	0	0
			53	16	25	2	10			

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	L	1	Total	C	H	O	0	0
			10	2	6	2		
5	L	1	Total	C	H	O	0	0
			10	2	6	2		
5	H	1	Total	C	H	O	0	0
			10	2	6	2		
5	H	1	Total	C	H	O	0	0
			10	2	6	2		
5	H	1	Total	C	H	O	0	0
			10	2	6	2		
5	H	1	Total	C	H	O	0	0
			10	2	6	2		
5	M	1	Total	C	H	O	0	0
			10	2	6	2		
5	M	1	Total	C	H	O	0	0
			10	2	6	2		
5	M	1	Total	C	H	O	0	0
			10	2	6	2		
5	M	1	Total	C	H	O	0	0
			10	2	6	2		
5	I	1	Total	C	H	O	0	0
			10	2	6	2		
5	I	1	Total	C	H	O	0	0
			10	2	6	2		
5	I	1	Total	C	H	O	0	0
			10	2	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	I	1	Total	C	H	O	0	0
			10	2	6	2		
5	I	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	101	Total	O	0	0
			101	101		
6	H	197	Total	O	0	0
			197	197		
6	M	91	Total	O	0	0
			91	91		
6	I	213	Total	O	0	0
			213	213		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MuSK1A light chain




- Molecule 1: MuSK1A light chain



- Molecule 2: MuSK1A heavy chain



- Molecule 2: MuSK1A heavy chain



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:

100%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.87Å 71.72Å 106.18Å 90.00° 104.37° 90.00°	Depositor
Resolution (Å)	49.04 – 1.80 49.04 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.04-1.80) 91.5 (49.04-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.63 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.180 , 0.213 0.185 , 0.217	Depositor DCC
R_{free} test set	2005 reflections (2.27%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtriage
Anisotropy	0.246	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 58.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14243	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, BMA, PCA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.11	0/1735	0.31	0/2362
1	M	0.11	0/1682	0.31	0/2293
2	H	0.13	0/1778	0.33	0/2421
2	I	0.14	0/1840	0.35	0/2506
All	All	0.12	0/7035	0.33	0/9582

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1669	1612	1590	5	0
1	M	1635	1574	1574	6	0
2	H	1711	1690	1666	8	0
2	I	1748	1716	1660	12	0
3	A	39	34	34	0	0
4	B	28	25	25	0	0
5	H	20	30	30	0	0
5	I	20	30	30	0	0
5	L	8	12	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	M	16	24	24	0	0
6	H	197	0	0	1	0
6	I	213	0	0	2	0
6	L	101	0	0	0	0
6	M	91	0	0	1	0
All	All	7496	6747	6645	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:58:PRO:O	2:I:80:ARG:NH1	2.36	0.58
2:I:59:TYR:C	2:I:80:ARG:HH12	2.20	0.49
2:I:193:LEU:HD12	2:I:193:LEU:C	2.39	0.48
1:L:13:LEU:HD21	1:L:19:ALA:HB2	1.96	0.47
1:M:107:TRP:HZ2	2:I:107:GLU:OE2	1.98	0.46
1:M:116:VAL:HB	2:I:52:TRP:CD2	2.50	0.46
2:I:165[C]:VAL:HG23	2:I:193:LEU:HD21	1.97	0.46
2:H:193:LEU:C	2:H:193:LEU:HD12	2.41	0.45
1:L:123:ARG:CZ	1:L:163[B]:ARG:HH22	2.29	0.45
1:M:125:THR:HG21	1:M:162:PRO:HB2	1.99	0.45
1:L:163[B]:ARG:HD3	1:L:194:TYR:CE2	2.53	0.44
2:I:59:TYR:O	2:I:80:ARG:NH1	2.51	0.43
2:H:37:TYR:CE2	2:H:108:VAL:HG22	2.53	0.43
1:L:107:TRP:HZ2	2:H:107:GLU:OE2	2.01	0.43
1:M:75:ARG:HB2	1:M:92:SER:O	2.18	0.43
2:I:229:LYS:NZ	6:I:410:HOH:O	2.51	0.43
1:M:163:ARG:NH2	6:M:409:HOH:O	2.51	0.43
2:I:63:GLY:N	2:I:80:ARG:NH1	2.67	0.42
2:H:160:TYR:CE1	2:H:165[B]:VAL:HG23	2.54	0.42
1:M:205:ALA:O	1:M:209:LYS:HG3	2.19	0.42
2:H:69:GLN:NE2	6:H:402:HOH:O	2.45	0.41
2:I:106:ARG:O	2:I:115:SER:HA	2.20	0.41
1:L:116:VAL:HB	2:H:52:TRP:CG	2.55	0.41
2:I:158:LYS:NZ	2:I:186:GLN:OE1	2.54	0.41
2:H:106:ARG:O	2:H:115:SER:HA	2.20	0.41
2:I:95:ARG:NH1	6:I:414:HOH:O	2.54	0.41
2:H:165[B]:VAL:HG22	2:H:215:HIS:CD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	222/214 (104%)	215 (97%)	6 (3%)	1 (0%)	24	14
1	M	216/214 (101%)	209 (97%)	7 (3%)	0	100	100
2	H	229/232 (99%)	225 (98%)	4 (2%)	0	100	100
2	I	241/232 (104%)	236 (98%)	5 (2%)	0	100	100
All	All	908/892 (102%)	885 (98%)	22 (2%)	1 (0%)	48	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	109	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	195/186 (105%)	193 (99%)	2 (1%)	68	64
1	M	190/186 (102%)	185 (97%)	5 (3%)	40	28
2	H	196/194 (101%)	193 (98%)	3 (2%)	57	49
2	I	205/194 (106%)	205 (100%)	0	100	100
All	All	786/760 (103%)	776 (99%)	10 (1%)	61	54

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	29	ASP
1	L	211	LYS
2	H	108	VAL
2	H	224	LYS
2	H	227	GLU
1	M	13	LEU
1	M	125	THR
1	M	129	GLN
1	M	186	GLU
1	M	190	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	37	ASN
1	L	187	GLN
2	H	69	GLN
1	M	168	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PCA	I	1	2	7,8,9	2.30	2 (28%)	9,10,12	2.04	4 (44%)
2	PCA	H	1	2	7,8,9	2.29	2 (28%)	9,10,12	2.09	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	I	1	2	-	0/0/11/13	0/1/1/1
2	PCA	H	1	2	-	0/0/11/13	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	1	PCA	CD-N	4.96	1.46	1.34
2	H	1	PCA	CD-N	4.92	1.46	1.34
2	H	1	PCA	CA-N	3.42	1.50	1.46
2	I	1	PCA	CA-N	3.40	1.50	1.46

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	PCA	CA-N-CD	-3.10	102.96	113.58
2	I	1	PCA	CA-N-CD	-3.10	102.96	113.58
2	H	1	PCA	OE-CD-CG	-2.89	121.57	126.72
2	I	1	PCA	OE-CD-CG	-2.83	121.67	126.72
2	H	1	PCA	CB-CA-N	2.76	110.82	103.24
2	I	1	PCA	CB-CA-N	2.62	110.46	103.24
2	H	1	PCA	CG-CD-N	2.58	114.70	108.39
2	I	1	PCA	CG-CD-N	2.56	114.65	108.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1	3,2	14,14,15	0.36	0	17,19,21	0.45	0
3	NAG	A	2	3	14,14,15	0.19	0	17,19,21	0.48	0
3	BMA	A	3	3	11,11,12	0.51	0	15,15,17	0.82	0
4	NAG	B	1	2,4	14,14,15	0.43	0	17,19,21	0.46	0
4	NAG	B	2	4	14,14,15	0.28	0	17,19,21	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1	3,2	-	2/6/23/26	0/1/1/1
3	NAG	A	2	3	-	2/6/23/26	0/1/1/1
3	BMA	A	3	3	-	2/2/19/22	0/1/1/1
4	NAG	B	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	B	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	2	NAG	O5-C5-C6-O6
3	A	2	NAG	C4-C5-C6-O6
4	B	2	NAG	O5-C5-C6-O6
4	B	2	NAG	C4-C5-C6-O6
3	A	3	BMA	O5-C5-C6-O6
3	A	3	BMA	C4-C5-C6-O6
3	A	1	NAG	C4-C5-C6-O6
3	A	1	NAG	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	H	308	-	3,3,3	0.43	0	2,2,2	0.33	0
5	EDO	M	302	-	3,3,3	0.43	0	2,2,2	0.36	0
5	EDO	M	301	-	3,3,3	0.41	0	2,2,2	0.38	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	I	303	-	3,3,3	0.40	0	2,2,2	0.56	0
5	EDO	L	302	-	3,3,3	0.41	0	2,2,2	0.37	0
5	EDO	I	306	-	3,3,3	0.41	0	2,2,2	0.50	0
5	EDO	L	301	-	3,3,3	0.45	0	2,2,2	0.34	0
5	EDO	H	304	-	3,3,3	0.41	0	2,2,2	0.40	0
5	EDO	M	303	-	3,3,3	0.43	0	2,2,2	0.34	0
5	EDO	I	307	-	3,3,3	0.38	0	2,2,2	0.67	0
5	EDO	M	304	-	3,3,3	0.40	0	2,2,2	0.59	0
5	EDO	I	305	-	3,3,3	0.44	0	2,2,2	0.32	0
5	EDO	H	306	-	3,3,3	0.44	0	2,2,2	0.39	0
5	EDO	H	307	-	3,3,3	0.41	0	2,2,2	0.47	0
5	EDO	I	304	-	3,3,3	0.43	0	2,2,2	0.37	0
5	EDO	H	305	-	3,3,3	0.41	0	2,2,2	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	H	308	-	-	0/1/1/1	-
5	EDO	M	302	-	-	0/1/1/1	-
5	EDO	M	301	-	-	1/1/1/1	-
5	EDO	I	303	-	-	0/1/1/1	-
5	EDO	L	302	-	-	0/1/1/1	-
5	EDO	I	306	-	-	0/1/1/1	-
5	EDO	L	301	-	-	1/1/1/1	-
5	EDO	H	304	-	-	0/1/1/1	-
5	EDO	M	303	-	-	0/1/1/1	-
5	EDO	I	307	-	-	0/1/1/1	-
5	EDO	M	304	-	-	0/1/1/1	-
5	EDO	I	305	-	-	0/1/1/1	-
5	EDO	H	306	-	-	0/1/1/1	-
5	EDO	H	307	-	-	0/1/1/1	-
5	EDO	I	304	-	-	0/1/1/1	-
5	EDO	H	305	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	M	301	EDO	O1-C1-C2-O2
5	L	301	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	L	213/214 (99%)	0.47	12 (5%)	30	29	19, 46, 95, 166	8 (3%)
1	M	212/214 (99%)	0.60	22 (10%)	11	10	19, 50, 96, 185	6 (2%)
2	H	221/232 (95%)	-0.18	9 (4%)	41	41	14, 31, 66, 134	8 (3%)
2	I	225/232 (96%)	-0.24	10 (4%)	39	38	13, 29, 64, 251	10 (4%)
All	All	871/892 (97%)	0.15	53 (6%)	27	25	13, 38, 90, 251	32 (3%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	109	GLY	6.8
2	I	231	CYS	6.5
1	L	107	TRP	6.4
1	M	107	TRP	5.7
2	H	67[A]	TYR	5.7
2	I	112(A)	TYR	5.4
2	I	110	GLY	4.8
2	I	112(B)	GLY	4.4
1	L	235	CYS	4.4
2	H	112(A)	TYR	4.3
2	H	230	SER	4.2
2	I	111(A)	THR	3.9
1	L	2	TYR	3.9
1	M	235	CYS	3.9
1	M	109	GLY	3.8
1	M	234	GLU	3.8
2	I	229	LYS	3.6
2	I	111	ASP	3.6
1	M	127	LEU	3.5
1	L	108	ASP	3.4
2	I	109	ARG	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	L	115	THR	3.3
1	M	20	THR	3.1
2	H	112	GLY	3.1
1	L	114	SER	3.1
1	M	126	VAL	3.0
1	L	96	ALA	3.0
1	M	93	GLY	2.8
2	I	230	SER	2.7
1	L	233	GLY	2.7
1	M	128	GLY	2.7
1	L	125[A]	THR	2.6
2	H	229	LYS	2.6
1	M	13	LEU	2.6
1	M	96	ALA	2.5
1	L	13	LEU	2.5
1	M	114	SER	2.4
1	M	27	LYS	2.4
2	H	109	ARG	2.4
1	M	17	GLN	2.4
1	M	125	THR	2.4
2	I	206	THR	2.3
1	M	11	VAL	2.3
1	M	166	LYS	2.3
1	M	69	SER	2.3
2	H	112(B)	GLY	2.3
1	L	209	LYS	2.3
1	M	97	VAL	2.2
2	H	205	GLY	2.2
2	H	23	CYS	2.1
1	M	147	LYS	2.1
1	M	92	SER	2.1
1	M	226	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	I	1	8/9	0.87	0.16	32,66,80,80	0

Continued on next page...

Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PCA	H	1	8/9	0.89	0.13	27,61,69,75	0


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	1	14/15	-	-	37,59,74,96	0
3	NAG	A	2	14/15	-	-	114,165,316,316	0
3	BMA	A	3	11/12	-	-	142,154,182,185	0
4	NAG	B	1	14/15	-	-	64,92,111,120	0
4	NAG	B	2	14/15	-	-	127,147,173,191	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	M	301	4/4	0.67	0.23	55,66,79,87	0
5	EDO	M	302	4/4	0.77	0.17	43,54,59,64	0
5	EDO	H	308	4/4	0.80	0.30	100,120,124,125	0
5	EDO	L	302	4/4	0.81	0.14	42,50,61,70	0
5	EDO	I	305	4/4	0.86	0.15	31,38,45,54	0
5	EDO	M	304	4/4	0.87	0.27	83,99,111,111	0
5	EDO	M	303	4/4	0.90	0.12	54,66,79,79	0
5	EDO	H	307	4/4	0.90	0.15	29,48,58,58	0
5	EDO	H	306	4/4	0.90	0.12	32,38,46,51	0
5	EDO	L	301	4/4	0.92	0.09	30,38,46,53	0
5	EDO	I	304	4/4	0.94	0.08	34,40,45,50	0
5	EDO	H	305	4/4	0.94	0.15	26,33,61,73	0
5	EDO	I	307	4/4	0.94	0.15	23,41,74,74	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EDO	H	304	4/4	0.95	0.09	33,41,64,64	0
5	EDO	I	306	4/4	0.97	0.07	29,39,48,48	0
5	EDO	I	303	4/4	0.97	0.09	32,40,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.