



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 10:24 PM UTC

PDB ID : 2WZI / pdb_00002wzi
Title : BtGH84 D243N in complex with 5F-oxazoline
Authors : He, Y.; Davies, G.J.
Deposited on : 2009-11-30
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	737	3% 77% 9% 13%
1	B	737	4% 77% 9% 14%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 11649 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called O-GLCNACASE BT_4395.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	640	Total	C	N	O	S	0	11	0
			5273	3379	889	985	20			
1	B	636	Total	C	N	O	S	0	4	0
			5194	3328	877	971	18			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	243	ASN	ASP	engineered mutation	UNP Q89ZI2
B	243	ASN	ASP	engineered mutation	UNP Q89ZI2

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	B	1	Total	Ca	0	0
			1	1		

- Molecule 4 is (3AS,5S,6S,7R,7AR)-5-FLUORO-5-(HYDROXYMETHYL)-2-METHYL-5,6,7,7A-TETRAHYDRO-3AH-PYRANO[3,2-D][1,3]OXAZOLE-6,7-DIOL (CCD ID: 5FN) (formula: C₈H₁₂FO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0
			15	8	1	1	5	0
4	B	1	Total	C	F	N	O	0
			15	8	1	1	5	0

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total 2	Na 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	584	Total 584	O 584	0	0
6	B	534	Total 534	O 534	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: O-GLCNACASE BT 4395



- Molecule 1: O-GLCNACASE BT 4395





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	51.38Å 93.82Å 98.88Å 104.14° 94.06° 103.08°	Depositor
Resolution (Å)	34.33 – 1.90 34.33 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.3 (34.33-1.90) 96.3 (34.33-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.184 , 0.223 0.193 , 0.230	Depositor DCC
R_{free} test set	6620 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	17.4	Xtriage
Anisotropy	0.175	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11649	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 5FN, NA, CA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/5410	0.89	1/7331 (0.0%)
1	B	0.82	2/5326 (0.0%)	0.89	5/7218 (0.1%)
All	All	0.81	2/10736 (0.0%)	0.89	6/14549 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	371	THR	C-O	-5.62	1.17	1.24
1	B	370	VAL	C-N	-5.20	1.26	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	561	ILE	N-CA-C	5.91	116.09	110.53
1	B	89	GLU	N-CA-C	5.67	116.88	108.60
1	B	22	PRO	N-CA-C	5.30	119.19	110.55
1	B	561	ILE	N-CA-C	5.30	115.51	110.42
1	B	209	GLY	N-CA-C	5.24	120.50	114.16
1	B	336	ILE	N-CA-C	5.00	115.56	109.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5273	0	5184	57	0
1	B	5194	0	5104	48	0
2	A	18	0	24	1	0
2	B	12	0	16	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	15	0	12	1	0
4	B	15	0	12	1	0
5	B	2	0	0	0	0
6	A	584	0	0	12	0
6	B	534	0	0	8	0
All	All	11649	0	10352	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:GLN:HG2	6:A:2074:HOH:O	1.32	1.24
1:A:179:PRO:HG3	6:A:2196:HOH:O	1.78	0.83
1:A:431:ASN:H	2:A:1718:GOL:C1	1.99	0.75
1:B:536:GLN:HG2	1:B:590:TYR:CD1	2.21	0.74
1:B:6:GLN:H	1:B:109:GLN:HE22	1.38	0.70
1:B:88:LYS:NZ	6:B:2069:HOH:O	2.25	0.69
1:A:253:LYS:HE3	6:A:2281:HOH:O	1.92	0.69
1:B:253:LYS:HE3	6:B:2251:HOH:O	1.92	0.68
1:B:174:PRO:HD2	6:B:2149:HOH:O	1.94	0.67
1:B:643:ASN:CA	1:B:684:LEU:HD12	2.26	0.65
1:A:308:MET:HA	1:A:335:TYR:O	1.98	0.64
1:A:519:ARG:HD2	1:A:519:ARG:H	1.63	0.63
1:A:337:TRP:C	1:A:337:TRP:CD1	2.75	0.62
1:B:337:TRP:CD1	1:B:337:TRP:C	2.76	0.62
1:A:228[A]:LYS:NZ	6:A:2266:HOH:O	2.35	0.60
1:B:643:ASN:N	1:B:684:LEU:HD12	2.18	0.59
1:B:445:PRO:HB2	6:B:2417:HOH:O	2.03	0.58
1:A:608:GLN:HB2	1:A:615:LEU:HD11	1.85	0.57
1:A:337:TRP:HE1	1:A:372:ASN:HB2	1.70	0.56
1:B:643:ASN:HA	1:B:684:LEU:HD12	1.88	0.55
1:A:337:TRP:C	1:A:337:TRP:HD1	2.14	0.55
1:A:336:ILE:HG21	1:A:369:PHE:CE1	2.41	0.55
1:B:645:GLN:HB3	1:B:708:GLN:HE22	1.70	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:337:TRP:C	1:B:337:TRP:HD1	2.15	0.54
1:B:574:LYS:HE2	1:B:578:GLN:NE2	2.22	0.54
1:A:303:PRO:HD2	6:A:2317:HOH:O	2.07	0.53
1:A:444[B]:GLN:NE2	1:A:448:GLU:OE2	2.42	0.53
1:A:615:LEU:C	1:A:615:LEU:HD12	2.34	0.53
1:A:337:TRP:NE1	1:A:372:ASN:HB2	2.24	0.52
1:B:19:ILE:HG12	1:B:87:GLU:HG3	1.92	0.52
1:A:397:TYR:HE2	1:A:399[B]:THR:HG22	1.75	0.51
1:A:441[A]:MET:SD	1:A:444[A]:GLN:OE1	2.68	0.51
1:A:399[B]:THR:CG2	6:A:2398:HOH:O	2.58	0.51
1:B:557:ALA:HB1	1:B:561:ILE:HB	1.93	0.51
1:B:70:TYR:OH	1:B:89:GLU:OE2	2.28	0.51
1:B:385:TYR:CD2	1:B:406:ALA:HB2	2.47	0.50
1:B:83:LEU:HD23	1:B:83:LEU:C	2.37	0.49
1:A:336:ILE:HB	1:A:369:PHE:HA	1.94	0.49
1:A:242:ASP:OD2	4:A:1719:5FN:N1	2.45	0.49
1:A:608:GLN:HB2	1:A:615:LEU:CD1	2.42	0.49
1:B:591:MET:HG3	1:B:593:HIS:O	2.13	0.49
1:B:308:MET:HA	1:B:335:TYR:O	2.11	0.49
1:A:340:PHE:O	1:A:371:THR:OG1	2.31	0.48
1:A:397:TYR:CE2	1:A:399[B]:THR:HG22	2.49	0.48
1:B:536:GLN:NE2	6:B:2476:HOH:O	2.45	0.48
1:A:341:PRO:O	1:A:342:VAL:C	2.56	0.48
1:B:574:LYS:HE2	1:B:578:GLN:HE22	1.79	0.48
1:A:131:GLY:HA3	1:A:160:THR:O	2.14	0.47
1:B:112:LYS:HE3	6:B:2086:HOH:O	2.14	0.47
1:B:125:PRO:HB3	1:B:392:TRP:CE3	2.50	0.47
1:B:444:GLN:HB3	1:B:445:PRO:HD3	1.95	0.47
1:B:337:TRP:NE1	1:B:372:ASN:HB2	2.30	0.47
1:A:331:LYS:HE3	6:A:2308:HOH:O	2.13	0.46
1:B:214:TRP:HB3	1:B:253:LYS:HD2	1.97	0.46
1:A:462:LYS:HG3	1:A:466:GLU:OE1	2.15	0.46
1:B:72:ARG:HG3	6:B:2049:HOH:O	2.15	0.46
1:B:228[B]:LYS:HE2	1:B:231:GLN:OE1	2.16	0.46
1:A:15:GLN:HB2	1:A:118:GLU:HG2	1.98	0.46
1:B:242:ASP:OD2	4:B:1721:5FN:N1	2.49	0.46
1:B:149:GLN:HB3	1:B:153:TYR:CZ	2.51	0.45
1:B:238:ALA:HA	1:B:276:VAL:O	2.15	0.45
1:B:456:GLU:OE2	1:B:458:LYS:HD2	2.16	0.45
1:A:581:ASN:O	1:A:581:ASN:CG	2.61	0.44
1:B:684:LEU:C	1:B:685:GLN:HG2	2.42	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:THR:HG21	1:A:115:LYS:HE2	1.99	0.44
1:A:170:TYR:HB2	1:A:180:TYR:CE1	2.52	0.44
1:A:27:LEU:HD23	6:A:2042:HOH:O	2.17	0.44
1:A:81:TYR:CE2	1:A:123:ASP:HB3	2.52	0.44
1:A:98:ARG:HG2	1:A:102:TYR:CE2	2.52	0.44
1:A:81:TYR:CZ	1:A:123:ASP:HB3	2.53	0.44
1:A:644:ILE:O	1:A:681:SER:HA	2.17	0.44
1:A:125:PRO:HB3	1:A:392:TRP:CE3	2.53	0.44
1:A:536:GLN:HG2	1:A:590:TYR:CD1	2.53	0.43
1:B:81:TYR:CZ	1:B:123:ASP:HB3	2.53	0.43
1:A:288:ASN:HD22	1:A:289:PRO:HD2	1.83	0.43
1:B:341:PRO:O	1:B:342:VAL:C	2.62	0.43
1:B:337:TRP:CD1	1:B:337:TRP:O	2.71	0.43
1:B:337:TRP:HE1	1:B:372:ASN:HB2	1.83	0.43
1:B:532:LYS:HA	1:B:532:LYS:HD2	1.91	0.42
1:A:65:LYS:HG3	6:A:2031:HOH:O	2.19	0.42
1:A:25:TYR:CZ	1:A:45:LEU:HD13	2.54	0.42
1:A:288:ASN:ND2	6:A:2303:HOH:O	2.52	0.42
1:A:442[A]:ASP:OD1	1:A:443:ILE:N	2.51	0.42
1:B:469:GLN:NE2	6:B:2461:HOH:O	2.51	0.42
1:A:228[B]:LYS:HD3	6:A:2267:HOH:O	2.20	0.42
1:B:293:TYR:CD2	1:B:293:TYR:C	2.98	0.42
1:B:41:LEU:HB2	1:B:104:LEU:HD11	2.01	0.42
1:B:356:TYR:CZ	2:B:1717:GOL:H32	2.55	0.42
1:A:337:TRP:CD1	1:A:337:TRP:O	2.73	0.41
1:A:336:ILE:HG21	1:A:369:PHE:CD1	2.55	0.41
1:A:487:GLU:HG2	6:A:2413:HOH:O	2.20	0.41
1:A:356:TYR:HB3	1:A:399[A]:THR:HG21	2.01	0.41
1:A:462:LYS:HD2	1:A:466:GLU:OE1	2.20	0.41
1:A:238:ALA:HA	1:A:276:VAL:O	2.20	0.41
1:A:355:VAL:O	1:A:399[A]:THR:HG23	2.21	0.41
1:A:385:TYR:CD2	1:A:406:ALA:HB2	2.56	0.41
1:A:62:LYS:O	1:A:62:LYS:HG3	2.21	0.41
1:B:17:LYS:HG2	1:B:18:THR:N	2.35	0.41
1:B:17:LYS:HD3	1:B:118:GLU:OE1	2.21	0.41
1:B:170:TYR:HB2	1:B:180:TYR:CE1	2.56	0.41
1:A:33:ALA:O	1:A:34:ASN:C	2.64	0.40
1:A:293:TYR:CD2	1:A:293:TYR:C	3.00	0.40
1:B:300:LYS:HD3	1:B:300:LYS:HA	1.88	0.40
1:B:489:LYS:N	1:B:490:PRO:CD	2.85	0.40
1:A:532:LYS:HA	1:A:532:LYS:HD2	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	639/737 (87%)	613 (96%)	26 (4%)	0	100	100
1	B	628/737 (85%)	605 (96%)	23 (4%)	0	100	100
All	All	1267/1474 (86%)	1218 (96%)	49 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	572/647 (88%)	568 (99%)	4 (1%)	76	78
1	B	563/647 (87%)	561 (100%)	2 (0%)	84	87
All	All	1135/1294 (88%)	1129 (100%)	6 (0%)	81	84

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ILE
1	A	451	LEU
1	A	462	LYS
1	A	469	GLN
1	B	178	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	708	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	GLN
1	A	252	GLN
1	A	288	ASN
1	A	302	ASN
1	A	529	ASN
1	A	543	GLN
1	A	608	GLN
1	B	109	GLN
1	B	243	ASN
1	B	292	ASN
1	B	529	ASN
1	B	535	GLN
1	B	543	GLN
1	B	578	GLN
1	B	581	ASN
1	B	708	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	1718	-	5,5,5	0.28	0	5,5,5	1.53	1 (20%)
2	GOL	B	1717	-	5,5,5	0.32	0	5,5,5	0.51	0
2	GOL	A	1720	-	5,5,5	0.47	0	5,5,5	0.57	0
2	GOL	B	1716	-	5,5,5	0.39	0	5,5,5	0.66	0
4	5FN	B	1721	-	12,16,16	1.40	1 (8%)	14,25,25	2.22	4 (28%)
2	GOL	A	1716	-	5,5,5	0.39	0	5,5,5	0.67	0
4	5FN	A	1719	-	12,16,16	1.33	1 (8%)	14,25,25	1.57	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1718	-	-	2/4/4/4	-
2	GOL	B	1717	-	-	0/4/4/4	-
2	GOL	A	1720	-	-	2/4/4/4	-
2	GOL	B	1716	-	-	2/4/4/4	-
4	5FN	B	1721	-	-	0/2/34/34	0/2/2/2
2	GOL	A	1716	-	-	0/4/4/4	-
4	5FN	A	1719	-	-	0/2/34/34	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1721	5FN	O23-C5	3.83	1.45	1.37
4	A	1719	5FN	O23-C5	3.38	1.44	1.37

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1721	5FN	C2-N1-C7	6.47	110.96	106.43
4	A	1719	5FN	C2-N1-C7	4.02	109.24	106.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1718	GOL	C3-C2-C1	-2.68	101.95	111.80
4	B	1721	5FN	O23-C1-O1	2.44	113.18	109.37
4	B	1721	5FN	C9-C7-N1	-2.32	123.70	127.05
4	A	1719	5FN	C9-C7-N1	-2.10	124.02	127.05
4	B	1721	5FN	C1-O1-C7	-2.05	104.06	106.88

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1718	GOL	O1-C1-C2-C3
2	A	1720	GOL	O1-C1-C2-C3
2	B	1716	GOL	O1-C1-C2-C3
2	A	1718	GOL	O1-C1-C2-O2
2	A	1720	GOL	O1-C1-C2-O2
2	B	1716	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1718	GOL	1	0
2	B	1717	GOL	1	0
4	B	1721	5FN	1	0
4	A	1719	5FN	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	640/737 (86%)	-0.14	25 (3%)	43 46	8, 21, 58, 89	12 (1%)
1	B	636/737 (86%)	-0.08	29 (4%)	37 40	9, 23, 64, 87	6 (0%)
All	All	1276/1474 (86%)	-0.11	54 (4%)	40 43	8, 21, 60, 89	18 (1%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	582	ALA	4.3
1	B	605	LEU	4.2
1	B	3	VAL	4.1
1	B	19	ILE	4.1
1	A	453	ALA	4.0
1	A	694	PHE	4.0
1	A	46	SER	4.0
1	B	21	LEU	3.7
1	B	454	PHE	3.4
1	A	518	GLY	3.4
1	B	618	PRO	3.3
1	A	605	LEU	3.2
1	A	647	ASN	3.1
1	A	15	GLN	3.1
1	A	454	PHE	3.1
1	B	45	LEU	3.1
1	B	20	ASP	3.1
1	A	457	GLY	3.1
1	A	618	PRO	3.0
1	A	644	ILE	2.9
1	B	692	VAL	2.9
1	B	23	ALA	2.9
1	B	54	GLY	2.9
1	A	709	PHE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	681	SER	2.8
1	B	647	ASN	2.8
1	A	646	ILE	2.6
1	B	638	ILE	2.6
1	A	679	ARG	2.5
1	B	709	PHE	2.5
1	A	616	ILE	2.5
1	B	616	ILE	2.5
1	A	689	VAL	2.5
1	B	685	GLN	2.4
1	A	455	LYS	2.4
1	B	646	ILE	2.4
1	A	456	GLU	2.4
1	B	2	ASN	2.4
1	B	24	VAL	2.4
1	A	179	PRO	2.3
1	B	27	LEU	2.3
1	A	708	GLN	2.2
1	B	708	GLN	2.2
1	B	691	PHE	2.2
1	B	16	ASN	2.2
1	A	53	LYS	2.2
1	B	617	SER	2.1
1	A	451	LEU	2.1
1	B	453	ALA	2.1
1	B	18	THR	2.1
1	A	517	GLU	2.1
1	B	682	ALA	2.1
1	A	611	ALA	2.0
1	B	470	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	1720	6/6	0.85	0.13	34,34,35,35	0
2	GOL	A	1718	6/6	0.86	0.12	25,30,31,33	0
2	GOL	B	1717	6/6	0.90	0.12	25,28,29,30	0
2	GOL	B	1716	6/6	0.93	0.10	22,27,32,33	0
2	GOL	A	1716	6/6	0.94	0.08	27,31,31,32	0
5	NA	B	1719	1/1	0.95	0.07	25,25,25,25	0
3	CA	B	1718	1/1	0.97	0.14	31,31,31,31	1
4	5FN	A	1719	15/15	0.97	0.05	12,13,16,16	0
3	CA	A	1717	1/1	0.97	0.05	30,30,30,30	1
4	5FN	B	1721	15/15	0.98	0.05	11,13,15,16	0
5	NA	B	1720	1/1	0.98	0.05	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.