



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:02 AM UTC

PDB ID : 5WZQ / pdb_00005wzq
Title : Alpha-N-acetylgalactosaminidase NagBb from Bifidobacterium bifidum - quadruple mutant
Authors : Sato, M.; Arakawa, T.; Ashida, H.; Fushinobu, S.
Deposited on : 2017-01-18
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	640	
1	B	640	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10504 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-N-acetylgalactosaminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	610	Total	C	N	O	S	0	0	0
			4820	3034	854	897	35			
1	B	611	Total	C	N	O	S	0	0	0
			4838	3043	859	901	35			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	271	ALA	HIS	engineered mutation	UNP G5ELM1
A	320	ALA	HIS	engineered mutation	UNP G5ELM1
A	322	ALA	ASP	engineered mutation	UNP G5ELM1
A	366	ALA	HIS	engineered mutation	UNP G5ELM1
A	635	HIS	-	expression tag	UNP G5ELM1
A	636	HIS	-	expression tag	UNP G5ELM1
A	637	HIS	-	expression tag	UNP G5ELM1
A	638	HIS	-	expression tag	UNP G5ELM1
A	639	HIS	-	expression tag	UNP G5ELM1
A	640	HIS	-	expression tag	UNP G5ELM1
B	271	ALA	HIS	engineered mutation	UNP G5ELM1
B	320	ALA	HIS	engineered mutation	UNP G5ELM1
B	322	ALA	ASP	engineered mutation	UNP G5ELM1
B	366	ALA	HIS	engineered mutation	UNP G5ELM1
B	635	HIS	-	expression tag	UNP G5ELM1
B	636	HIS	-	expression tag	UNP G5ELM1
B	637	HIS	-	expression tag	UNP G5ELM1
B	638	HIS	-	expression tag	UNP G5ELM1
B	639	HIS	-	expression tag	UNP G5ELM1
B	640	HIS	-	expression tag	UNP G5ELM1

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	452	Total	O	0	0
			452	452		
4	B	380	Total	O	0	0
			380	380		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alpha-N-acetylgalactosaminidase



• Molecule 1: Alpha-N-acetylgalactosaminidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.88Å 127.84Å 176.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	103.51 – 1.90 103.51 – 1.90	Depositor EDS
% Data completeness (in resolution range)	68.6 (103.51-1.90) 68.6 (103.51-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.172 , 0.225 0.172 , 0.225	Depositor DCC
R_{free} test set	3847 reflections (3.42%)	wwPDB-VP
Wilson B-factor (Å ²)	17.9	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 31.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10504	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	4/4955 (0.1%)	1.07	7/6737 (0.1%)
1	B	1.06	3/4973 (0.1%)	1.08	8/6760 (0.1%)
All	All	1.08	7/9928 (0.1%)	1.07	15/13497 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	135	GLU	C-O	-6.26	1.17	1.24
1	B	237	THR	N-CA	5.37	1.52	1.46
1	B	622	HIS	N-CA	5.30	1.52	1.46
1	A	30	SER	CA-C	-5.26	1.46	1.52
1	B	140	ILE	N-CA	5.25	1.50	1.46
1	A	461	ARG	C-O	5.22	1.30	1.24
1	A	365	THR	C-O	5.13	1.29	1.23

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	482	ASP	N-CA-C	8.67	123.62	112.34
1	A	289	GLN	CA-C-N	7.68	127.23	119.24
1	A	289	GLN	C-N-CA	7.68	127.23	119.24
1	B	44	ARG	N-CA-C	6.46	120.10	108.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	485	VAL	CA-C-N	-6.18	113.26	119.56
1	B	485	VAL	C-N-CA	-6.18	113.26	119.56
1	A	446	SER	N-CA-C	-5.77	105.44	112.88
1	B	35	ILE	N-CA-C	5.75	116.07	107.51
1	B	45	SER	N-CA-C	5.43	118.16	110.50
1	A	218	ARG	CB-CA-C	-5.25	101.00	109.72
1	A	38	GLN	N-CA-C	5.22	117.61	109.52
1	A	44	ARG	N-CA-C	5.18	117.06	109.24
1	B	291	GLY	N-CA-C	5.17	119.14	112.83
1	A	289	GLN	O-C-N	5.08	124.90	121.19
1	B	125	ASP	N-CA-C	-5.07	106.35	112.88

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	197	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4820	0	4532	35	0
1	B	4838	0	4550	50	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	452	0	0	5	0
4	B	380	0	0	8	0
All	All	10504	0	9098	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:44:ARG:HD2	4:B:801:HOH:O	1.09	1.23
1:B:44:ARG:CD	4:B:801:HOH:O	1.69	1.13
1:B:2:MET:HE2	1:B:58:ARG:HG2	1.49	0.94
1:A:451:ARG:HD2	4:A:1165:HOH:O	1.71	0.89
1:B:40:THR:HG22	1:B:45:SER:HB3	1.57	0.86
1:A:123:ARG:HB2	4:A:909:HOH:O	1.81	0.79
1:B:44:ARG:NE	4:B:801:HOH:O	1.94	0.78
1:B:14:ASP:CG	1:B:17:THR:HG22	2.10	0.75
1:B:292:LYS:HE2	1:B:292:LYS:H	1.49	0.75
1:B:17:THR:HG23	1:B:19:ARG:H	1.53	0.74
1:B:39:LEU:CD1	1:B:113:VAL:HG22	2.18	0.72
1:A:288:ALA:C	1:A:289:GLN:HG2	2.14	0.71
1:A:1:MET:N	4:A:801:HOH:O	2.24	0.70
1:B:292:LYS:HE2	1:B:292:LYS:N	2.09	0.67
1:B:123:ARG:HG3	1:B:128:ASP:OD2	1.94	0.67
1:A:122:ASP:OD1	1:A:122:ASP:N	2.31	0.64
1:B:37:PRO:HB3	1:B:115:TRP:HD1	1.61	0.64
1:A:19:ARG:HD3	1:A:33:ASP:OD1	2.00	0.62
1:A:96:CYS:HB3	1:A:115:TRP:CH2	2.36	0.61
1:A:286:ASP:OD2	1:A:289:GLN:HG3	2.00	0.61
1:A:280:PRO:HA	1:A:285:TYR:CG	2.36	0.61
1:B:2:MET:HE2	1:B:58:ARG:CG	2.27	0.60
1:B:2:MET:CE	1:B:58:ARG:HG2	2.27	0.60
1:B:2:MET:HG3	4:B:916:HOH:O	2.01	0.59
1:A:37:PRO:HB3	1:A:115:TRP:HD1	1.68	0.58
1:B:537:ASP:OD1	1:B:578:ARG:NH2	2.36	0.58
1:B:614:GLY:O	1:B:631:ASN:HB3	2.05	0.57
1:B:39:LEU:HD12	1:B:113:VAL:HG22	1.85	0.56
1:A:22:PHE:CE1	1:A:225:LEU:HD11	2.40	0.56
1:B:123:ARG:NE	1:B:128:ASP:OD2	2.30	0.55
1:A:286:ASP:CG	1:A:289:GLN:HG3	2.32	0.55
1:B:144:TRP:O	1:B:198:GLY:HA2	2.09	0.53
1:B:58:ARG:HD2	4:B:916:HOH:O	2.10	0.52
1:A:333:HIS:CE1	1:A:368:GLN:HB3	2.45	0.51
1:B:265:LEU:HD21	1:B:591:MET:HG2	1.91	0.51
1:A:435:ASP:O	1:A:436:VAL:HG22	2.10	0.51
1:B:145:PRO:O	1:B:197:ALA:HB1	2.10	0.51
1:B:14:ASP:HB3	1:B:17:THR:CG2	2.40	0.51
1:B:44:ARG:CZ	4:B:801:HOH:O	2.51	0.51
1:A:277:ASN:ND2	1:A:279:GLN:OE1	2.43	0.51
1:A:122:ASP:O	1:A:124:ALA:N	2.45	0.50
1:B:159:PHE:CD1	1:B:167:PRO:HA	2.47	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2:MET:CE	1:B:58:ARG:CG	2.86	0.49
1:A:110:ILE:O	1:A:210:PRO:HG3	2.12	0.49
1:B:588:ARG:NH2	4:B:805:HOH:O	2.46	0.48
1:A:588:ARG:NH2	4:A:809:HOH:O	2.47	0.48
1:B:22:PHE:CD1	1:B:225:LEU:HD21	2.49	0.48
1:B:122:ASP:OD1	1:B:201:TYR:HB2	2.14	0.47
1:B:39:LEU:HD11	1:B:110:ILE:HD11	1.96	0.47
1:B:465:PHE:HB3	1:B:475:THR:HG21	1.97	0.47
1:A:512:VAL:HG23	1:A:514:LEU:HG	1.97	0.46
1:B:5:THR:HG22	1:B:10:MET:HG2	1.97	0.46
1:B:1:MET:O	1:B:1:MET:HG2	2.15	0.46
1:A:289:GLN:O	1:A:292:LYS:HB2	2.16	0.46
1:A:353:ILE:CD1	1:A:427:ILE:HG23	2.46	0.46
1:B:333:HIS:CE1	1:B:368:GLN:HB3	2.51	0.45
1:A:102:ARG:HA	1:A:102:ARG:HD2	1.65	0.45
1:A:527:TRP:CD1	1:A:547:GLY:HA3	2.51	0.45
1:B:14:ASP:CB	1:B:17:THR:HG22	2.47	0.45
1:B:36:GLU:O	1:B:38:GLN:HG2	2.17	0.45
1:B:323:GLY:HA2	1:B:336:TYR:OH	2.17	0.45
1:B:71:PHE:N	1:B:71:PHE:CD1	2.86	0.44
1:A:19:ARG:HD2	1:A:31:GLY:O	2.18	0.44
1:A:280:PRO:HA	1:A:285:TYR:CD2	2.52	0.44
1:B:125:ASP:C	1:B:127:HIS:H	2.27	0.43
1:A:36:GLU:O	1:A:38:GLN:HG3	2.18	0.43
1:A:25:ASP:CG	1:A:227:ASP:OD2	2.62	0.42
1:A:476:SER:HA	1:A:490:PHE:O	2.19	0.42
1:A:59:ILE:HD13	1:B:382:ARG:NH1	2.35	0.42
1:B:14:ASP:HB3	1:B:17:THR:HG22	2.01	0.42
1:B:512:VAL:HG23	1:B:514:LEU:HG	2.01	0.42
1:A:74:PHE:O	1:A:77:ALA:HB3	2.20	0.42
1:A:199:GLY:HA3	1:A:200:PRO:HA	1.81	0.42
1:A:20:PHE:HE1	1:A:22:PHE:CE2	2.37	0.41
1:B:96:CYS:HB3	1:B:115:TRP:CH2	2.54	0.41
1:B:537:ASP:CG	1:B:578:ARG:NH2	2.79	0.41
1:B:123:ARG:HE	1:B:128:ASP:CG	2.24	0.41
1:B:135:GLU:O	1:B:136:GLN:C	2.63	0.41
1:A:10:MET:HE3	4:A:816:HOH:O	2.20	0.40
1:B:3:GLN:HG2	1:B:4:PHE:N	2.36	0.40
1:B:308:THR:O	1:B:312:MET:HG3	2.21	0.40
1:A:531:ARG:HH21	1:A:578:ARG:CD	2.34	0.40
1:A:122:ASP:C	1:A:124:ALA:N	2.80	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:122:ASP:HB2	4:B:1118:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	604/640 (94%)	575 (95%)	23 (4%)	6 (1%)	12	5
1	B	605/640 (94%)	581 (96%)	23 (4%)	1 (0%)	43	36
All	All	1209/1280 (94%)	1156 (96%)	46 (4%)	7 (1%)	21	13

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	ARG
1	A	292	LYS
1	A	76	GLY
1	B	126	ALA
1	A	126	ALA
1	A	436	VAL
1	A	216	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/516 (96%)	480 (97%)	13 (3%)	40	35
1	B	496/516 (96%)	475 (96%)	21 (4%)	26	19
All	All	989/1032 (96%)	955 (97%)	34 (3%)	32	25

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ARG
1	A	42	GLU
1	A	59	ILE
1	A	102	ARG
1	A	122	ASP
1	A	123	ARG
1	A	138	VAL
1	A	192	ASP
1	A	276	THR
1	A	289	GLN
1	A	292	LYS
1	A	493	TYR
1	A	627	GLU
1	B	1	MET
1	B	3	GLN
1	B	21	SER
1	B	32	CYS
1	B	38	GLN
1	B	41	ARG
1	B	43	ASP
1	B	44	ARG
1	B	45	SER
1	B	53	THR
1	B	102	ARG
1	B	112	ARG
1	B	123	ARG
1	B	175	ASP
1	B	204	VAL
1	B	292	LYS
1	B	420	SER
1	B	482	ASP
1	B	493	TYR
1	B	517	LEU
1	B	631	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	277	ASN
1	A	293	ASN
1	A	331	ASN
1	A	383	ASN
1	B	331	ASN
1	B	631	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	702	-	5,5,5	0.29	0	5,5,5	0.51	0
3	GOL	B	702	-	5,5,5	0.66	0	5,5,5	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	702	-	-	2/4/4/4	-
3	GOL	B	702	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	702	GOL	O1-C1-C2-O2
3	A	702	GOL	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	610/640 (95%)	-0.37	12 (1%) 65 69	8, 16, 39, 59	0
1	B	611/640 (95%)	-0.20	11 (1%) 67 71	10, 20, 40, 65	0
All	All	1221/1280 (95%)	-0.29	23 (1%) 66 70	8, 18, 40, 65	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	124	ALA	5.1
1	B	124	ALA	4.7
1	A	126	ALA	4.1
1	A	43	ASP	4.1
1	A	558	ALA	4.0
1	B	43	ASP	3.9
1	A	288	ALA	3.3
1	B	110	ILE	3.2
1	B	44	ARG	3.1
1	B	46	PHE	2.9
1	B	78	ASP	2.9
1	B	1	MET	2.9
1	B	126	ALA	2.8
1	A	1	MET	2.8
1	A	286	ASP	2.8
1	B	79	TYR	2.7
1	A	287	PRO	2.7
1	B	39	LEU	2.7
1	B	630	ALA	2.6
1	A	78	ASP	2.3
1	A	75	ALA	2.3
1	A	46	PHE	2.2
1	A	102	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	702	6/6	0.95	0.07	21,22,23,25	0
3	GOL	B	702	6/6	0.95	0.07	21,22,24,24	0
2	ZN	A	701	1/1	1.00	0.01	11,11,11,11	0
2	ZN	B	701	1/1	1.00	0.01	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.