



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 28, 2026 – 03:41 PM JST

PDB ID : 9WZ2 / pdb_00009wz2
Title : Crystal structure of Cas13a-B complex
Authors : Chen, H.; Huang, L.; Chen, J.; Liu, L.
Deposited on : 2025-09-28
Resolution : 3.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.2
Xtriage (Phenix) : 2.2
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.51


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1131 (3.80-3.60)
Clashscore	190562	1171 (3.80-3.60)
Ramachandran outliers	187476	1129 (3.80-3.60)
Sidechain outliers	187428	1126 (3.80-3.60)
RSRZ outliers	180081	1130 (3.80-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	140	4% 71% 27% .
1	B	140	2% 70% 27% ..
1	C	140	4% 76% 22% .
1	D	140	4% 58% 39% ..
1	E	140	8% 64% 34% .
1	F	140	% 73% 26% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	140	6% 64% 34%
1	H	140	2% 76% 24%
2	I	1159	26% 65% 28% 7%
2	J	1159	13% 69% 23% 7%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 27046 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HicB-like antitoxin of toxin-antitoxin system domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	138	Total	C	N	O	S	0	0	0
			1120	729	172	216	3			
1	B	138	Total	C	N	O	S	0	0	0
			1120	729	172	216	3			
1	C	138	Total	C	N	O	S	0	0	0
			1117	728	172	214	3			
1	D	138	Total	C	N	O	S	0	0	0
			1120	729	172	216	3			
1	E	138	Total	C	N	O	S	0	0	0
			1120	729	172	216	3			
1	F	138	Total	C	N	O	S	0	0	0
			1120	729	172	216	3			
1	G	138	Total	C	N	O	S	0	0	0
			1117	728	172	214	3			
1	H	140	Total	C	N	O	S	0	0	0
			1130	734	174	219	3			

- Molecule 2 is a protein called CRISPR-associated endoribonuclease Cas13a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	1083	Total	C	N	O	S	0	0	0
			9033	5805	1519	1686	23			
2	J	1073	Total	C	N	O	S	0	0	0
			8957	5762	1506	1666	23			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	O	0	0
			4	4		
3	B	7	Total	O	0	0
			7	7		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	4	Total 4	O 4	0	0
3	D	7	Total 7	O 7	0	0
3	E	7	Total 7	O 7	0	0
3	F	3	Total 3	O 3	0	0
3	G	12	Total 12	O 12	0	0
3	H	6	Total 6	O 6	0	0
3	I	21	Total 21	O 21	0	0
3	J	21	Total 21	O 21	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HicB-like antitoxin of toxin-antitoxin system domain-containing protein




- Molecule 1: HicB-like antitoxin of toxin-antitoxin system domain-containing protein




- Molecule 1: HicB-like antitoxin of toxin-antitoxin system domain-containing protein




- Molecule 1: HicB-like antitoxin of toxin-antitoxin system domain-containing protein








- Molecule 1: HicB-like antitoxin of toxin-antitoxin system domain-containing protein




- Molecule 1: HicB-like antitoxin of toxin-antitoxin system domain-containing protein



- Molecule 1: HicB-like antitoxin of toxin-antitoxin system domain-containing protein




- Molecule 1: HicB-like antitoxin of toxin-antitoxin system domain-containing protein




- Molecule 2: CRISPR-associated endoribonuclease Cas13a





● Molecule 2: CRISPR-associated endoribonuclease Cas13a

Chain J:



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	200.16Å 200.16Å 348.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.53 – 3.70 49.53 – 3.70	Depositor EDS
% Data completeness (in resolution range)	85.2 (49.53-3.70) 85.5 (49.53-3.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 3.67Å)	Xtriage
Refinement program	PHENIX dev_svn	Depositor
R, R_{free}	0.322 , 0.330 0.322 , 0.331	Depositor DCC
R_{free} test set	2000 reflections (2.62%)	wwPDB-VP
Wilson B-factor (Å ²)	3.6	Xtriage
Anisotropy	0.615	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.11 , 0.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.28$, $\langle L^2 \rangle = 0.13$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.54	EDS
Total number of atoms	27046	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/1142	0.61	0/1545
1	B	0.24	0/1142	0.63	0/1545
1	C	0.26	0/1139	0.63	0/1541
1	D	0.23	0/1142	0.62	0/1545
1	E	0.25	0/1142	0.61	0/1545
1	F	0.22	0/1142	0.61	0/1545
1	G	0.19	0/1139	0.53	0/1541
1	H	0.21	0/1152	0.63	0/1558
2	I	0.20	0/9184	0.54	2/12293 (0.0%)
2	J	0.18	0/9105	0.52	2/12180 (0.0%)
All	All	0.20	0/27429	0.56	4/36838 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	I	0	2
2	J	0	2
All	All	0	4

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	549	ASN	CA-C-N	5.93	128.01	122.85
2	I	549	ASN	C-N-CA	5.93	128.01	122.85
2	J	544	GLU	CA-C-N	5.48	131.57	121.70
2	J	544	GLU	C-N-CA	5.48	131.57	121.70

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	I	1052	ALA	Peptide
2	I	913	ARG	Peptide
2	J	227	TYR	Peptide
2	J	232	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1120	0	1129	35	0
1	B	1120	0	1129	40	0
1	C	1117	0	1127	34	0
1	D	1120	0	1129	57	0
1	E	1120	0	1129	39	0
1	F	1120	0	1129	30	0
1	G	1117	0	1127	45	0
1	H	1130	0	1137	31	0
2	I	9033	0	9098	262	0
2	J	8957	0	9033	218	0
3	A	4	0	0	0	0
3	B	7	0	0	0	0
3	C	4	0	0	0	0
3	D	7	0	0	0	0
3	E	7	0	0	0	0
3	F	3	0	0	0	0
3	G	12	0	0	0	0
3	H	6	0	0	0	0
3	I	21	0	0	0	0
3	J	21	0	0	0	0
All	All	27046	0	27167	715	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 715 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:867:MET:CE	2:I:1057:ILE:HD11	1.72	1.17
2:J:135:LYS:HA	2:J:138:LEU:HD22	1.27	1.09
2:I:215:LEU:HB3	2:I:222:LYS:HD3	1.37	1.04
2:I:771:HIS:HA	2:I:774:LEU:HD22	1.40	1.03
2:J:135:LYS:HA	2:J:138:LEU:CD2	1.87	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	136/140 (97%)	127 (93%)	9 (7%)	0	100	100
1	B	136/140 (97%)	128 (94%)	5 (4%)	3 (2%)	5	31
1	C	136/140 (97%)	125 (92%)	10 (7%)	1 (1%)	18	50
1	D	136/140 (97%)	132 (97%)	2 (2%)	2 (2%)	8	36
1	E	136/140 (97%)	130 (96%)	5 (4%)	1 (1%)	18	50
1	F	136/140 (97%)	130 (96%)	5 (4%)	1 (1%)	18	50
1	G	136/140 (97%)	133 (98%)	3 (2%)	0	100	100
1	H	138/140 (99%)	130 (94%)	8 (6%)	0	100	100
2	I	1077/1159 (93%)	1026 (95%)	49 (4%)	2 (0%)	43	72
2	J	1061/1159 (92%)	1004 (95%)	56 (5%)	1 (0%)	48	78
All	All	3228/3438 (94%)	3065 (95%)	152 (5%)	11 (0%)	36	65

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	120	ASN
1	B	136	GLY
1	E	120	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	120	ASN
2	J	808	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	125/126 (99%)	125 (100%)	0	100	100
1	B	125/126 (99%)	125 (100%)	0	100	100
1	C	124/126 (98%)	124 (100%)	0	100	100
1	D	125/126 (99%)	125 (100%)	0	100	100
1	E	125/126 (99%)	125 (100%)	0	100	100
1	F	125/126 (99%)	125 (100%)	0	100	100
1	G	124/126 (98%)	124 (100%)	0	100	100
1	H	126/126 (100%)	126 (100%)	0	100	100
2	I	982/1081 (91%)	982 (100%)	0	100	100
2	J	973/1081 (90%)	972 (100%)	1 (0%)	88	88
All	All	2954/3170 (93%)	2953 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	J	1003	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
2	J	696	ASN
2	J	895	ASN
2	I	808	ASN
2	I	719	GLN
2	J	1000	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	138/140 (98%)	0.83	5 (3%)	46	29	13, 22, 39, 58	0
1	B	138/140 (98%)	0.62	3 (2%)	62	39	9, 23, 47, 68	0
1	C	138/140 (98%)	0.78	6 (4%)	40	25	9, 26, 47, 69	0
1	D	138/140 (98%)	0.69	5 (3%)	46	29	10, 25, 43, 78	0
1	E	138/140 (98%)	0.86	11 (7%)	18	14	8, 20, 33, 43	0
1	F	138/140 (98%)	0.67	2 (1%)	73	48	10, 22, 38, 52	0
1	G	138/140 (98%)	0.73	8 (5%)	29	20	8, 19, 40, 63	0
1	H	140/140 (100%)	0.64	3 (2%)	63	40	8, 22, 38, 85	0
2	I	1083/1159 (93%)	1.54	297 (27%)	1	2	9, 45, 117, 194	0
2	J	1073/1159 (92%)	1.08	151 (14%)	6	8	7, 30, 95, 158	0
All	All	3262/3438 (94%)	1.11	491 (15%)	5	7	7, 29, 97, 194	0

The worst 5 of 491 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	J	239	ASN	14.2
2	I	943	ASN	11.3
2	I	1033	ALA	8.5
2	I	949	GLU	8.4
2	I	946	GLU	8.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.