



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 07:31 PM UTC

PDB ID : 1X0P / pdb_00001x0p
Title : Structure of a cyanobacterial BLUF protein, Tll0078
Authors : Kita, A.; Okajima, K.; Morimoto, Y.; Ikeuchi, M.; Miki, K.
Deposited on : 2005-03-27
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	143	85% 14%
1	B	143	78% 19%
1	C	143	76% 22%
1	D	143	80% 17%
1	E	143	77% 21%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	143	2% 71% 27% ...
1	G	143	6% 78% 18% • •
1	H	143	8% 73% 24% •
1	I	143	14% 67% 31% • •
1	J	143	6% 66% 29% • •

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11816 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hypothetical protein Tll0078.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	142	Total	C	N	O	S	0	0	0
			1144	716	204	214	10			
1	B	139	Total	C	N	O	S	0	0	0
			1128	708	200	210	10			
1	C	139	Total	C	N	O	S	0	0	0
			1124	706	200	208	10			
1	D	139	Total	C	N	O	S	0	0	0
			1128	708	200	210	10			
1	E	141	Total	C	N	O	S	0	0	0
			1135	712	200	213	10			
1	F	142	Total	C	N	O	S	0	0	0
			1148	718	204	216	10			
1	G	139	Total	C	N	O	S	0	0	0
			1116	701	199	206	10			
1	H	138	Total	C	N	O	S	0	0	0
			1118	703	199	206	10			
1	I	141	Total	C	N	O	S	0	0	0
			1101	691	194	207	9			
1	J	139	Total	C	N	O	S	0	0	0
			1128	708	200	210	10			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			27	17	4	6		
2	B	1	Total	C	N	O	0	0
			27	17	4	6		
2	C	1	Total	C	N	O	0	0
			27	17	4	6		
2	D	1	Total	C	N	O	0	0
			27	17	4	6		
2	E	1	Total	C	N	O	0	0
			27	17	4	6		
2	F	1	Total	C	N	O	0	0
			27	17	4	6		
2	G	1	Total	C	N	O	0	0
			27	17	4	6		
2	H	1	Total	C	N	O	0	0
			27	17	4	6		
2	I	1	Total	C	N	O	0	0
			27	17	4	6		
2	J	1	Total	C	N	O	0	0
			27	17	4	6		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	39	Total	O	0	0
			39	39		
3	B	32	Total	O	0	0
			32	32		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	30	Total 30	O 30	0	0
3	D	25	Total 25	O 25	0	0
3	E	22	Total 22	O 22	0	0
3	F	35	Total 35	O 35	0	0
3	G	29	Total 29	O 29	0	0
3	H	22	Total 22	O 22	0	0
3	I	20	Total 20	O 20	0	0
3	J	22	Total 22	O 22	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: hypothetical protein Tll0078




- Molecule 1: hypothetical protein Tll0078




- Molecule 1: hypothetical protein Tll0078




- Molecule 1: hypothetical protein Tll0078




- Molecule 1: hypothetical protein Tll0078




- Molecule 1: hypothetical protein Tll0078




- Molecule 1: hypothetical protein Tll0078




- Molecule 1: hypothetical protein Tll0078



- Molecule 1: hypothetical protein Tll0078



- Molecule 1: hypothetical protein Tll0078



R9122
T9125
Q9128
F9132
E9135
L9136
Y9137
Q9138
H9139
S9140
GLN
GLY
SER

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.51Å 109.85Å 169.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 50.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.00) 90.2 (50.00-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 1.91Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.231 , 0.268 0.227 , 0.264	Depositor DCC
R_{free} test set	5752 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	26.3	Xtriage
Anisotropy	0.276	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11816	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.45	0/1164	0.91	3/1562 (0.2%)
1	B	0.43	0/1148	0.90	4/1542 (0.3%)
1	C	0.43	0/1144	0.91	5/1537 (0.3%)
1	D	0.42	0/1148	0.86	3/1542 (0.2%)
1	E	0.39	0/1155	0.83	4/1552 (0.3%)
1	F	0.43	0/1168	0.88	5/1567 (0.3%)
1	G	0.38	0/1136	0.79	2/1528 (0.1%)
1	H	0.44	1/1138 (0.1%)	0.83	3/1529 (0.2%)
1	I	0.37	0/1119	0.77	0/1507
1	J	0.38	0/1148	0.84	3/1542 (0.2%)
All	All	0.41	1/11468 (0.0%)	0.85	32/15408 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	7089	ILE	C-N	-6.11	1.25	1.33

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	ASN	N-CA-C	9.62	122.77	111.71
1	C	2090	ASN	N-CA-C	9.40	122.39	111.11
1	C	2089	ILE	N-CA-C	8.72	119.28	110.82
1	B	1090	ASN	N-CA-C	8.65	121.80	111.33
1	D	3089	ILE	N-CA-C	8.40	118.97	110.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1144	0	1133	13	0
1	B	1128	0	1121	16	0
1	C	1124	0	1117	21	0
1	D	1128	0	1121	23	0
1	E	1135	0	1121	22	0
1	F	1148	0	1137	24	0
1	G	1116	0	1102	26	0
1	H	1118	0	1112	29	0
1	I	1101	0	1069	37	0
1	J	1128	0	1121	32	0
2	A	27	0	19	1	0
2	B	27	0	19	0	0
2	C	27	0	19	1	0
2	D	27	0	19	0	0
2	E	27	0	19	0	0
2	F	27	0	19	0	0
2	G	27	0	19	0	0
2	H	27	0	19	1	0
2	I	27	0	19	1	0
2	J	27	0	19	0	0
3	A	39	0	0	0	0
3	B	32	0	0	1	0
3	C	30	0	0	1	0
3	D	25	0	0	0	0
3	E	22	0	0	0	0
3	F	35	0	0	0	0
3	G	29	0	0	2	0
3	H	22	0	0	0	0
3	I	20	0	0	2	0
3	J	22	0	0	1	0
All	All	11816	0	11344	225	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 225 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:7040:MET:HE2	1:H:7094:ARG:HD3	1.46	0.94
1:G:6098:LEU:HB3	1:G:6106:ILE:HD12	1.53	0.88
1:J:9125:THR:H	1:J:9128:GLN:HE21	1.27	0.81
1:D:3042:CYS:SG	1:D:3098:LEU:HD21	2.21	0.80
1:J:9125:THR:H	1:J:9128:GLN:NE2	1.78	0.80

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	140/143 (98%)	140 (100%)	0	0	100	100
1	B	137/143 (96%)	135 (98%)	2 (2%)	0	100	100
1	C	137/143 (96%)	134 (98%)	3 (2%)	0	100	100
1	D	137/143 (96%)	135 (98%)	2 (2%)	0	100	100
1	E	139/143 (97%)	136 (98%)	2 (1%)	1 (1%)	18	14
1	F	140/143 (98%)	138 (99%)	1 (1%)	1 (1%)	18	14
1	G	137/143 (96%)	135 (98%)	2 (2%)	0	100	100
1	H	136/143 (95%)	131 (96%)	5 (4%)	0	100	100
1	I	139/143 (97%)	130 (94%)	7 (5%)	2 (1%)	9	4
1	J	137/143 (96%)	133 (97%)	3 (2%)	1 (1%)	18	14
All	All	1379/1430 (96%)	1347 (98%)	27 (2%)	5 (0%)	30	27

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	5093	MET
1	I	8093	MET
1	J	9103	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	I	8101	MET
1	E	4141	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/126 (98%)	123 (99%)	1 (1%)	73	80
1	B	123/126 (98%)	122 (99%)	1 (1%)	73	80
1	C	122/126 (97%)	122 (100%)	0	100	100
1	D	123/126 (98%)	122 (99%)	1 (1%)	73	80
1	E	123/126 (98%)	121 (98%)	2 (2%)	55	62
1	F	125/126 (99%)	122 (98%)	3 (2%)	43	47
1	G	120/126 (95%)	118 (98%)	2 (2%)	53	60
1	H	121/126 (96%)	120 (99%)	1 (1%)	73	80
1	I	115/126 (91%)	113 (98%)	2 (2%)	53	60
1	J	123/126 (98%)	119 (97%)	4 (3%)	33	34
All	All	1219/1260 (97%)	1202 (99%)	17 (1%)	59	66

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	9094	ARG
1	J	9109	LEU
1	F	5093	MET
1	G	6095	LEU
1	G	6106	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	8128	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	J	9057	GLN
1	J	9128	GLN
1	E	4120	GLN
1	E	4057	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	E	9154	-	29,29,58	2.68	13 (44%)	42,43,89	2.46	12 (28%)
2	FAD	H	9157	-	29,29,58	2.58	12 (41%)	42,43,89	2.51	13 (30%)
2	FAD	B	9151	-	29,29,58	2.55	11 (37%)	42,43,89	2.55	13 (30%)
2	FAD	G	9156	-	29,29,58	2.57	10 (34%)	42,43,89	2.54	13 (30%)
2	FAD	J	9159	-	29,29,58	2.71	10 (34%)	42,43,89	2.52	13 (30%)
2	FAD	D	9153	-	29,29,58	2.61	13 (44%)	42,43,89	2.46	13 (30%)
2	FAD	A	9150	-	29,29,58	2.48	10 (34%)	42,43,89	2.43	14 (33%)
2	FAD	C	9152	-	29,29,58	2.45	10 (34%)	42,43,89	2.40	11 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	I	9158	-	29,29,58	2.58	10 (34%)	42,43,89	2.49	14 (33%)
2	FAD	F	9155	-	29,29,58	2.73	13 (44%)	42,43,89	2.59	13 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	E	9154	-	-	3/14/14/50	0/3/3/6
2	FAD	H	9157	-	-	3/14/14/50	0/3/3/6
2	FAD	B	9151	-	-	5/14/14/50	0/3/3/6
2	FAD	G	9156	-	-	3/14/14/50	0/3/3/6
2	FAD	J	9159	-	-	2/14/14/50	0/3/3/6
2	FAD	D	9153	-	-	2/14/14/50	0/3/3/6
2	FAD	A	9150	-	-	3/14/14/50	0/3/3/6
2	FAD	C	9152	-	-	1/14/14/50	0/3/3/6
2	FAD	I	9158	-	-	3/14/14/50	0/3/3/6
2	FAD	F	9155	-	-	5/14/14/50	0/3/3/6

The worst 5 of 112 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	9155	FAD	C1'-C2'	-9.40	1.39	1.52
2	J	9159	FAD	C1'-C2'	-8.61	1.40	1.52
2	G	9156	FAD	C1'-C2'	-8.52	1.40	1.52
2	H	9157	FAD	C1'-C2'	-8.09	1.41	1.52
2	E	9154	FAD	C1'-C2'	-7.94	1.41	1.52

The worst 5 of 129 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	9155	FAD	O5'-C5'-C4'	9.26	130.59	111.16
2	G	9156	FAD	O5'-C5'-C4'	9.22	130.52	111.16
2	J	9159	FAD	O5'-C5'-C4'	9.15	130.38	111.16
2	B	9151	FAD	O5'-C5'-C4'	9.14	130.35	111.16
2	E	9154	FAD	O5'-C5'-C4'	8.93	129.91	111.16

There are no chirality outliers.

5 of 30 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	9150	FAD	N10-C1'-C2'-C3'
2	B	9151	FAD	N10-C1'-C2'-O2'
2	B	9151	FAD	N10-C1'-C2'-C3'
2	C	9152	FAD	N10-C1'-C2'-C3'
2	D	9153	FAD	N10-C1'-C2'-C3'

There are no ring outliers.


4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	9157	FAD	1	0
2	A	9150	FAD	1	0
2	C	9152	FAD	1	0
2	I	9158	FAD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand FAD H 9157




Bond lengths




Bond angles





Torsions





Rings




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	142/143 (99%)	0.01	2 (1%) 73 73	16, 24, 33, 39	0
1	B	139/143 (97%)	0.15	2 (1%) 73 73	16, 26, 36, 42	0
1	C	139/143 (97%)	0.15	3 (2%) 62 61	20, 27, 36, 43	0
1	D	139/143 (97%)	0.30	4 (2%) 53 52	20, 29, 36, 43	0
1	E	141/143 (98%)	0.41	6 (4%) 40 39	21, 30, 39, 46	0
1	F	142/143 (99%)	0.15	3 (2%) 63 63	17, 27, 38, 47	0
1	G	139/143 (97%)	0.39	8 (5%) 29 27	19, 29, 42, 48	0
1	H	138/143 (96%)	0.55	12 (8%) 16 15	22, 31, 41, 46	0
1	I	141/143 (98%)	0.96	20 (14%) 6 5	22, 35, 45, 49	0
1	J	139/143 (97%)	0.62	8 (5%) 29 27	22, 32, 42, 46	0
All	All	1399/1430 (97%)	0.37	68 (4%) 35 34	16, 29, 41, 49	0

The worst 5 of 68 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	8103	SER	5.8
1	I	8100	GLU	5.7
1	I	8101	MET	5.2
1	G	6018	TYR	4.6
1	I	8099	GLY	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FAD	I	9158	27/53	0.69	0.15	35,42,47,48	0
2	FAD	D	9153	27/53	0.73	0.15	31,34,41,46	0
2	FAD	E	9154	27/53	0.74	0.15	29,36,43,46	0
2	FAD	B	9151	27/53	0.74	0.14	29,35,38,41	0
2	FAD	F	9155	27/53	0.78	0.14	30,33,38,39	0
2	FAD	J	9159	27/53	0.78	0.13	35,38,43,47	0
2	FAD	H	9157	27/53	0.82	0.12	31,35,42,43	0
2	FAD	G	9156	27/53	0.83	0.12	31,37,43,48	0
2	FAD	C	9152	27/53	0.83	0.11	23,29,35,37	0
2	FAD	A	9150	27/53	0.87	0.11	25,29,34,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FAD I 9158:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD D 9153:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD E 9154:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD B 9151:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD F 9155:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD J 9159:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD H 9157:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD G 9156:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD C 9152:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FAD A 9150:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.