



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 08:24 PM UTC

PDB ID : 2X08 / pdb_00002x08
Title : cytochrome c peroxidase: ascorbate bound to the engineered ascorbate binding site
Authors : Murphy, E.J.; Metcalfe, C.L.; Gumiero, A.; Raven, E.L.; Moody, P.C.E.
Deposited on : 2009-12-07
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYTOCHROME C PEROXIDASE, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	1	0
			2319	1480	387	446	6			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	36	ALA	TYR	engineered mutation	UNP P00431
A	184	ARG	ASN	engineered mutation	UNP P00431
A	191	PHE	TRP	engineered mutation	UNP P00431
A	257	ARG	LYS	conflict	UNP P00431

- Molecule 2 is ASCORBIC ACID (CCD ID: ASC) (formula: C₆H₈O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $\text{C}_{34}\text{H}_{32}\text{FeN}_4\text{O}_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	226	Total O 226 226	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYTOCHROME C PEROXIDASE, MITOCHONDRIAL



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.00Å 74.49Å 106.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.92 – 2.01 41.92 – 2.02	Depositor EDS
% Data completeness (in resolution range)	100.0 (41.92-2.01) 94.1 (41.92-2.02)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.201 , (Not available) 0.348 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	30.5	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 78.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	2600	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ASC, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	3/2390 (0.1%)	1.07	7/3243 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	124	ILE	CA-CB	6.91	1.60	1.54
1	A	154	VAL	CA-CB	5.39	1.60	1.54
1	A	113	VAL	CA-CB	5.12	1.60	1.54

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	124	ILE	CB-CA-C	-6.30	105.09	111.08
1	A	166	ARG	CG-CD-NE	-5.96	98.88	112.00
1	A	280	ALA	CA-C-N	-5.71	113.89	119.78
1	A	280	ALA	C-N-CA	-5.71	113.89	119.78
1	A	166	ARG	NE-CZ-NH2	-5.69	114.08	119.20
1	A	2	THR	CA-C-N	5.36	126.54	119.84
1	A	2	THR	C-N-CA	5.36	126.54	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2319	0	2136	32	0
2	A	12	0	7	2	0
3	A	43	0	30	0	0
4	A	226	0	0	7	0
All	All	2600	0	2173	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:ASP:HB2	4:A:2163:HOH:O	1.35	1.27
1:A:184:ARG:HH22	2:A:1253:ASC:H61	1.40	0.87
1:A:257:ARG:NE	4:A:2193:HOH:O	2.19	0.74
1:A:166:ARG:NH2	1:A:250:GLU:OE1	2.21	0.74
1:A:166:ARG:HH22	1:A:250:GLU:CD	1.97	0.71
1:A:216:ASN:HD22	1:A:222:GLN:HE21	1.42	0.68
1:A:210:ASP:CG	4:A:2165:HOH:O	2.39	0.64
1:A:20:GLN:HE22	1:A:287:LYS:H	1.45	0.64
1:A:44:PRO:HD2	2:A:1253:ASC:O1	1.98	0.63
1:A:216:ASN:HD22	1:A:222:GLN:NE2	1.98	0.62
1:A:98:GLU:OE2	4:A:2084:HOH:O	2.17	0.57
1:A:6:HIS:HE1	4:A:2059:HOH:O	1.88	0.57
1:A:257:ARG:NH2	4:A:2193:HOH:O	2.38	0.56
1:A:257:ARG:CZ	4:A:2193:HOH:O	2.55	0.54
1:A:188:GLU:H	1:A:222:GLN:HE22	1.56	0.54
1:A:53:THR:HG22	1:A:71:TYR:HB2	1.90	0.53
1:A:29:LYS:HE2	1:A:91:PHE:O	2.11	0.50
1:A:99:PHE:O	1:A:102:ILE:HG22	2.12	0.49
1:A:84:GLY:HA3	1:A:184:ARG:HD3	1.97	0.47
1:A:20:GLN:HE22	1:A:287:LYS:N	2.13	0.46
1:A:67:TYR:HA	1:A:130:ARG:HG2	1.98	0.46
1:A:58:ASP:OD1	1:A:60:HIS:HD2	1.99	0.45
1:A:23:TYR:CD2	1:A:23:TYR:C	2.94	0.45
1:A:127:ARG:HG3	1:A:271:GLU:OE1	2.17	0.45
1:A:179:LYS:NZ	1:A:188:GLU:OE1	2.49	0.44
1:A:57:TRP:CD2	1:A:272:ASN:HB3	2.53	0.43
1:A:201:GLU:HG2	1:A:205:ASN:ND2	2.35	0.42
1:A:215:LYS:HD3	1:A:219:ASN:OD1	2.19	0.42
1:A:58:ASP:OD1	1:A:60:HIS:CD2	2.73	0.42
1:A:200:ASN:H	1:A:200:ASN:HD22	1.69	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:MET:HE2	1:A:172:MET:HB2	1.92	0.40
1:A:277:PRO:HD2	1:A:280:ALA:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/293 (100%)	287 (98%)	4 (1%)	1 (0%)	36	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/251 (94%)	231 (98%)	4 (2%)	53	60

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	77	PHE
1	A	200	ASN
1	A	257	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	HIS
1	A	20	GLN
1	A	24	ASN
1	A	159	GLN
1	A	200	ASN
1	A	208	ASN
1	A	222	GLN
1	A	292	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HEM	A	1254	1,4	50,50,50	1.91	9 (18%)	67,82,82	1.51	13 (19%)
2	ASC	A	1253	-	12,12,12	1.79	3 (25%)	17,17,17	2.46	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	1254	1,4	-	2/14/54/54	-
2	ASC	A	1253	-	-	4/6/22/22	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1254	HEM	C3D-C2D	7.05	1.52	1.36
3	A	1254	HEM	FE-NA	5.32	2.12	1.95
2	A	1253	ASC	O4-C1	5.00	1.44	1.36
3	A	1254	HEM	FE-ND	3.77	2.06	1.94
3	A	1254	HEM	CAB-C3B	3.32	1.56	1.47
3	A	1254	HEM	CAC-C3C	3.31	1.56	1.47
3	A	1254	HEM	FE-NC	3.14	2.05	1.95
3	A	1254	HEM	FE-NB	3.03	2.04	1.94
3	A	1254	HEM	CMA-C3A	2.50	1.55	1.50
3	A	1254	HEM	CMC-C2C	2.45	1.55	1.50
2	A	1253	ASC	O4-C4	-2.11	1.42	1.45
2	A	1253	ASC	O1-C1	-2.10	1.17	1.21

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1253	ASC	C1-C2-C3	-6.10	100.29	107.77
3	A	1254	HEM	C4D-ND-C1D	4.66	110.72	105.21
2	A	1253	ASC	C4-C3-C2	4.08	115.78	109.55
3	A	1254	HEM	C3B-C2B-C1B	3.44	108.99	106.41
3	A	1254	HEM	CAA-CBA-CGA	-3.18	105.23	113.67
2	A	1253	ASC	O3-C3-C2	-3.14	124.35	132.40
2	A	1253	ASC	O4-C1-C2	2.98	112.50	109.86
3	A	1254	HEM	C4C-NC-C1C	2.94	110.61	105.82
3	A	1254	HEM	CMD-C2D-C1D	2.88	129.53	125.03
3	A	1254	HEM	CHD-C4C-NC	2.63	127.31	124.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1253	ASC	O2-C2-C1	2.50	128.32	122.22
2	A	1253	ASC	C4-O4-C1	-2.46	106.57	109.27
3	A	1254	HEM	CAD-C3D-C4D	2.42	128.92	124.70
2	A	1253	ASC	O4-C4-C3	-2.27	101.48	103.76
3	A	1254	HEM	C2A-C1A-NA	-2.22	107.69	110.15
3	A	1254	HEM	CHA-C4D-ND	2.18	127.06	124.37
3	A	1254	HEM	O2D-CGD-CBD	2.18	120.88	114.00
3	A	1254	HEM	O1D-CGD-CBD	-2.16	116.25	123.09
3	A	1254	HEM	O1A-CGA-CBA	-2.12	116.38	123.09
3	A	1254	HEM	O2A-CGA-CBA	2.07	120.53	114.00

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1253	ASC	C3-C4-C5-C6
2	A	1253	ASC	C3-C4-C5-O5
2	A	1253	ASC	O4-C4-C5-C6
2	A	1253	ASC	O4-C4-C5-O5
3	A	1254	HEM	CAA-CBA-CGA-O2A
3	A	1254	HEM	CAA-CBA-CGA-O1A

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1253	ASC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/293 (100%)	2.33	165 (56%) 0 0	14, 26, 39, 53	1 (0%)

All (165) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	101	TRP	6.1
1	A	16	TYR	6.1
1	A	194	ALA	5.8
1	A	99	PHE	5.6
1	A	71	TYR	5.1
1	A	2	THR	5.0
1	A	15	SER	5.0
1	A	95	ILE	4.9
1	A	247	ILE	4.9
1	A	10	VAL	4.9
1	A	293	GLY	4.8
1	A	39	TYR	4.7
1	A	13	GLY	4.7
1	A	73	PHE	4.5
1	A	133	THR	4.5
1	A	252	ALA	4.5
1	A	23	TYR	4.4
1	A	3	PRO	4.4
1	A	285	ILE	4.4
1	A	100	PRO	4.3
1	A	7	VAL	4.2
1	A	8	ALA	4.2
1	A	281	PRO	4.1
1	A	276	PHE	4.1
1	A	286	PHE	4.1
1	A	14	ARG	4.0
1	A	213	LEU	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	22	VAL	3.9
1	A	77	PHE	3.9
1	A	4	LEU	3.9
1	A	248	VAL	3.8
1	A	256	ASP	3.8
1	A	98	GLU	3.8
1	A	96	HIS	3.7
1	A	294	LEU	3.7
1	A	134	PRO	3.7
1	A	275	THR	3.7
1	A	280	ALA	3.7
1	A	207	LEU	3.6
1	A	291	GLU	3.6
1	A	245	LEU	3.6
1	A	19	PHE	3.5
1	A	40	ILE	3.5
1	A	182	LEU	3.5
1	A	69	GLY	3.5
1	A	128	CYS	3.5
1	A	137	THR	3.5
1	A	265	ALA	3.4
1	A	277	PRO	3.4
1	A	287	LYS	3.4
1	A	116	VAL	3.4
1	A	250	GLU	3.4
1	A	94	PRO	3.4
1	A	206	LEU	3.4
1	A	38	ASN	3.3
1	A	89	PHE	3.3
1	A	102	ILE	3.3
1	A	24	ASN	3.3
1	A	20	GLN	3.3
1	A	129	GLY	3.2
1	A	258	PHE	3.2
1	A	217	ASP	3.2
1	A	254	ASP	3.2
1	A	70	THR	3.2
1	A	211	TRP	3.1
1	A	136	ASP	3.1
1	A	88	GLY	3.1
1	A	292	GLN	3.1
1	A	18	ASP	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	244	TYR	3.1
1	A	97	LYS	3.1
1	A	198	PHE	3.0
1	A	259	PHE	3.0
1	A	17	GLU	3.0
1	A	246	SER	3.0
1	A	283	PRO	3.0
1	A	242	PRO	3.0
1	A	289	LEU	2.9
1	A	72	ARG	2.9
1	A	103	SER	2.9
1	A	126	TRP	2.9
1	A	193	ALA	2.9
1	A	212	LYS	2.9
1	A	131	VAL	2.8
1	A	249	LYS	2.8
1	A	218	ALA	2.8
1	A	251	TYR	2.8
1	A	226	LYS	2.7
1	A	26	ILE	2.7
1	A	132	ASP	2.7
1	A	108	PHE	2.7
1	A	87	ASN	2.7
1	A	45	VAL	2.7
1	A	284	PHE	2.7
1	A	263	SER	2.7
1	A	34	ASP	2.7
1	A	210	ASP	2.7
1	A	28	LEU	2.6
1	A	233	PRO	2.6
1	A	9	SER	2.6
1	A	274	ILE	2.6
1	A	36	ALA	2.6
1	A	91	PHE	2.6
1	A	255	GLN	2.6
1	A	190	PRO	2.6
1	A	214	GLU	2.5
1	A	169	VAL	2.5
1	A	270	LEU	2.5
1	A	221	GLU	2.5
1	A	30	LEU	2.5
1	A	170	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	110	LEU	2.5
1	A	215	LYS	2.5
1	A	208	ASN	2.5
1	A	5	VAL	2.4
1	A	85	LEU	2.4
1	A	120	GLN	2.4
1	A	42	TYR	2.4
1	A	92	LEU	2.4
1	A	138	THR	2.4
1	A	203	TYR	2.3
1	A	253	ASN	2.3
1	A	279	ASP	2.3
1	A	197	VAL	2.3
1	A	195	ASN	2.3
1	A	228	GLY	2.3
1	A	33	ASP	2.3
1	A	223	TRP	2.3
1	A	6	HIS	2.3
1	A	290	GLU	2.3
1	A	53	THR	2.3
1	A	288	THR	2.3
1	A	220	ASN	2.3
1	A	178	GLY	2.3
1	A	172	MET	2.3
1	A	80	PRO	2.2
1	A	122	PRO	2.2
1	A	219	ASN	2.2
1	A	12	LYS	2.2
1	A	65	GLY	2.2
1	A	144	LEU	2.2
1	A	257	ARG	2.2
1	A	268	LYS	2.2
1	A	272	ASN	2.2
1	A	261	ASP	2.2
1	A	113	VAL	2.2
1	A	86	GLN	2.2
1	A	47	VAL	2.1
1	A	264	LYS	2.1
1	A	35	GLU	2.1
1	A	241	ASP	2.1
1	A	93	GLU	2.1
1	A	135	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	67	TYR	2.1
1	A	266	PHE	2.1
1	A	123	LYS	2.1
1	A	151	ALA	2.1
1	A	171	LEU	2.1
1	A	186	GLY	2.1
1	A	46	LEU	2.0
1	A	236	TYR	2.0
1	A	243	LYS	2.0
1	A	57	TRP	2.0
1	A	127	ARG	2.0
1	A	21	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ASC	A	1253	12/12	0.55	0.32	13,25,34,35	12
3	HEM	A	1254	43/43	0.91	0.12	17,23,25,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HEM A 1254:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.