



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 12:48 PM UTC

PDB ID : 1X11 / pdb_00001x11
Title : X11 PTB DOMAIN
Authors : Lee, C.-H.; Zhang, Z.; Kuriyan, J.
Deposited on : 1997-07-28
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	172	% 42% 30% 7% 22%
1	B	172	% 36% 30% 5% 29%
2	C	13	62% 31% 8%
2	D	13	62% 31% 8%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3196 atoms, of which 814 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called X11.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	135	Total	C	H	N	O	S	Se	0	0	0
			1288	661	237	180	200	1	9			
1	B	122	Total	C	H	N	O	S	Se	0	0	0
			1151	602	204	161	175	1	8			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	354	MSE	MET	modified residue	UNP Q02410
A	355	MSE	MET	modified residue	UNP Q02410
A	366	MSE	MET	modified residue	UNP Q02410
A	385	MSE	MET	modified residue	UNP Q02410
A	408	MSE	MET	modified residue	UNP Q02410
A	409	MSE	MET	modified residue	UNP Q02410
A	429	MSE	MET	modified residue	UNP Q02410
A	458	MSE	MET	modified residue	UNP Q02410
B	354	MSE	MET	modified residue	UNP Q02410
B	355	MSE	MET	modified residue	UNP Q02410
B	366	MSE	MET	modified residue	UNP Q02410
B	385	MSE	MET	modified residue	UNP Q02410
B	408	MSE	MET	modified residue	UNP Q02410
B	409	MSE	MET	modified residue	UNP Q02410
B	429	MSE	MET	modified residue	UNP Q02410
B	458	MSE	MET	modified residue	UNP Q02410

- Molecule 2 is a protein called 13-MER PEPTIDE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	12	Total	C	H	N	O	0	0	0
			132	71	25	16	20			
2	D	13	Total	C	H	N	O	0	0	0
			142	76	26	17	23			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	83	Total	H	O	0	0
			249	166	83		
3	C	8	Total	H	O	0	0
			24	16	8		
3	B	55	Total	H	O	0	0
			165	110	55		
3	D	15	Total	H	O	0	0
			45	30	15		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: X11



• Molecule 1: X11



• Molecule 2: 13-MER PEPTIDE



• Molecule 2: 13-MER PEPTIDE





4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	74.60Å 74.60Å 155.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.50 6.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	82.7 (6.00-2.50) 85.9 (6.00-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 2.39Å)	Xtriage
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.214 , 0.304 0.235 , 0.319	Depositor DCC
R_{free} test set	1638 reflections (9.99%)	wwPDB-VP
Wilson B-factor (Å ²)	45.9	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.16 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3196	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	6/1054 (0.6%)	1.10	6/1406 (0.4%)
1	B	0.95	7/949 (0.7%)	1.00	2/1267 (0.2%)
2	C	0.47	0/111	0.82	0/149
2	D	0.61	0/120	1.09	1/161 (0.6%)
All	All	0.91	13/2234 (0.6%)	1.05	9/2983 (0.3%)

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	460	CYS	CB-SG	-11.37	1.43	1.81
1	A	460	CYS	CB-SG	-11.30	1.44	1.81
1	A	429	MSE	SE-CE	-6.36	1.76	1.95
1	A	323	MSE	SE-CE	-6.32	1.76	1.95
1	B	354	MSE	SE-CE	-6.31	1.76	1.95

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	456	TYR	N-CA-C	7.21	121.79	112.12
2	D	11	PHE	N-CA-C	6.74	120.61	112.38
1	B	392	ILE	N-CA-C	6.25	116.92	108.17
1	A	466	GLU	N-CA-C	-6.22	104.96	112.54
1	B	412	PRO	N-CA-C	-5.91	101.87	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1051	237	1052	56	0
1	B	947	204	950	43	0
2	C	107	25	94	7	0
2	D	116	26	100	4	0
3	A	83	166	0	4	0
3	B	55	110	0	2	0
3	C	8	16	0	1	0
3	D	15	30	0	0	0
All	All	2382	814	2196	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 102 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:393:LEU:HD21	1:B:396:ARG:HG2	1.56	0.87
1:B:429:MSE:HB3	1:B:458:MSE:HE1	1.60	0.81
1:A:397:ILE:HG21	1:A:428:LEU:HD11	1.63	0.80
1:B:391:PHE:CE2	1:B:393:LEU:HD22	2.17	0.80
1:A:395:GLN:HE21	1:A:396:ARG:NH1	1.87	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/172 (75%)	120 (93%)	8 (6%)	1 (1%)	16	31
1	B	114/172 (66%)	107 (94%)	5 (4%)	2 (2%)	6	12

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	10/13 (77%)	9 (90%)	1 (10%)	0	100	100
2	D	11/13 (85%)	11 (100%)	0	0	100	100
All	All	264/370 (71%)	247 (94%)	14 (5%)	3 (1%)	11	22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	457	LYS
1	A	367	ALA
1	B	372	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/140 (81%)	101 (89%)	12 (11%)	6	14
1	B	100/140 (71%)	90 (90%)	10 (10%)	7	15
2	C	11/12 (92%)	11 (100%)	0	100	100
2	D	12/12 (100%)	12 (100%)	0	100	100
All	All	236/304 (78%)	214 (91%)	22 (9%)	8	18

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	358	GLN
1	B	395	GLN
1	B	387	GLU
1	B	396	ARG
1	A	419	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	405	GLN
1	B	469	GLN
1	A	477	GLN
1	A	490	ASN
2	C	1	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	126/172 (73%)	-0.08	1 (0%) 82 80	18, 31, 57, 92	0
1	B	114/172 (66%)	-0.08	2 (1%) 67 64	18, 32, 58, 70	0
2	C	12/13 (92%)	-0.07	0 100 100	23, 35, 48, 53	0
2	D	13/13 (100%)	-0.02	0 100 100	23, 35, 50, 62	0
All	All	265/370 (71%)	-0.08	3 (1%) 78 75	18, 32, 58, 92	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	372	LYS	2.5
1	B	494	PRO	2.5
1	A	454	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.