



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:43 AM UTC

PDB ID : 1X13 / pdb_00001x13
Title : Crystal structure of E. coli transhydrogenase domain I
Authors : Johansson, T.; Oswald, C.; Pedersen, A.; Tornroth, S.; Okvist, M.; Karlsson, B.G.; Rydstrom, J.; Kregel, U.
Deposited on : 2005-03-31
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	85% 5% • 8%
1	B	401	2% 82% 8% • 9%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5946 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD(P) transhydrogenase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	0	0	0
			2745	1741	465	527	12			
1	B	364	Total	C	N	O	S	0	0	0
			2721	1724	464	522	11			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	994	MET	-	expression tag	UNP P07001
A	995	HIS	-	expression tag	UNP P07001
A	996	HIS	-	expression tag	UNP P07001
A	997	HIS	-	expression tag	UNP P07001
A	998	HIS	-	expression tag	UNP P07001
A	999	HIS	-	expression tag	UNP P07001
A	1000	HIS	-	expression tag	UNP P07001
A	1001	GLY	-	expression tag	UNP P07001
B	994	MET	-	expression tag	UNP P07001
B	995	HIS	-	expression tag	UNP P07001
B	996	HIS	-	expression tag	UNP P07001
B	997	HIS	-	expression tag	UNP P07001
B	998	HIS	-	expression tag	UNP P07001
B	999	HIS	-	expression tag	UNP P07001
B	1000	HIS	-	expression tag	UNP P07001
B	1001	GLY	-	expression tag	UNP P07001


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	227	Total	O	0	0
			227	227		
2	B	253	Total	O	0	0
			253	253		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NAD(P) transhydrogenase subunit alpha



- Molecule 1: NAD(P) transhydrogenase subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	38.76Å 66.90Å 76.30Å 67.11° 80.66° 81.23°	Depositor
Resolution (Å)	30.00 – 1.90 30.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-1.90) 94.3 (30.00-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.179 , 0.224 0.188 , 0.233	Depositor DCC
R_{free} test set	3836 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	22.4	Xtriage
Anisotropy	0.366	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5946	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/2787	0.90	3/3785 (0.1%)
1	B	0.84	0/2764	0.93	1/3756 (0.0%)
All	All	0.79	0/5551	0.91	4/7541 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1061	ASN	N-CA-C	5.63	118.96	111.75
1	A	1342	LYS	CA-C-N	5.36	131.35	121.70
1	A	1342	LYS	C-N-CA	5.36	131.35	121.70
1	A	1248	GLU	N-CA-C	5.31	120.62	113.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2745	0	2804	7	0
1	B	2721	0	2772	23	1
2	A	227	0	0	0	0
2	B	253	0	0	6	1
All	All	5946	0	5576	30	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1100:GLU:N	1:B:1100:GLU:OE1	2.31	0.64
1:B:1048:LYS:HG2	2:B:73:HOH:O	1.97	0.64
1:B:1115:MET:HE2	1:B:1331:LEU:HD23	1.83	0.60
1:B:1375:VAL:HG12	1:B:1376:SER:N	2.17	0.58
1:B:1100:GLU:CD	1:B:1100:GLU:H	2.13	0.56
1:A:1341:GLU:O	1:A:1342:LYS:C	2.52	0.53
1:B:1041:GLN:NE2	2:B:410:HOH:O	2.43	0.52
1:A:1239:MET:HA	1:A:1239:MET:HE2	1.93	0.49
1:B:1141:ALA:HB2	1:B:1284:LEU:HD11	1.95	0.48
1:A:1342:LYS:N	1:A:1343:ASP:HB3	2.29	0.48
1:B:1375:VAL:HG12	1:B:1376:SER:H	1.81	0.46
1:B:1156:GLN:HE21	1:B:1158:THR:HG22	1.82	0.45
1:B:1185:ASN:ND2	2:B:86:HOH:O	2.50	0.45
1:B:1256:ALA:H	1:B:1291:ASN:HD21	1.64	0.45
1:B:1254:THR:HG21	1:B:1291:ASN:HB2	2.00	0.44
1:B:1120:ARG:HB2	1:B:1375:VAL:CG1	2.48	0.44
1:A:1169:MET:HA	1:A:1192:ARG:O	2.17	0.44
1:B:1120:ARG:O	1:B:1375:VAL:HG13	2.17	0.44
1:A:1338:LEU:CD1	1:A:1355:ILE:HD12	2.49	0.43
1:B:1156:GLN:NE2	2:B:319:HOH:O	2.52	0.43
1:A:1239:MET:HE2	1:A:1239:MET:CA	2.48	0.42
1:B:1251:ILE:HA	1:B:1280:VAL:O	2.20	0.41
1:B:1156:GLN:CD	2:B:319:HOH:O	2.63	0.41
1:B:1287:GLN:C	1:B:1288:ASN:HD22	2.28	0.41
1:A:1284:LEU:HD22	1:A:1315:LEU:CD2	2.50	0.41
1:B:1119:PRO:HG2	1:B:1354:VAL:HA	2.02	0.41
1:B:1048:LYS:HG3	2:B:254:HOH:O	2.21	0.41
1:B:1235:ILE:O	1:B:1239:MET:HG2	2.22	0.40
1:B:1185:ASN:HD22	1:B:1185:ASN:HA	1.72	0.40
1:B:1149:PHE:CE2	1:B:1151:ARG:HB2	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:999:HIS:NE2	2:B:261:HOH:O[1_565]	2.12	0.08

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/401 (91%)	357 (98%)	5 (1%)	2 (0%)	24	16
1	B	360/401 (90%)	352 (98%)	7 (2%)	1 (0%)	36	29
All	All	724/802 (90%)	709 (98%)	12 (2%)	3 (0%)	30	22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1342	LYS
1	A	1343	ASP
1	B	1375	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/317 (92%)	271 (93%)	20 (7%)	14	7
1	B	289/317 (91%)	280 (97%)	9 (3%)	35	29
All	All	580/634 (92%)	551 (95%)	29 (5%)	22	14

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1021	LYS
1	A	1048	LYS
1	A	1059	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1083	LEU
1	A	1101	LEU
1	A	1104	LYS
1	A	1122	SER
1	A	1153	PHE
1	A	1199	GLU
1	A	1207	MET
1	A	1228	LYS
1	A	1241	LEU
1	A	1257	LEU
1	A	1265	LYS
1	A	1284	LEU
1	A	1287	GLN
1	A	1315	LEU
1	A	1338	LEU
1	A	1343	ASP
1	A	1346	ILE
1	B	1041	GLN
1	B	1100	GLU
1	B	1162	LYS
1	B	1203	GLN
1	B	1215	ASP
1	B	1231	SER
1	B	1261	LYS
1	B	1337	LEU
1	B	1345	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1052	GLN
1	A	1084	ASN
1	A	1135	ASN
1	A	1156	GLN
1	A	1205	GLN
1	A	1287	GLN
1	A	1288	ASN
1	A	1330	ASN
1	B	1073	ASN
1	B	1084	ASN
1	B	1156	GLN
1	B	1185	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1203	GLN
1	B	1288	ASN
1	B	1291	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	368/401 (91%)	0.23	6 (1%) 70 74	17, 31, 47, 55	0
1	B	364/401 (90%)	0.04	10 (2%) 56 60	15, 27, 45, 61	0
All	All	732/802 (91%)	0.13	16 (2%) 62 66	15, 29, 47, 61	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	999	HIS	3.9
1	B	1377	ALA	3.9
1	B	1375	VAL	3.5
1	A	1342	LYS	3.3
1	B	1214	LEU	2.8
1	B	1376	SER	2.7
1	B	1259	PRO	2.7
1	B	1372	PRO	2.7
1	A	1344	GLY	2.4
1	A	1258	ILE	2.4
1	A	1157	ILE	2.3
1	A	1078	ASP	2.3
1	B	1260	GLY	2.2
1	B	1258	ILE	2.1
1	B	1231	SER	2.0
1	A	1214	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.