



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 07:35 AM UTC

PDB ID : 2X1D / pdb_00002x1d
Title : The crystal structure of mature acyl coenzyme A:isopenicillin N acyltransferase from *Penicillium chrysogenum*
Authors : Bokhove, M.; Yoshida, H.; Hensgens, C.M.H.; van der Laan, J.M.; Sutherland, J.D.; Dijkstra, B.W.
Deposited on : 2009-12-23
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1141 (1.64-1.64)
Clashscore	190562	1171 (1.64-1.64)
Ramachandran outliers	187476	1151 (1.64-1.64)
Sidechain outliers	187428	1150 (1.64-1.64)
RSRZ outliers	180081	1141 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	357	6% 94% 5%
1	B	357	8% 90% 8%
1	C	357	10% 93% 6%
1	D	357	5% 93% 6%

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 12327 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACYL-COENZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	354	Total	C	N	O	S	0	8	0
			2848	1784	513	538	13			
1	B	352	Total	C	N	O	S	0	10	0
			2856	1790	513	538	15			
1	C	351	Total	C	N	O	S	0	8	0
			2835	1777	509	534	15			
1	D	351	Total	C	N	O	S	0	8	0
			2828	1776	503	532	17			

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	B	2	Total	Cl	0	0
			2	2		
4	C	1	Total	Cl	0	0
			1	1		
4	D	1	Total	Cl	0	0
			1	1		

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is (4S,5S)-1,2-DITHIANE-4,5-DIOL (CCD ID: D1D) (formula: C₄H₈O₂S₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	D	1	Total	C	O	S	0	0
			8	4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	202	Total 202	O 202	0	0
7	B	260	Total 260	O 260	0	0
7	C	226	Total 226	O 226	0	0
7	D	221	Total 221	O 221	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ACYL-COENZYME




• Molecule 1: ACYL-COENZYME



• Molecule 1: ACYL-COENZYME



• Molecule 1: ACYL-COENZYME





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	198.71Å 68.22Å 146.96Å 90.00° 128.64° 90.00°	Depositor
Resolution (Å)	34.10 – 1.64 34.10 – 1.64	Depositor EDS
% Data completeness (in resolution range)	98.2 (34.10-1.64) 98.2 (34.10-1.64)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 1.64Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.179 , 0.207 0.190 , 0.215	Depositor DCC
R_{free} test set	9288 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.134	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12327	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, CSD, GOL, PO4, D1D, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/2898	0.79	1/3913 (0.0%)
1	B	0.64	0/2905	0.79	0/3923
1	C	0.62	0/2884	0.79	0/3891
1	D	0.58	0/2877	0.79	0/3884
All	All	0.60	0/11564	0.79	1/15611 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	152	ILE	N-CA-C	5.50	118.28	112.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2848	0	2785	9	0
1	B	2856	0	2799	23	0
1	C	2835	0	2777	8	0
1	D	2828	0	2774	14	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	10	0	0	0	0
3	A	6	0	8	0	0
4	A	1	0	0	0	0
4	B	2	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	B	4	0	3	1	0
5	C	4	0	3	0	0
5	D	4	0	3	0	0
6	D	8	0	8	0	0
7	A	202	0	0	0	0
7	B	260	0	0	1	0
7	C	226	0	0	1	0
7	D	221	0	0	0	0
All	All	12327	0	11160	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:62:TRP:CZ2	1:B:179[B]:VAL:HG12	2.02	0.95
1:B:62:TRP:HZ2	1:B:179[B]:VAL:HG12	1.38	0.86
1:D:214:MET:HE2	1:D:242[B]:MET:HE2	1.65	0.78
1:C:94:TYR:HA	1:C:97:LYS:HB2	1.70	0.73
1:C:96:LEU:HG	1:C:174:LEU:HD22	1.73	0.70
1:B:256:GLU:OE2	5:B:1358:ACT:H2	1.92	0.69
1:B:179[B]:VAL:HG11	1:B:204:GLN:HB3	1.75	0.66
1:B:177[B]:THR:O	1:B:177[B]:THR:HG22	1.97	0.64
1:B:179[B]:VAL:HG11	1:B:204:GLN:CB	2.28	0.64
1:A:236:LEU:HD23	1:A:242:MET:HG3	1.80	0.63
1:D:179[B]:VAL:HG11	1:D:204:GLN:CB	2.30	0.62
1:A:95:GLY:HA2	1:A:174:LEU:CD1	2.34	0.56
1:B:179[B]:VAL:CG1	1:B:204:GLN:HB3	2.37	0.55
1:B:236:LEU:HD23	1:B:242[B]:MET:HG3	1.87	0.55
1:C:42:LYS:HE2	1:C:96:LEU:HD22	1.92	0.51
1:A:203:GLU:HG3	1:B:232:ARG:HD2	1.92	0.51
1:A:350:ARG:HG3	1:A:355:ALA:HB3	1.93	0.50
1:C:350:ARG:HG3	1:C:355:ALA:HB3	1.93	0.50
1:B:207:MET:HG3	1:B:226:PHE:CD1	2.47	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:67:GLU:OE1	1:D:70:ARG:NH2	2.42	0.50
1:B:60[A]:GLU:HG3	7:B:2061:HOH:O	2.12	0.49
1:A:252:HIS:HB3	1:A:256:GLU:HG3	1.95	0.49
1:A:95:GLY:HA2	1:A:174:LEU:HD13	1.95	0.49
1:C:45:GLU:O	1:C:49:GLN:HG2	2.14	0.48
1:B:94:TYR:HA	1:B:97:LYS:HB2	1.96	0.47
1:B:112:ASN:HB3	7:C:2112:HOH:O	2.14	0.46
1:A:36:ILE:HG23	1:A:96:LEU:CD1	2.45	0.46
1:D:241:ARG:HD2	1:D:276:LEU:HD13	1.98	0.46
1:D:275:LEU:HB3	1:D:289:LEU:HD22	1.97	0.46
1:D:252:HIS:HB3	1:D:256:GLU:HG3	1.98	0.46
1:B:144:LYS:HE3	1:B:343[A]:ARG:O	2.16	0.45
1:B:177[B]:THR:O	1:B:177[B]:THR:CG2	2.64	0.45
1:B:133:LEU:HD12	1:B:145:PHE:CZ	2.52	0.45
1:D:85[B]:MET:HE3	1:D:85[B]:MET:HB3	1.87	0.44
1:D:179[B]:VAL:HG11	1:D:204:GLN:HB2	1.98	0.44
1:D:236:LEU:HA	1:D:242[B]:MET:HG3	2.00	0.44
1:B:179[B]:VAL:HG23	1:B:179[B]:VAL:O	2.19	0.43
1:C:215:VAL:HB	1:C:222:PHE:CE1	2.53	0.43
1:B:49:GLN:O	1:B:53[B]:GLN:HG3	2.18	0.43
1:D:179[B]:VAL:CG1	1:D:204:GLN:HB3	2.48	0.43
1:B:261:PRO:HG2	1:B:266:TRP:HH2	1.84	0.43
1:B:42:LYS:HE3	1:B:96:LEU:HD22	2.00	0.43
1:C:36:ILE:HG12	1:C:92:PHE:HB3	1.99	0.43
1:D:236:LEU:HD23	1:D:242[B]:MET:HG3	2.00	0.42
1:C:232:ARG:NE	1:C:251:GLN:O	2.49	0.42
1:A:211:ALA:HB3	1:A:226:PHE:CZ	2.55	0.41
1:A:160:ALA:HA	1:D:112:ASN:HA	2.01	0.41
1:B:179[B]:VAL:HG11	1:B:204:GLN:HG3	2.02	0.41
1:B:258:GLU:HG2	1:B:261:PRO:HG3	2.03	0.41
1:D:133:LEU:HD12	1:D:145:PHE:CZ	2.55	0.41
1:D:179[B]:VAL:O	1:D:179[B]:VAL:CG2	2.69	0.41
1:B:85[B]:MET:HE3	1:B:85[B]:MET:HB3	1.98	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	358/357 (100%)	346 (97%)	12 (3%)	0	100	100
1	B	358/357 (100%)	349 (98%)	8 (2%)	1 (0%)	36	20
1	C	355/357 (99%)	347 (98%)	7 (2%)	1 (0%)	36	20
1	D	354/357 (99%)	345 (98%)	8 (2%)	1 (0%)	36	20
All	All	1425/1428 (100%)	1387 (97%)	35 (2%)	3 (0%)	43	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	154	LYS
1	D	154	LYS
1	C	154	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/291 (102%)	296 (100%)	1 (0%)	86	79
1	B	298/291 (102%)	294 (99%)	4 (1%)	61	38
1	C	296/291 (102%)	293 (99%)	3 (1%)	68	49
1	D	295/291 (101%)	294 (100%)	1 (0%)	86	79
All	All	1186/1164 (102%)	1177 (99%)	9 (1%)	76	58

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	157	PHE
1	B	157	PHE
1	B	179[A]	VAL
1	B	179[B]	VAL
1	B	299	SER
1	C	44	ASP
1	C	157	PHE
1	C	299	SER
1	D	157	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	53	GLN
1	A	109	GLN
1	A	112	ASN
1	B	109	GLN
1	B	219	HIS
1	B	288	GLN
1	C	53	GLN
1	C	219	HIS
1	D	112	ASN
1	D	172	GLN
1	D	234	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	A	103	1	4,7,8	3.01	1 (25%)	1,8,10	0.79	0
1	CSD	C	103	1	4,7,8	2.74	1 (25%)	1,8,10	0.69	0
1	CSD	B	103	1	4,7,8	2.31	1 (25%)	1,8,10	2.19	1 (100%)
1	CSD	D	103	1	4,7,8	3.06	1 (25%)	1,8,10	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	103	1	-	0/2/6/8	-
1	CSD	C	103	1	-	0/2/6/8	-
1	CSD	B	103	1	-	0/2/6/8	-
1	CSD	D	103	1	-	0/2/6/8	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	103	CSD	OD1-SG	-5.70	1.42	1.47
1	A	103	CSD	OD1-SG	-5.61	1.42	1.47
1	C	103	CSD	OD1-SG	-5.04	1.43	1.47
1	B	103	CSD	OD1-SG	-4.01	1.44	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	103	CSD	OD1-SG-CB	2.19	109.64	105.60

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 14 ligands modelled in this entry, 5 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	D	1357	-	4,4,4	0.85	0	6,6,6	0.42	0
5	ACT	B	1358	-	3,3,3	0.74	0	3,3,3	1.12	0
2	PO4	B	1357	-	4,4,4	0.90	0	6,6,6	0.51	0
5	ACT	D	1358	-	3,3,3	0.88	0	3,3,3	1.04	0
2	PO4	D	1356	-	4,4,4	0.84	0	6,6,6	0.74	0
3	GOL	A	1357	-	5,5,5	0.37	0	5,5,5	0.15	0
5	ACT	C	1357	-	3,3,3	0.75	0	3,3,3	1.33	0
2	PO4	A	1356	-	4,4,4	0.46	0	6,6,6	0.72	0
6	D1D	D	1359	-	8,8,8	0.54	0	6,10,10	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1357	-	-	0/4/4/4	-
6	D1D	D	1359	-	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1358	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	353/357 (98%)	0.44	20 (5%) 29 33	9, 25, 40, 63	8 (2%)
1	B	351/357 (98%)	0.34	29 (8%) 17 20	10, 23, 41, 53	10 (2%)
1	C	350/357 (98%)	0.47	35 (10%) 12 14	9, 24, 49, 66	8 (2%)
1	D	350/357 (98%)	0.45	19 (5%) 31 35	10, 26, 39, 47	8 (2%)
All	All	1404/1428 (98%)	0.43	103 (7%) 21 24	9, 24, 42, 66	34 (2%)

All (103) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	94	TYR	5.8
1	C	43	THR	5.1
1	B	307	GLY	5.0
1	B	98	ALA	4.2
1	C	47	LEU	4.0
1	C	253	GLY	3.9
1	A	111	PRO	3.9
1	C	254	LYS	3.9
1	C	262	LEU	3.9
1	B	112	ASN	3.8
1	C	120	TRP	3.8
1	C	259	LEU	3.7
1	D	98	ALA	3.7
1	B	304	TYR	3.7
1	C	96	LEU	3.6
1	B	41	LYS	3.6
1	A	120	TRP	3.5
1	A	99	ALA	3.5
1	C	255	ASN	3.5
1	C	44	ASP	3.5
1	C	50	VAL	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	304	TYR	3.4
1	C	41	LYS	3.3
1	C	260	ASP	3.3
1	C	307	GLY	3.3
1	B	310	ARG	3.3
1	D	307	GLY	3.2
1	C	310	ARG	3.2
1	C	46	GLU	3.2
1	A	277	ASP	3.2
1	A	56	ARG	3.2
1	B	356	ARG	3.2
1	C	40	THR	3.2
1	D	96	LEU	3.1
1	C	93	ALA	3.1
1	A	43	THR	3.1
1	D	41	LYS	3.1
1	B	260	ASP	3.1
1	B	254	LYS	3.0
1	D	120	TRP	3.0
1	B	120	TRP	3.0
1	C	277	ASP	2.9
1	A	170[A]	HIS	2.9
1	B	306	GLU	2.9
1	C	49	GLN	2.9
1	A	255	ASN	2.9
1	D	112	ASN	2.9
1	B	277	ASP	2.8
1	C	94	TYR	2.8
1	B	111	PRO	2.8
1	B	262	LEU	2.8
1	A	112	ASN	2.8
1	C	42	LYS	2.8
1	A	98	ALA	2.8
1	D	94	TYR	2.8
1	B	309	SER	2.8
1	D	109[A]	GLN	2.6
1	A	238	ALA	2.6
1	C	92	PHE	2.6
1	B	97	LYS	2.6
1	C	356	ARG	2.6
1	A	101	ASP	2.5
1	D	277	ASP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	251	GLN	2.5
1	C	251	GLN	2.5
1	C	222	PHE	2.5
1	B	96	LEU	2.5
1	B	259	LEU	2.5
1	C	257	LYS	2.5
1	C	308	LYS	2.5
1	B	34	ASP	2.5
1	B	255	ASN	2.5
1	D	111	PRO	2.5
1	A	262	LEU	2.4
1	C	45	GLU	2.4
1	D	49	GLN	2.4
1	D	306	GLU	2.3
1	D	355	ALA	2.3
1	B	305	GLU	2.3
1	A	38	GLY	2.3
1	B	308	LYS	2.3
1	B	93	ALA	2.3
1	C	288	GLN	2.3
1	D	251	GLN	2.3
1	C	97	LYS	2.3
1	D	38	GLY	2.3
1	D	33	VAL	2.3
1	B	90	THR	2.2
1	D	346	GLU	2.2
1	A	40	THR	2.2
1	D	43	THR	2.2
1	B	251	GLN	2.2
1	B	257	LYS	2.2
1	A	96	LEU	2.2
1	C	261	PRO	2.2
1	B	261	PRO	2.1
1	C	172	GLN	2.1
1	A	34	ASP	2.1
1	A	100	ARG	2.1
1	B	263	PRO	2.1
1	A	97	LYS	2.1
1	D	15	TYR	2.1
1	C	311	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSD	C	103	8/9	0.96	0.09	21,23,29,30	0
1	CSD	A	103	8/9	0.97	0.07	20,22,25,26	0
1	CSD	D	103	8/9	0.97	0.08	20,21,26,26	0
1	CSD	B	103	8/9	0.98	0.06	19,21,26,30	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	D	1357	5/5	0.67	0.15	56,56,57,58	0
5	ACT	D	1358	4/4	0.75	0.21	37,38,38,38	0
5	ACT	B	1358	4/4	0.76	0.23	34,35,36,36	0
4	CL	A	1358	1/1	0.78	0.24	69,69,69,69	0
5	ACT	C	1357	4/4	0.85	0.16	30,31,31,31	0
4	CL	B	1359	1/1	0.87	0.16	56,56,56,56	0
4	CL	D	1360	1/1	0.88	0.18	52,52,52,52	0
4	CL	C	1358	1/1	0.91	0.16	48,48,48,48	0
6	D1D	D	1359	8/8	0.91	0.17	25,26,28,28	8
2	PO4	A	1356	5/5	0.92	0.14	27,28,33,34	0
3	GOL	A	1357	6/6	0.93	0.11	27,29,30,31	0
2	PO4	B	1357	5/5	0.94	0.14	29,31,32,33	0
4	CL	B	1360	1/1	0.94	0.10	32,32,32,32	1
2	PO4	D	1356	5/5	0.98	0.12	23,25,27,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.