



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:15 PM UTC

PDB ID : 3X1L / pdb_00003x1l
Title : Crystal Structure of the CRISPR-Cas RNA Silencing Cmr Complex Bound to a Target Analog
Authors : Osawa, T.; Numata, T.
Deposited on : 2014-11-20
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)
RNA backbone	3983	1006 (2.42-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	677	 6% 79% 6% 15%
2	B	322	 6% 94% 5%
3	C	357	 6% 84% 6% 10%
3	D	357	 9% 86% 7% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	357	
4	F	155	
4	G	155	
5	H	349	
6	I	39	
7	J	31	

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 19952 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CRISPR system Cmr subunit Cmr2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	573	Total	C	N	O	S	0	0	0
			4470	2920	749	789	12			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	195	MET	-	expression tag	UNP Q8U1S6
A	196	GLY	-	expression tag	UNP Q8U1S6
A	197	SER	-	expression tag	UNP Q8U1S6
A	198	SER	-	expression tag	UNP Q8U1S6
A	199	HIS	-	expression tag	UNP Q8U1S6
A	200	HIS	-	expression tag	UNP Q8U1S6
A	201	HIS	-	expression tag	UNP Q8U1S6
A	202	HIS	-	expression tag	UNP Q8U1S6
A	203	HIS	-	expression tag	UNP Q8U1S6
A	204	HIS	-	expression tag	UNP Q8U1S6
A	205	SER	-	expression tag	UNP Q8U1S6
A	206	SER	-	expression tag	UNP Q8U1S6
A	207	GLY	-	expression tag	UNP Q8U1S6
A	208	LEU	-	expression tag	UNP Q8U1S6
A	209	VAL	-	expression tag	UNP Q8U1S6
A	210	PRO	-	expression tag	UNP Q8U1S6
A	211	ARG	-	expression tag	UNP Q8U1S6
A	212	GLY	-	expression tag	UNP Q8U1S6
A	213	SER	-	expression tag	UNP Q8U1S6
A	214	HIS	-	expression tag	UNP Q8U1S6
A	215	MET	-	expression tag	UNP Q8U1S6

- Molecule 2 is a protein called CRISPR system Cmr subunit Cmr3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	319	Total	C	N	O	S	0	0	0
			2458	1611	394	449	4			

- Molecule 3 is a protein called Cmr4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	322	Total	C	N	O	S	0	0	0
			2405	1549	411	444	1			
3	D	331	Total	C	N	O	S	0	0	0
			2428	1571	413	442	2			
3	E	330	Total	C	N	O	S	0	0	0
			2395	1550	410	434	1			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	MET	-	expression tag	UNP O28416
C	0	GLY	-	expression tag	UNP O28416
D	-1	MET	-	expression tag	UNP O28416
D	0	GLY	-	expression tag	UNP O28416
E	-1	MET	-	expression tag	UNP O28416
E	0	GLY	-	expression tag	UNP O28416

- Molecule 4 is a protein called CRISPR system Cmr subunit Cmr5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	146	Total	C	N	O	S	0	0	0
			1110	722	180	206	2			
4	G	152	Total	C	N	O	S	0	0	0
			1167	753	186	226	2			

- Molecule 5 is a protein called Cmr6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	H	297	Total	C	N	O	S	0	0	0
			2260	1469	363	425	3			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	344	HIS	-	expression tag	UNP O28418
H	345	HIS	-	expression tag	UNP O28418

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	346	HIS	-	expression tag	UNP O28418
H	347	HIS	-	expression tag	UNP O28418
H	348	HIS	-	expression tag	UNP O28418
H	349	HIS	-	expression tag	UNP O28418

- Molecule 6 is a RNA chain called RNA (32-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	I	32	Total	C	N	O	P	0	0	0
			679	304	117	227	31			

- Molecule 7 is a DNA chain called DNA (5'-D(*TP*GP*CP*TP*CP*TP*CP*AP*GP*CP*CP*GP*CP*AP*AP*GP*GP*AP*CP*CP*GP*CP*AP*TP*AP*CP*TP*AP*CP*AP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	J	22	Total	C	N	O	P	0	0	0
			448	212	88	126	22			

- Molecule 8 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Zn	0	0
			1	1		

- Molecule 9 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	1	Total	Mg	0	0
			1	1		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	32	Total	O	0	0
			32	32		
10	B	23	Total	O	0	0
			23	23		
10	C	16	Total	O	0	0
			16	16		
10	D	14	Total	O	0	0
			14	14		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	E	13	Total 13	O 13	0	0
10	G	5	Total 5	O 5	0	0
10	H	4	Total 4	O 4	0	0
10	I	23	Total 23	O 23	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CRISPR system Cmr subunit Cmr2




• Molecule 2: CRISPR system Cmr subunit Cmr3





• Molecule 3: Cmr4









• Molecule 3: Cmr4




• Molecule 3: Cmr4




• Molecule 4: CRISPR system Cmr subunit Cmr5




• Molecule 4: CRISPR system Cmr subunit Cmr5




• Molecule 5: Cmr6






● Molecule 6: RNA (32-MER)



● Molecule 7: DNA (5'-D(*TP*GP*CP*TP*CP*TP*CP*AP*GP*CP*CP*GP*CP*AP*AP*GP*GP*AP*CP*CP*GP*CP*AP*TP*AP*CP*TP*AP*CP*AP*A)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	75.49Å 76.22Å 139.19Å 90.32° 104.83° 118.58°	Depositor
Resolution (Å)	44.34 – 2.10 44.34 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.4 (44.34-2.10) 97.4 (44.34-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 2.10Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.208 , 0.246 0.210 , 0.247	Depositor DCC
R_{free} test set	7443 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	32.8	Xtriage
Anisotropy	0.397	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 34.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.005 for k,-h-k,h+l 0.005 for -h-k,h,h+k+l 0.017 for h,-h-k,-h-l 0.017 for -h-k,k,-l 0.016 for k,h,-h-k-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	19952	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/4561	0.66	0/6184
2	B	0.29	0/2512	0.66	0/3401
3	C	0.30	0/2445	0.68	0/3320
3	D	0.29	0/2467	0.67	0/3348
3	E	0.29	0/2436	0.70	0/3315
4	F	0.30	0/1131	0.67	0/1527
4	G	0.29	0/1190	0.66	0/1609
5	H	0.29	0/2306	0.64	0/3124
6	I	0.15	0/758	0.40	0/1180
7	J	0.20	0/503	0.36	0/772
All	All	0.29	0/20309	0.65	0/27780

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4470	0	4434	23	0
2	B	2458	0	2480	9	0
3	C	2405	0	2362	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	2428	0	2351	14	0
3	E	2395	0	2289	16	0
4	F	1110	0	1097	9	0
4	G	1167	0	1137	11	0
5	H	2260	0	2132	10	0
6	I	679	0	343	5	0
7	J	448	0	245	3	0
8	A	1	0	0	0	0
9	B	1	0	0	0	0
10	A	32	0	0	0	0
10	B	23	0	0	0	0
10	C	16	0	0	0	0
10	D	14	0	0	0	0
10	E	13	0	0	0	0
10	G	5	0	0	0	0
10	H	4	0	0	0	0
10	I	23	0	0	0	0
All	All	19952	0	18870	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 95 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:331:GLY:O	6:I:20:G:N2	2.27	0.68
3:C:201:ILE:HD13	4:F:131:ILE:HG12	1.77	0.66
5:H:29:ILE:HD13	5:H:35:ILE:HB	1.80	0.64
2:B:42:ILE:HD13	2:B:178:ILE:HD11	1.82	0.62
1:A:422:LYS:HB3	2:B:108:PRO:HG3	1.83	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	559/677 (83%)	551 (99%)	8 (1%)	0	100	100
2	B	315/322 (98%)	304 (96%)	11 (4%)	0	100	100
3	C	316/357 (88%)	309 (98%)	7 (2%)	0	100	100
3	D	319/357 (89%)	310 (97%)	9 (3%)	0	100	100
3	E	322/357 (90%)	313 (97%)	9 (3%)	0	100	100
4	F	142/155 (92%)	141 (99%)	1 (1%)	0	100	100
4	G	150/155 (97%)	143 (95%)	7 (5%)	0	100	100
5	H	291/349 (83%)	283 (97%)	7 (2%)	1 (0%)	36	36
All	All	2414/2729 (88%)	2354 (98%)	59 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	H	217	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	457/604 (76%)	455 (100%)	2 (0%)	84	89
2	B	257/280 (92%)	255 (99%)	2 (1%)	73	81
3	C	243/309 (79%)	241 (99%)	2 (1%)	73	81
3	D	238/309 (77%)	238 (100%)	0	100	100
3	E	228/309 (74%)	225 (99%)	3 (1%)	61	69
4	F	111/131 (85%)	111 (100%)	0	100	100
4	G	119/131 (91%)	119 (100%)	0	100	100
5	H	224/309 (72%)	222 (99%)	2 (1%)	70	78
All	All	1877/2382 (79%)	1866 (99%)	11 (1%)	78	86

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	E	268	ASP
3	E	338	LYS
5	H	89	TYR
5	H	8	LYS
3	C	39	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	239	GLN
1	A	297	ASN
3	E	218	ASN
4	F	113	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
6	I	31/39 (79%)	3 (9%)	0

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
6	I	2	U
6	I	14	G
6	I	27	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	573/677 (84%)	0.54	38 (6%) 24 26	21, 35, 58, 65	0
2	B	319/322 (99%)	0.42	20 (6%) 26 27	21, 33, 51, 64	0
3	C	322/357 (90%)	0.44	20 (6%) 26 28	21, 34, 52, 65	0
3	D	331/357 (92%)	0.72	32 (9%) 13 14	23, 37, 64, 81	0
3	E	330/357 (92%)	0.88	55 (16%) 4 4	25, 39, 64, 77	0
4	F	146/155 (94%)	0.72	15 (10%) 12 12	24, 37, 60, 72	0
4	G	152/155 (98%)	0.44	9 (5%) 28 30	24, 35, 52, 61	0
5	H	297/349 (85%)	0.77	31 (10%) 11 12	25, 39, 68, 77	0
6	I	32/39 (82%)	-0.59	0 100 100	25, 29, 61, 79	0
7	J	22/31 (70%)	0.33	0 100 100	32, 46, 60, 62	0
All	All	2524/2799 (90%)	0.60	220 (8%) 16 16	21, 36, 61, 81	0

The worst 5 of 220 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	196	GLY	5.6
2	B	50	GLY	5.3
1	A	527	VAL	5.3
4	F	103	GLY	5.1
3	E	80	THR	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	MG	B	401	1/1	0.98	0.04	24,24,24,24	0
8	ZN	A	901	1/1	1.00	0.04	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.