



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 09:52 PM UTC

PDB ID : 5X1J / pdb_00005x1j
Title : Vanillate/3-O-methylgallate O-demethylase, LigM, vanillate complex form
Authors : Harada, A.; Senda, T.
Deposited on : 2017-01-26
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	474	0% 95% 5%
1	B	474	0% 92% 5%
1	C	474	3% 90% 6%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11482 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Vanillate/3-O-methylgallate O-demethylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	455	Total	C	N	O	S	0	2	0
			3568	2285	605	665	13			
1	B	451	Total	C	N	O	S	0	4	0
			3550	2276	598	663	13			
1	C	455	Total	C	N	O	S	0	2	0
			3550	2280	599	658	13			

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP G2IQS7
A	-1	SER	-	expression tag	UNP G2IQS7
A	0	SER	-	expression tag	UNP G2IQS7
B	-2	GLY	-	expression tag	UNP G2IQS7
B	-1	SER	-	expression tag	UNP G2IQS7
B	0	SER	-	expression tag	UNP G2IQS7
C	-2	GLY	-	expression tag	UNP G2IQS7
C	-1	SER	-	expression tag	UNP G2IQS7
C	0	SER	-	expression tag	UNP G2IQS7

- Molecule 2 is 4-HYDROXY-3-METHOXYBENZOATE (CCD ID: VNL) (formula: C₈H₇O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	8	4		
2	B	1	Total	C	O	0	0
			12	8	4		
2	C	1	Total	C	O	0	0
			12	8	4		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	4	1	3		
4	B	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	319	Total	O	0	0
			319	319		
5	B	254	Total	O	0	0
			254	254		
5	C	169	Total	O	0	0
			169	169		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Vanillate/3-O-methylgallate O-demethylase



- Molecule 1: Vanillate/3-O-methylgallate O-demethylase



- Molecule 1: Vanillate/3-O-methylgallate O-demethylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	102.82Å 118.04Å 130.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.82 – 1.90 47.82 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.9 (47.82-1.90) 93.6 (47.82-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 1.90Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.172 , 0.204 0.173 , 0.205	Depositor DCC
R_{free} test set	6147 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtriage
Anisotropy	0.261	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11482	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VNL, EDO, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/3673	0.54	0/5007
1	B	0.29	0/3652	0.53	0/4979
1	C	0.25	0/3652	0.47	0/4981
All	All	0.29	0/10977	0.52	0/14967

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3568	0	3432	2	0
1	B	3550	0	3401	9	0
1	C	3550	0	3397	16	0
2	A	12	0	7	0	0
2	B	12	0	7	0	0
2	C	12	0	7	0	0
3	A	8	0	12	0	0
3	B	8	0	12	0	0
3	C	4	0	6	0	0
4	A	8	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	8	0	12	2	0
5	A	319	0	0	0	0
5	B	254	0	0	2	0
5	C	169	0	0	0	0
All	All	11482	0	10305	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:430:THR:O	4:B:504:TRS:H32	1.87	0.75
1:B:435:VAL:O	4:B:504:TRS:H31	1.87	0.74
1:B:452:TYR:O	5:B:601:HOH:O	2.11	0.66
1:C:64:LEU:HD13	1:C:126:ALA:HB1	1.80	0.63
1:C:104:ILE:HG12	1:C:264[A]:ILE:HD11	1.87	0.57
1:B:64:LEU:HB2	1:B:119:TYR:HB3	1.88	0.54
1:C:345:ASP:HB3	1:C:441:MET:HE1	1.88	0.54
1:B:37:GLU:OE2	5:B:602:HOH:O	2.18	0.53
1:C:64:LEU:CD1	1:C:126:ALA:HB1	2.41	0.51
1:C:261:LEU:HD12	1:C:262:PRO:HD2	1.93	0.50
1:A:64:LEU:HB2	1:A:119:TYR:HB3	1.96	0.48
1:C:119:TYR:CE1	1:C:126:ALA:HB2	2.50	0.47
1:B:251:THR:HA	1:B:254:SER:OG	2.16	0.46
1:B:112:LEU:HD23	1:B:191:MET:HE1	2.00	0.43
1:C:110:PHE:HB2	1:C:118:VAL:CG1	2.48	0.43
1:C:363:LYS:HB2	1:C:438:HIS:CD2	2.53	0.43
1:C:378:ASP:HB2	1:C:391:SER:HB3	2.00	0.43
1:C:172:TRP:CZ2	1:C:185:LYS:HG2	2.53	0.43
1:C:441:MET:HE2	1:C:443:VAL:HG22	2.01	0.43
1:C:45:GLN:HG2	1:C:48:TRP:CH2	2.54	0.43
1:C:305:TYR:CD2	1:C:330:PRO:HB3	2.54	0.42
1:C:3:ALA:HA	1:C:4:PRO:HD3	1.89	0.42
1:B:45:GLN:HG2	1:B:48:TRP:CH2	2.55	0.41
1:A:303:ASN:HB2	1:A:304:PRO:HD2	2.02	0.41
1:C:86:TRP:CD2	1:C:92:LYS:HD2	2.55	0.41
1:B:264:ILE:HA	1:B:270:LEU:HD13	2.03	0.41
1:C:420:THR:HA	1:C:443:VAL:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	455/474 (96%)	445 (98%)	10 (2%)	0	100	100
1	B	453/474 (96%)	444 (98%)	9 (2%)	0	100	100
1	C	455/474 (96%)	444 (98%)	11 (2%)	0	100	100
All	All	1363/1422 (96%)	1333 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/382 (96%)	365 (100%)	0	100	100
1	B	362/382 (95%)	362 (100%)	0	100	100
1	C	357/382 (94%)	357 (100%)	0	100	100
All	All	1084/1146 (95%)	1084 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	223	GLN
1	A	386	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	158	GLN
1	B	411	HIS
1	C	250	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	VNL	C	501	-	12,12,12	1.26	2 (16%)	16,16,16	1.42	2 (12%)
3	EDO	A	502	-	3,3,3	0.44	0	2,2,2	0.24	0
2	VNL	A	501	-	12,12,12	1.11	1 (8%)	16,16,16	1.85	3 (18%)
3	EDO	A	503	-	3,3,3	0.51	0	2,2,2	0.56	0
2	VNL	B	501	-	12,12,12	1.14	1 (8%)	16,16,16	1.49	3 (18%)
3	EDO	C	502	-	3,3,3	0.50	0	2,2,2	0.18	0
3	EDO	B	503	-	3,3,3	0.47	0	2,2,2	0.19	0
4	TRS	A	504	-	7,7,7	0.27	0	9,9,9	0.93	0
3	EDO	B	502	-	3,3,3	0.45	0	2,2,2	0.51	0
4	TRS	B	504	-	7,7,7	0.32	0	9,9,9	1.13	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	VNL	C	501	-	-	2/6/6/6	0/1/1/1
3	EDO	A	502	-	-	0/1/1/1	-
2	VNL	A	501	-	-	2/6/6/6	0/1/1/1
3	EDO	A	503	-	-	0/1/1/1	-
2	VNL	B	501	-	-	2/6/6/6	0/1/1/1
3	EDO	C	502	-	-	0/1/1/1	-
3	EDO	B	503	-	-	0/1/1/1	-
4	TRS	A	504	-	-	7/9/9/9	-
3	EDO	B	502	-	-	0/1/1/1	-
4	TRS	B	504	-	-	3/9/9/9	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	VNL	OM-CM2	2.48	1.41	1.37
2	C	501	VNL	OM-CM2	2.43	1.41	1.37
2	C	501	VNL	O3-CZ	2.17	1.40	1.36
2	B	501	VNL	O3-CZ	2.03	1.40	1.36

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	VNL	OM-CM2-CZ	4.83	121.79	114.55
2	C	501	VNL	OM-CM2-CZ	3.92	120.43	114.55
2	B	501	VNL	OM-CM2-CZ	3.69	120.08	114.55
2	A	501	VNL	OM-CM2-CO2	-3.08	118.78	124.08
4	B	504	TRS	O3-C3-C	2.57	118.03	110.88
2	B	501	VNL	OM-CM2-CO2	-2.33	120.06	124.08
2	C	501	VNL	OM-CM2-CO2	-2.32	120.09	124.08
2	A	501	VNL	O1-CC-C1	2.19	120.45	114.84
2	B	501	VNL	CV-OM-CM2	-2.06	114.50	117.51

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	504	TRS	C2-C-C1-O1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	504	TRS	C3-C-C1-O1
4	B	504	TRS	N-C-C1-O1
2	C	501	VNL	CO2-CM2-OM-CV
2	A	501	VNL	CO2-CM2-OM-CV
2	B	501	VNL	CO2-CM2-OM-CV
4	A	504	TRS	C1-C-C3-O3
2	C	501	VNL	CZ-CM2-OM-CV
2	A	501	VNL	CZ-CM2-OM-CV
2	B	501	VNL	CZ-CM2-OM-CV
4	A	504	TRS	C1-C-C2-O2
4	A	504	TRS	C3-C-C2-O2
4	A	504	TRS	N-C-C3-O3
4	A	504	TRS	C2-C-C3-O3
4	A	504	TRS	N-C-C2-O2
4	A	504	TRS	N-C-C1-O1

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	504	TRS	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	455/474 (95%)	-0.37	4 (0%)	81 83	8, 18, 30, 52	2 (0%)
1	B	451/474 (95%)	-0.20	4 (0%)	81 83	8, 21, 35, 58	4 (0%)
1	C	455/474 (95%)	0.32	16 (3%)	47 50	14, 28, 44, 54	2 (0%)
All	All	1361/1422 (95%)	-0.08	24 (1%)	67 71	8, 22, 39, 58	8 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	ALA	4.6
1	B	208	MET	4.4
1	C	456	ALA	4.0
1	B	209	ALA	3.5
1	C	415	VAL	3.1
1	B	3	ALA	3.0
1	C	109[A]	ILE	3.0
1	A	456	ALA	2.9
1	A	455	THR	2.7
1	C	292	PHE	2.6
1	C	318	ASP	2.5
1	C	64	LEU	2.5
1	C	209	ALA	2.4
1	C	3	ALA	2.3
1	C	210	GLY	2.3
1	C	137	GLY	2.3
1	C	383	ALA	2.2
1	C	315	PHE	2.2
1	C	264[A]	ILE	2.2
1	A	457	ARG	2.2
1	C	135	THR	2.1
1	B	13	ALA	2.1
1	C	455	THR	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	387	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	TRS	B	504	8/8	0.85	0.14	26,32,38,40	0
4	TRS	A	504	8/8	0.90	0.11	24,35,42,44	0
2	VNL	C	501	12/12	0.95	0.07	22,27,30,31	0
3	EDO	B	503	4/4	0.95	0.08	19,20,22,22	0
2	VNL	A	501	12/12	0.96	0.06	14,17,22,23	0
2	VNL	B	501	12/12	0.97	0.06	16,20,23,24	0
3	EDO	C	502	4/4	0.97	0.06	25,26,26,29	0
3	EDO	A	502	4/4	0.97	0.06	16,17,20,23	0
3	EDO	A	503	4/4	0.97	0.07	17,18,21,23	0
3	EDO	B	502	4/4	0.98	0.04	20,20,21,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.