



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2026 – 01:48 AM UTC

PDB ID : 5X2G / pdb_00005x2g
Title : Crystal structure of Campylobacter jejuni Cas9 in complex with sgRNA and target DNA (AGAAACC PAM)
Authors : Yamada, M.; Watanabe, Y.; Hirano, H.; Nakane, T.; Ishitani, R.; Nishimasu, H.; Nureki, O.
Deposited on : 2017-01-31
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)
RNA backbone	3983	1155 (2.70-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	835	8% 81% 8% 10%
2	B	93	85% 14%
3	C	28	93% 7%
4	D	8	88% 12%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8967 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CRISPR-associated endonuclease Cas9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	749	Total	C	N	O	S	0	0	0
			5929	3824	1009	1085	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	SER	-	expression tag	UNP Q0P897
A	-4	SER	-	expression tag	UNP Q0P897
A	-3	GLY	-	expression tag	UNP Q0P897
A	-2	SER	-	expression tag	UNP Q0P897
A	-1	GLY	-	expression tag	UNP Q0P897
A	0	HIS	-	expression tag	UNP Q0P897
A	481	GLY	-	linker	UNP Q0P897
A	637	GLY	-	linker	UNP Q0P897
A	638	GLY	-	linker	UNP Q0P897
A	639	SER	-	linker	UNP Q0P897
A	640	GLY	-	linker	UNP Q0P897
A	641	GLY	-	linker	UNP Q0P897

- Molecule 2 is a RNA chain called sgRNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	93	Total	C	N	O	P	0	0	0
			1990	889	361	647	93			

- Molecule 3 is a DNA chain called Target DNA strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	28	Total	C	N	O	P	0	0	0
			562	270	96	169	27			

- Molecule 4 is a DNA chain called Non-target DNA strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	8	Total 163	C 78	N 36	O 42	P 7	0	0	0

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total 4	C 2	O 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	166	Total 166	O 166	0	0
6	B	108	Total 108	O 108	0	0
6	C	30	Total 30	O 30	0	0
6	D	15	Total 15	O 15	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CRISPR-associated endonuclease Cas9




• Molecule 2: sgRNA



• Molecule 3: Target DNA strand



• Molecule 4: Non-target DNA strand





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	104.34Å 105.09Å 136.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.55 – 2.40 52.55 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (52.55-2.40) 99.9 (52.55-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 2.40Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.190 , 0.221 0.194 , 0.225	Depositor DCC
R_{free} test set	2833 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	39.9	Xtriage
Anisotropy	0.722	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.012 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8967	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.10	0/6034	0.29	0/8114
2	B	0.11	0/2227	0.28	0/3470
3	C	0.22	0/627	0.40	0/964
4	D	0.24	0/184	0.49	0/282
All	All	0.12	0/9072	0.30	0/12830

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5929	0	5866	35	0
2	B	1990	0	1002	3	0
3	C	562	0	318	3	0
4	D	163	0	90	1	0
5	B	4	0	6	0	0
6	A	166	0	0	0	0
6	B	108	0	0	0	0
6	C	30	0	0	0	0
6	D	15	0	0	0	0
All	All	8967	0	7282	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:866:ARG:NH1	3:C:-6:DG:N7	2.31	0.79
1:A:99:ILE:O	1:A:120:ARG:NH2	2.17	0.77
1:A:3:ARG:NH2	1:A:470:TYR:O	2.32	0.62
1:A:134:ILE:HD11	3:C:6:DG:H4'	1.85	0.57
1:A:108:ALA:HB2	1:A:113:LEU:HD11	1.87	0.56
1:A:318:ASP:HA	1:A:321:PHE:HE2	1.69	0.56
1:A:32:PHE:CG	1:A:458:GLU:HG3	2.43	0.53
1:A:710:ASP:O	1:A:714:ILE:HG12	2.09	0.53
1:A:700:ASP:O	1:A:706:HIS:HB3	2.08	0.52
1:A:288:LYS:O	1:A:292:ASN:ND2	2.40	0.52
1:A:893:LEU:HD22	1:A:965:VAL:HG22	1.93	0.50
1:A:20:GLU:OE1	1:A:25:LYS:NZ	2.45	0.50
1:A:135:LYS:HG3	3:C:6:DG:H5''	1.94	0.49
1:A:439:ASN:HA	1:A:444:LYS:HD3	1.95	0.48
1:A:472:LYS:NZ	1:A:656:ASP:OD2	2.40	0.48
1:A:712:VAL:HG11	1:A:766:VAL:HG21	1.95	0.48
1:A:134:ILE:HD13	1:A:134:ILE:H	1.79	0.48
1:A:867:SER:HB3	1:A:873:LYS:H	1.78	0.47
1:A:212:ARG:NH2	1:A:223:GLU:OE2	2.48	0.47
1:A:246:CYS:HB2	1:A:397:SER:HB3	1.95	0.47
1:A:413:LYS:HB2	1:A:416:GLU:HG3	1.97	0.46
1:A:833:VAL:HG13	1:A:888:LEU:HB2	1.98	0.46
1:A:294:LEU:HD22	1:A:309:THR:HG22	1.98	0.46
1:A:306:TYR:O	1:A:310:LYS:HG3	2.17	0.45
1:A:687:THR:HG22	1:A:714:ILE:HD11	1.99	0.44
1:A:868:LYS:HE2	1:A:868:LYS:HB3	1.88	0.44
1:A:14:ILE:HG12	1:A:32:PHE:HE1	1.81	0.43
1:A:56:LYS:NZ	2:B:67:U:OP2	2.51	0.43
1:A:44:LEU:HD12	1:A:450:PRO:HB2	1.99	0.43
1:A:780:ARG:HH22	4:D:2:DG:P	2.42	0.43
1:A:353:ASN:HB3	1:A:398:PHE:CZ	2.53	0.43
2:B:75:U:O2'	2:B:76:A:H5'	2.19	0.43
1:A:401:LEU:O	1:A:405:THR:OG1	2.25	0.42
1:A:683:SER:O	1:A:687:THR:HG23	2.20	0.42
1:A:298:VAL:HG22	1:A:304:LEU:HB2	2.02	0.41
1:A:450:PRO:HB3	2:B:71:G:H4'	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	735/835 (88%)	715 (97%)	19 (3%)	1 (0%)	48 64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	608/740 (82%)	596 (98%)	12 (2%)	48 70

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ILE
1	A	22	ASP
1	A	27	CYS
1	A	72	ILE
1	A	81	GLU
1	A	134	ILE
1	A	287	THR
1	A	296	ASN
1	A	303	THR
1	A	392	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	690	LEU
1	A	933	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	197	GLN
1	A	381	GLN
1	A	395	ASN
1	A	704	HIS
1	A	706	HIS
1	A	910	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	92/93 (98%)	10 (10%)	1 (1%)

All (10) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	38	G
2	B	49	A
2	B	51	A
2	B	58	G
2	B	64	C
2	B	76	A
2	B	77	C
2	B	79	A
2	B	86	A
2	B	88	A

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	B	63	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	B	101	-	3,3,3	0.43	0	2,2,2	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	101	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	749/835 (89%)	0.49	70 (9%) 14 11	22, 57, 101, 126	0
2	B	93/93 (100%)	-0.62	0 100 100	29, 47, 73, 119	0
3	C	28/28 (100%)	-0.56	0 100 100	34, 42, 72, 75	0
4	D	8/8 (100%)	-0.75	0 100 100	28, 33, 48, 54	0
All	All	878/964 (91%)	0.33	70 (7%) 18 15	22, 55, 99, 126	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	640	GLY	6.3
1	A	136	ASN	6.3
1	A	86	PHE	5.4
1	A	481	GLY	5.1
1	A	315	LEU	5.0
1	A	42	GLU	4.8
1	A	2	ALA	4.7
1	A	316	SER	4.7
1	A	140	LYS	4.6
1	A	88	GLU	4.5
1	A	309	THR	4.5
1	A	41	GLY	4.3
1	A	662	ASP	4.2
1	A	983	LYS	4.2
1	A	716	TYR	4.0
1	A	763	ARG	3.7
1	A	320	GLU	3.7
1	A	1	MET	3.7
1	A	677	VAL	3.7
1	A	27	CYS	3.6
1	A	323	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	317	ASP	3.4
1	A	373	ALA	3.3
1	A	702	ASN	3.3
1	A	318	ASP	3.2
1	A	319	TYR	3.2
1	A	321	PHE	3.1
1	A	661	SER	3.1
1	A	328	TYR	3.0
1	A	324	GLU	3.0
1	A	313	LEU	3.0
1	A	765	LYS	3.0
1	A	698	ALA	2.9
1	A	480	LEU	2.9
1	A	11	ILE	2.9
1	A	376	ASP	2.9
1	A	305	THR	2.8
1	A	141	GLU	2.8
1	A	380	ASN	2.8
1	A	314	GLY	2.8
1	A	289	ASP	2.7
1	A	326	GLY	2.7
1	A	134	ILE	2.7
1	A	764	GLN	2.7
1	A	699	LYS	2.7
1	A	883	GLU	2.6
1	A	942	ALA	2.6
1	A	686	LEU	2.6
1	A	946	GLU	2.5
1	A	694	TRP	2.4
1	A	642	TYR	2.4
1	A	34	LYS	2.4
1	A	660	LEU	2.3
1	A	426	ALA	2.2
1	A	928	PHE	2.2
1	A	641	GLY	2.2
1	A	22	ASP	2.2
1	A	984	LYS	2.2
1	A	14	ILE	2.2
1	A	330	ILE	2.2
1	A	310	LYS	2.2
1	A	284	ILE	2.2
1	A	325	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	301	ASN	2.1
1	A	285	LEU	2.1
1	A	291	LEU	2.1
1	A	135	LYS	2.1
1	A	366	ILE	2.0
1	A	241	HIS	2.0
1	A	322	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EDO	B	101	4/4	0.98	0.07	19,21,23,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.