



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:08 AM UTC

PDB ID : 5X2W / pdb_00005x2w
Title : Crystal structure of Pseudomonas putida methionine gamma-lyase wild type with L-methionine intermediates
Authors : Shiba, T.; Sato, D.; Harada, S.
Deposited on : 2017-02-02
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	8% 91% 7% ..
1	B	398	5% 91% 7% ..
1	C	398	6% 94% ..
1	D	398	9% 92% 7% .

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 12009 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-methionine gamma-lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	0	0
			2951	1863	522	549	17			
1	B	392	Total	C	N	O	S	0	0	0
			2951	1863	522	549	17			
1	C	392	Total	C	N	O	S	0	0	0
			2951	1863	522	549	17			
1	D	398	Total	C	N	O	S	0	0	0
			2996	1889	532	557	18			

- Molecule 2 is (2E)-2-[(3-hydroxy-2-methyl-5-[(phosphonooxy)methyl]pyridin-4-yl)methyl]amino]-4-(methylsulfanyl)but-2-enoic acid (CCD ID: 3LM) (formula: C₁₃H₁₉N₂O₇PS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P S	0	0
			24	13	2	7	1 1		
2	B	1	Total	C	N	O	P S	0	0
			24	13	2	7	1 1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	S	0	0
			24	13	2	7	1	1		
2	D	1	Total	C	N	O	P	S	0	0
			24	13	2	7	1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	11	Total	O	0	0
			11	11		
3	B	22	Total	O	0	0
			22	22		
3	C	12	Total	O	0	0
			12	12		
3	D	19	Total	O	0	0
			19	19		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: L-methionine gamma-lyase



- Molecule 1: L-methionine gamma-lyase




- Molecule 1: L-methionine gamma-lyase



- Molecule 1: L-methionine gamma-lyase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	154.63Å 152.31Å 80.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70 20.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.5 (20.00-2.70) 99.5 (20.00-2.70)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.99 (at 2.71Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.195 , 0.246 0.198 , 0.243	Depositor DCC
R_{free} test set	2681 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtriage
Anisotropy	0.140	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 26.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.006 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12009	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 3LM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/3015	0.85	0/4093
1	B	0.62	0/3015	0.87	0/4093
1	C	0.62	0/3015	0.87	0/4093
1	D	0.64	0/3061	0.88	1/4153 (0.0%)
All	All	0.62	0/12106	0.87	1/16432 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	4	SER	N-CA-C	5.26	117.76	111.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2951	0	2921	9	0
1	B	2951	0	2921	7	0
1	C	2951	0	2921	3	0
1	D	2996	0	2967	13	0
2	A	24	0	15	0	0
2	B	24	0	16	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	24	0	15	0	0
2	D	24	0	15	2	0
3	A	11	0	0	0	0
3	B	22	0	0	0	0
3	C	12	0	0	0	0
3	D	19	0	0	0	0
All	All	12009	0	11791	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:99:THR:HG21	1:D:234:ILE:HA	1.73	0.70
1:A:99:THR:HG21	1:A:234:ILE:HA	1.78	0.66
1:B:70:LEU:HD21	1:B:255:LEU:HD23	1.78	0.63
1:B:99:THR:HG21	1:B:234:ILE:HA	1.82	0.62
1:A:95:SER:O	1:A:99:THR:HG23	2.04	0.58
1:D:300:TYR:CZ	1:D:304:ARG:HD2	2.43	0.54
1:D:368:GLY:O	1:D:369:ILE:HD12	2.08	0.54
1:D:137:MET:HE1	1:D:154:ILE:HG23	1.95	0.49
1:B:95:SER:O	1:B:99:THR:HG23	2.14	0.48
1:B:300:TYR:CZ	1:B:304:ARG:HD2	2.48	0.47
1:D:61:ARG:HG2	1:D:246:VAL:HG21	1.96	0.47
1:D:204:LEU:HD23	1:D:226:GLY:HA3	1.95	0.47
1:B:161:ASN:HD21	1:B:375:ARG:HH11	1.63	0.47
1:C:302:LEU:HA	1:C:305:GLN:HE21	1.80	0.47
1:A:329:MET:HE1	1:A:348:ALA:CB	2.44	0.46
1:A:154:ILE:HD12	1:A:176:ALA:HB2	1.98	0.45
1:D:211:LYS:HE2	2:D:401:3LM:H4A	1.99	0.45
1:D:211:LYS:CE	2:D:401:3LM:H4A	2.48	0.43
1:C:70:LEU:HD21	1:C:255:LEU:HD23	1.99	0.43
1:A:212:TYR:CE1	1:A:342:GLY:HA2	2.54	0.43
1:D:325:GLY:HA3	1:D:350:HIS:NE2	2.35	0.42
1:A:159:PRO:HG3	1:A:166:MET:HE1	2.02	0.42
1:A:339:VAL:O	1:A:340:SER:CB	2.68	0.41
1:D:95:SER:O	1:D:99:THR:HG23	2.20	0.41
1:B:339:VAL:O	1:B:340:SER:CB	2.69	0.41
1:D:152:ARG:NH2	1:D:180:GLY:O	2.53	0.41
2:B:401:3LM:O3	2:B:401:3LM:N	2.51	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:370:SER:OG	1:A:371:GLU:N	2.54	0.40
1:D:161:ASN:HA	1:D:162:PRO:HA	1.91	0.40
1:A:260:LYS:NZ	1:C:345:GLU:OE2	2.54	0.40
1:B:166:MET:HE2	1:B:303:ALA:HB2	2.03	0.40
1:D:78:GLU:OE1	1:D:207:HIS:CE1	2.74	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/398 (98%)	372 (95%)	17 (4%)	1 (0%)	36	60
1	B	390/398 (98%)	370 (95%)	18 (5%)	2 (0%)	24	48
1	C	390/398 (98%)	376 (96%)	13 (3%)	1 (0%)	36	60
1	D	396/398 (100%)	381 (96%)	13 (3%)	2 (0%)	24	48
All	All	1566/1592 (98%)	1499 (96%)	61 (4%)	6 (0%)	30	54

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	368	GLY
1	A	340	SER
1	C	211	LYS
1	B	340	SER
1	D	287	GLU
1	D	368	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/307 (98%)	289 (96%)	13 (4%)	26	54
1	B	302/307 (98%)	282 (93%)	20 (7%)	15	36
1	C	302/307 (98%)	292 (97%)	10 (3%)	33	63
1	D	307/307 (100%)	298 (97%)	9 (3%)	37	67
All	All	1213/1228 (99%)	1161 (96%)	52 (4%)	26	54

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	99	THR
1	A	131	LYS
1	A	133	ARG
1	A	140	LEU
1	A	161	ASN
1	A	200	LEU
1	A	230	LEU
1	A	263	ASN
1	A	280	LEU
1	A	288	LEU
1	A	341	LEU
1	A	374	VAL
1	B	44	GLU
1	B	63	SER
1	B	99	THR
1	B	107	VAL
1	B	111	ASN
1	B	141	GLN
1	B	152	ARG
1	B	161	ASN
1	B	166	MET
1	B	219	ILE
1	B	236	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	276	LEU
1	B	294	LEU
1	B	341	LEU
1	B	355	THR
1	B	360	THR
1	B	363	GLU
1	B	382	ASP
1	B	387	LEU
1	B	392	GLN
1	C	7	LEU
1	C	108	LEU
1	C	131	LYS
1	C	132	LEU
1	C	140	LEU
1	C	141	GLN
1	C	143	LEU
1	C	227	SER
1	C	285	GLN
1	C	349	GLN
1	D	22	GLN
1	D	99	THR
1	D	240	LYS
1	D	285	GLN
1	D	319	LYS
1	D	330	ASN
1	D	369	ILE
1	D	370	SER
1	D	394	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (33) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	57	HIS
1	A	68	ASN
1	A	111	ASN
1	A	122	HIS
1	A	134	HIS
1	A	161	ASN
1	A	263	ASN
1	A	283	GLN
1	A	290	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	68	ASN
1	B	111	ASN
1	B	161	ASN
1	B	274	GLN
1	B	290	HIS
1	B	299	GLN
1	B	392	GLN
1	C	17	HIS
1	C	54	GLN
1	C	161	ASN
1	C	165	HIS
1	C	187	ASN
1	C	305	GLN
1	C	306	GLN
1	D	24	HIS
1	D	34	GLN
1	D	68	ASN
1	D	187	ASN
1	D	195	GLN
1	D	207	HIS
1	D	237	GLN
1	D	283	GLN
1	D	330	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	3LM	C	401	-	22,24,24	1.47	4 (18%)	29,33,33	1.90	7 (24%)
2	3LM	A	401	-	22,24,24	1.40	1 (4%)	29,33,33	1.80	6 (20%)
2	3LM	D	401	-	22,24,24	1.54	5 (22%)	29,33,33	1.93	8 (27%)
2	3LM	B	401	-	22,24,24	1.41	1 (4%)	29,33,33	1.72	8 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3LM	C	401	-	-	5/18/19/19	0/1/1/1
2	3LM	A	401	-	-	5/18/19/19	0/1/1/1
2	3LM	D	401	-	-	5/18/19/19	0/1/1/1
2	3LM	B	401	-	-	5/18/19/19	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	3LM	C6-N1	3.15	1.40	1.34
2	A	401	3LM	C6-N1	3.04	1.40	1.34
2	D	401	3LM	C6-N1	2.95	1.40	1.34
2	B	401	3LM	C6-N1	2.73	1.40	1.34
2	D	401	3LM	C4A-N	2.21	1.50	1.46
2	D	401	3LM	CA-N	2.18	1.38	1.34
2	D	401	3LM	P-OP3	-2.14	1.46	1.54
2	C	401	3LM	C4A-N	2.04	1.50	1.46
2	D	401	3LM	C2A-C2	2.03	1.53	1.50
2	C	401	3LM	O2-C	-2.03	1.25	1.30
2	C	401	3LM	CA-N	2.01	1.37	1.34

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	401	3LM	C4A-N-CA	4.53	131.43	123.44
2	C	401	3LM	O2-C-CA	4.41	123.78	114.27
2	A	401	3LM	O2-C-CA	4.39	123.74	114.27
2	D	401	3LM	OP4-C5A-C5	-4.15	101.57	109.36
2	B	401	3LM	OP4-C5A-C5	-4.15	101.57	109.36
2	C	401	3LM	OP4-C5A-C5	-4.14	101.60	109.36
2	A	401	3LM	C4A-C4-C3	4.07	125.40	119.98
2	B	401	3LM	C4A-N-CA	3.75	130.06	123.44
2	C	401	3LM	C4A-N-CA	3.52	129.66	123.44
2	D	401	3LM	C4A-C4-C3	3.52	124.66	119.98
2	C	401	3LM	C4A-C4-C3	3.50	124.64	119.98
2	D	401	3LM	O2-C-O1	-3.33	115.98	123.90
2	C	401	3LM	O2-C-O1	-3.11	116.50	123.90
2	B	401	3LM	O2-C-O1	-3.07	116.59	123.90
2	A	401	3LM	O2-C-O1	-2.92	116.94	123.90
2	A	401	3LM	C4A-N-CA	2.91	128.58	123.44
2	A	401	3LM	OP4-C5A-C5	-2.88	103.96	109.36
2	B	401	3LM	C4A-C4-C3	2.72	123.60	119.98
2	D	401	3LM	C3-C4-C5	-2.48	116.48	118.73
2	C	401	3LM	C3-C4-C5	-2.42	116.54	118.73
2	D	401	3LM	O2-C-CA	2.35	119.33	114.27
2	C	401	3LM	C4-C3-C2	2.23	123.28	119.91
2	B	401	3LM	O2-C-CA	2.19	118.98	114.27
2	D	401	3LM	C4-C3-C2	2.18	123.20	119.91
2	D	401	3LM	C6-C5-C4	2.15	119.69	118.06
2	B	401	3LM	C3-C4-C5	-2.15	116.78	118.73
2	A	401	3LM	C3-C4-C5	-2.15	116.78	118.73
2	B	401	3LM	OP4-P-OP1	2.07	112.04	106.44
2	B	401	3LM	C6-C5-C4	2.05	119.61	118.06

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	3LM	C3-C4-C4A-N
2	A	401	3LM	C5-C4-C4A-N
2	B	401	3LM	C3-C4-C4A-N
2	B	401	3LM	C5-C4-C4A-N
2	C	401	3LM	C3-C4-C4A-N
2	C	401	3LM	C5-C4-C4A-N
2	D	401	3LM	C5-C4-C4A-N
2	D	401	3LM	C3-C4-C4A-N
2	B	401	3LM	CB-CG-SD-CE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	C	401	3LM	CB-CG-SD-CE
2	D	401	3LM	CB-CG-SD-CE
2	A	401	3LM	CB-CG-SD-CE
2	A	401	3LM	CB-CA-N-C4A
2	B	401	3LM	CB-CA-N-C4A
2	C	401	3LM	CB-CA-N-C4A
2	D	401	3LM	CB-CA-N-C4A
2	A	401	3LM	C-CA-N-C4A
2	B	401	3LM	C-CA-N-C4A
2	C	401	3LM	C-CA-N-C4A
2	D	401	3LM	C-CA-N-C4A


There are no ring outliers.


2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	401	3LM	2	0
2	B	401	3LM	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 3LM C 401









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/398 (98%)	0.10	31 (7%) 18 16	17, 32, 86, 130	0
1	B	392/398 (98%)	-0.20	20 (5%) 33 29	13, 24, 81, 123	0
1	C	392/398 (98%)	-0.15	25 (6%) 25 22	16, 27, 88, 108	0
1	D	398/398 (100%)	-0.04	36 (9%) 15 13	12, 25, 88, 120	0
All	All	1574/1592 (98%)	-0.07	112 (7%) 22 19	12, 27, 87, 130	0

All (112) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	367	TYR	5.5
1	A	366	HIS	5.5
1	C	367	TYR	5.4
1	D	366	HIS	5.4
1	B	55	ALA	5.4
1	B	359	TYR	5.4
1	A	365	ALA	5.3
1	A	55	ALA	5.2
1	D	369	ILE	4.9
1	A	356	HIS	4.8
1	D	365	ALA	4.7
1	D	1	MET	4.3
1	D	52	GLY	4.2
1	D	321	GLY	4.2
1	A	362	GLU	4.2
1	C	360	THR	4.1
1	C	359	TYR	4.0
1	A	368	GLY	3.9
1	B	45	TYR	3.9
1	A	360	THR	3.9
1	A	358	SER	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	50	PHE	3.9
1	D	55	ALA	3.8
1	C	358	SER	3.7
1	A	359	TYR	3.7
1	B	356	HIS	3.7
1	C	369	ILE	3.6
1	C	361	PRO	3.5
1	D	45	TYR	3.5
1	B	357	SER	3.4
1	B	358	SER	3.4
1	C	355	THR	3.4
1	C	365	ALA	3.3
1	B	369	ILE	3.3
1	C	51	ALA	3.3
1	C	55	ALA	3.2
1	B	362	GLU	3.2
1	C	47	ALA	3.2
1	A	357	SER	3.2
1	C	58	PHE	3.2
1	A	369	ILE	3.1
1	C	62	ILE	3.1
1	D	51	ALA	3.1
1	D	324	ALA	3.0
1	C	363	GLU	3.0
1	A	352	ALA	3.0
1	A	363	GLU	2.9
1	D	355	THR	2.9
1	B	366	HIS	2.9
1	B	351	PRO	2.8
1	D	398	ALA	2.8
1	B	360	THR	2.8
1	A	355	THR	2.8
1	C	366	HIS	2.8
1	D	58	PHE	2.8
1	D	370	SER	2.7
1	C	45	TYR	2.7
1	D	372	GLY	2.7
1	A	361	PRO	2.7
1	A	367	TYR	2.6
1	C	356	HIS	2.6
1	D	43	VAL	2.6
1	B	361	PRO	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	50	PHE	2.6
1	A	62	ILE	2.6
1	A	398	ALA	2.6
1	A	52	GLY	2.5
1	B	355	THR	2.5
1	B	365	ALA	2.5
1	D	371	GLU	2.5
1	D	41	PRO	2.5
1	D	364	ARG	2.5
1	A	163	ASN	2.4
1	D	360	THR	2.4
1	B	353	SER	2.4
1	B	367	TYR	2.4
1	A	353	SER	2.4
1	D	285	GLN	2.4
1	B	352	ALA	2.3
1	A	59	TYR	2.3
1	A	56	GLY	2.3
1	D	2	HIS	2.3
1	A	53	GLU	2.2
1	A	351	PRO	2.2
1	D	322	ILE	2.2
1	D	48	ALA	2.2
1	B	161	ASN	2.2
1	A	364	ARG	2.2
1	D	320	GLY	2.2
1	D	325	GLY	2.2
1	A	51	ALA	2.1
1	D	42	THR	2.1
1	A	350	HIS	2.1
1	A	7	LEU	2.1
1	C	354	MET	2.1
1	C	368	GLY	2.1
1	D	47	ALA	2.1
1	D	49	CYS	2.1
1	B	46	GLY	2.1
1	C	44	GLU	2.1
1	D	352	ALA	2.1
1	C	163	ASN	2.1
1	A	46	GLY	2.1
1	D	62	ILE	2.1
1	A	48	ALA	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	398	ALA	2.0
1	D	53	GLU	2.0
1	C	57	HIS	2.0
1	B	52	GLY	2.0
1	C	43	VAL	2.0
1	D	363	GLU	2.0
1	D	63	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	3LM	D	401	24/24	0.93	0.12	21,30,61,71	0
2	3LM	B	401	24/24	0.94	0.12	24,35,66,67	0
2	3LM	C	401	24/24	0.94	0.12	24,35,73,83	0
2	3LM	A	401	24/24	0.94	0.12	24,40,70,79	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around 3LM D 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3LM B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3LM C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 3LM A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.