



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 09:27 PM EDT

PDB ID : 5X2X / pdb_00005x2x
Title : Crystal structure of Pseudomonas putida methionine gamma-lyase wild type with L-homocysteine intermediates
Authors : Shiba, T.; Sato, D.; Harada, S.
Deposited on : 2017-02-02
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	7% 89% 9% .
1	B	398	3% 87% 10% ..
1	C	398	5% 91% 7% ..
1	D	398	8% 91% 8% .

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 12365 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-methionine gamma-lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	0	0
			2951	1863	522	549	17			
1	B	392	Total	C	N	O	S	0	0	0
			2951	1863	522	549	17			
1	C	392	Total	C	N	O	S	0	1	0
			2960	1868	524	551	17			
1	D	398	Total	C	N	O	S	0	1	0
			3006	1895	535	558	18			

- Molecule 2 is (2E)-2-[[[(1E)-{3-hydroxy-2-methyl-5-[(phosphonooxy)methyl]pyridin-4-yl}methylidene]amino]but-2-enoic acid (CCD ID: 4LM) (formula: C₁₂H₁₅N₂O₇P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			22	12	2	7	1		
2	B	1	Total	C	N	O	P	0	0
			22	12	2	7	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			22	12	2	7	1		
2	D	1	Total	C	N	O	P	0	0
			22	12	2	7	1		

- Molecule 3 is HYDROSULFURIC ACID (CCD ID: H2S) (formula: H₂S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	S	0	0
			1	1		
3	B	1	Total	S	0	0
			1	1		
3	C	1	Total	S	0	0
			1	1		
3	D	1	Total	S	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	71	Total	O	0	0
			71	71		
4	B	126	Total	O	0	0
			126	126		
4	C	95	Total	O	0	0
			95	95		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	113	Total 113	O 113	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: L-methionine gamma-lyase




- Molecule 1: L-methionine gamma-lyase




- Molecule 1: L-methionine gamma-lyase



- Molecule 1: L-methionine gamma-lyase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	154.41Å 152.88Å 80.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 20.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.7 (20.00-2.00) 96.7 (20.00-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.34 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.179 , 0.206 0.188 , 0.213	Depositor DCC
R_{free} test set	6310 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.099	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 33.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.013 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12365	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 4LM, H2S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.79	0/3015	0.91	0/4093
1	B	0.85	0/3015	0.93	1/4093 (0.0%)
1	C	0.84	0/3024	0.93	0/4105
1	D	0.85	0/3072	0.88	0/4168
All	All	0.83	0/12126	0.92	1/16459 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	371	GLU	N-CA-C	-8.03	99.94	110.53

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	207	HIS	Peptide
1	C	207	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2951	0	2921	17	0
1	B	2951	0	2921	28	0
1	C	2960	0	2928	14	0
1	D	3006	0	2973	15	0
2	A	22	0	11	1	0
2	B	22	0	12	0	0
2	C	22	0	12	2	0
2	D	22	0	11	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	71	0	0	1	0
4	B	126	0	0	3	0
4	C	95	0	0	2	0
4	D	113	0	0	1	0
All	All	12365	0	11789	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:112:THR:HG21	1:B:163:ASN:HD21	1.49	0.76
1:A:281:ALA:HA	1:A:289:ILE:HD11	1.68	0.74
1:C:112:THR:HG22	1:C:367:TYR:O	1.89	0.72
1:C:169:ILE:H	1:C:306:GLN:HE22	1.38	0.71
1:B:349:GLN:HE22	1:B:355:THR:CG2	2.05	0.69
1:C:211:LYS:HZ1	2:C:401:4LM:C4A	2.06	0.69
1:B:99:THR:HG21	1:B:234:ILE:HA	1.74	0.68
1:D:112:THR:HG21	1:D:163:ASN:HD21	1.58	0.68
1:B:349:GLN:HE22	1:B:355:THR:HG21	1.60	0.66
1:C:329:MET:HE1	1:C:349:GLN:C	2.21	0.66
1:B:166:MET:H	1:B:299:GLN:HE22	1.45	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:43:VAL:H	1:D:330:ASN:HD21	1.46	0.64
1:A:329:MET:HE1	1:A:350:HIS:HB2	1.81	0.60
1:C:62:ILE:HA	4:C:529:HOH:O	2.04	0.58
1:B:95:SER:O	1:B:99:THR:CG2	2.53	0.57
1:A:349:GLN:HE22	1:A:355:THR:CG2	2.18	0.56
1:A:349:GLN:HE22	1:A:355:THR:HG21	1.71	0.54
1:B:355:THR:HG23	1:B:356:HIS:ND1	2.24	0.53
1:A:43:VAL:H	1:B:330:ASN:HD21	1.55	0.53
1:C:211:LYS:NZ	2:C:401:4LM:C4A	2.73	0.52
1:A:242:MET:HE2	1:B:98:TRP:HZ2	1.75	0.52
1:D:349:GLN:HE22	1:D:355:THR:CG2	2.23	0.51
1:B:62:ILE:HA	4:B:597:HOH:O	2.10	0.51
1:B:354:MET:HE1	4:B:616:HOH:O	2.10	0.51
1:D:300:TYR:CZ	1:D:304:ARG:HD2	2.45	0.50
1:D:329:MET:HE1	1:D:350:HIS:HB2	1.94	0.50
1:D:349:GLN:HE22	1:D:355:THR:HG22	1.77	0.50
1:C:350:HIS:CE1	1:C:352:ALA:HB3	2.48	0.49
1:D:166:MET:HE3	1:D:297:PHE:CE2	2.48	0.48
1:B:78:GLU:OE2	1:B:207:HIS:CE1	2.66	0.48
4:A:533:HOH:O	1:B:339:VAL:HG22	2.14	0.48
1:B:78:GLU:OE2	1:B:207:HIS:HE1	1.96	0.48
1:B:148:THR:HB	1:B:149:PRO:HD2	1.97	0.47
1:B:349:GLN:HE22	1:B:355:THR:HG22	1.79	0.47
1:D:78:GLU:OE1	1:D:207:HIS:HE1	1.97	0.47
1:A:242:MET:HE2	1:B:98:TRP:CZ2	2.50	0.46
1:B:95:SER:O	1:B:99:THR:HG23	2.14	0.46
1:B:315:ALA:HB1	1:B:373:LEU:HD11	1.96	0.46
1:B:300:TYR:CZ	1:B:304:ARG:HD2	2.51	0.46
1:A:315:ALA:HB1	1:A:373:LEU:HD11	1.98	0.45
1:A:78:GLU:OE2	1:A:207:HIS:CE1	2.70	0.45
1:A:327:ARG:NH2	1:A:397:SER:O	2.50	0.45
1:D:78:GLU:OE1	1:D:207:HIS:CE1	2.70	0.45
1:A:355:THR:HG23	1:A:356:HIS:ND1	2.32	0.45
1:B:166:MET:CE	1:B:303:ALA:HA	2.48	0.44
4:C:510:HOH:O	1:D:37:THR:HG22	2.17	0.44
1:C:326:ARG:HD3	1:D:44:GLU:OE2	2.17	0.44
1:B:349:GLN:NE2	1:B:355:THR:HG21	2.31	0.44
1:C:166:MET:H	1:C:299:GLN:HE22	1.64	0.44
1:B:70:LEU:HD21	1:B:255:LEU:HD23	2.00	0.44
1:C:204:LEU:HD23	1:C:226:GLY:HA3	1.99	0.44
1:B:95:SER:O	1:B:99:THR:HG22	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:250:HIS:HD2	1:C:251:ASP:OD2	2.02	0.43
1:A:98:TRP:HZ2	1:B:242:MET:HE2	1.85	0.42
1:A:161:ASN:HA	1:A:162:PRO:HA	1.90	0.42
1:A:212:TYR:CE1	1:A:342:GLY:HA2	2.54	0.42
1:A:327:ARG:HB3	1:A:397:SER:HA	2.01	0.42
1:D:152:ARG:NE	1:D:152:ARG:HA	2.35	0.42
1:A:43:VAL:HG12	4:B:616:HOH:O	2.20	0.42
1:D:157:GLU:HG2	1:D:186:ASP:HB3	2.02	0.41
1:D:355:THR:HG23	1:D:356:HIS:ND1	2.35	0.41
1:C:37:THR:HG22	4:D:519:HOH:O	2.21	0.41
1:B:212:TYR:CE1	1:B:342:GLY:HA2	2.56	0.41
2:A:401:4LM:O3	2:A:401:4LM:N	2.54	0.41
1:D:350:HIS:HE1	1:D:371:GLU:O	2.04	0.41
1:A:351:PRO:HA	1:A:355:THR:HG22	2.03	0.40
1:B:329:MET:HE1	1:B:350:HIS:HB2	2.02	0.40
1:B:169:ILE:H	1:B:306:GLN:HE22	1.69	0.40
1:C:143:LEU:HD13	1:C:175:ILE:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/398 (98%)	377 (97%)	12 (3%)	1 (0%)	36	35
1	B	390/398 (98%)	379 (97%)	10 (3%)	1 (0%)	36	35
1	C	391/398 (98%)	381 (97%)	8 (2%)	2 (0%)	24	21
1	D	397/398 (100%)	384 (97%)	12 (3%)	1 (0%)	36	35
All	All	1568/1592 (98%)	1521 (97%)	42 (3%)	5 (0%)	36	35

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	340	SER
1	C	340	SER
1	D	340	SER
1	B	340	SER
1	C	211	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/307 (98%)	286 (95%)	16 (5%)	20	18
1	B	302/307 (98%)	289 (96%)	13 (4%)	26	25
1	C	303/307 (99%)	294 (97%)	9 (3%)	36	38
1	D	308/307 (100%)	293 (95%)	15 (5%)	22	20
All	All	1215/1228 (99%)	1162 (96%)	53 (4%)	25	24

All (53) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	22	GLN
1	A	112	THR
1	A	131	LYS
1	A	132	LEU
1	A	152	ARG
1	A	163	ASN
1	A	177	ARG
1	A	263	ASN
1	A	280	LEU
1	A	289	ILE
1	A	304	ARG
1	A	341	LEU
1	A	362	GLU
1	A	366	HIS
1	A	374	VAL
1	B	64	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	99	THR
1	B	108	LEU
1	B	111	ASN
1	B	112	THR
1	B	152	ARG
1	B	166	MET
1	B	213	LEU
1	B	236	LEU
1	B	276	LEU
1	B	294	LEU
1	B	341	LEU
1	B	392	GLN
1	C	22	GLN
1	C	132	LEU
1	C	140	LEU
1	C	141	GLN
1	C	143	LEU
1	C	163	ASN
1	C	285	GLN
1	C	341	LEU
1	C	395	LYS
1	D	4	SER
1	D	6	LYS
1	D	43	VAL
1	D	112	THR
1	D	131	LYS
1	D	133	ARG
1	D	135	VAL
1	D	191	THR
1	D	213	LEU
1	D	322	ILE
1	D	326	ARG
1	D	340	SER
1	D	341	LEU
1	D	370	SER
1	D	394	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (44) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	GLN
1	A	54	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	68	ASN
1	A	134	HIS
1	A	161	ASN
1	A	207	HIS
1	A	263	ASN
1	A	349	GLN
1	A	350	HIS
1	A	391	GLN
1	B	57	HIS
1	B	64	ASN
1	B	68	ASN
1	B	111	ASN
1	B	161	ASN
1	B	207	HIS
1	B	299	GLN
1	B	306	GLN
1	B	330	ASN
1	B	349	GLN
1	B	350	HIS
1	B	392	GLN
1	C	34	GLN
1	C	54	GLN
1	C	68	ASN
1	C	141	GLN
1	C	161	ASN
1	C	165	HIS
1	C	187	ASN
1	C	195	GLN
1	C	250	HIS
1	C	299	GLN
1	C	306	GLN
1	C	349	GLN
1	C	350	HIS
1	D	22	GLN
1	D	34	GLN
1	D	68	ASN
1	D	163	ASN
1	D	187	ASN
1	D	207	HIS
1	D	274	GLN
1	D	330	ASN
1	D	350	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 4 are modelled with single atom - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	4LM	A	401	-	21,22,22	2.39	7 (33%)	26,31,31	1.80	6 (23%)
2	4LM	B	401	-	21,22,22	2.24	6 (28%)	26,31,31	1.63	5 (19%)
2	4LM	D	401	-	21,22,22	1.87	5 (23%)	26,31,31	1.86	6 (23%)
2	4LM	C	401	-	21,22,22	2.25	6 (28%)	26,31,31	1.70	5 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4LM	A	401	-	-	3/15/17/17	0/1/1/1
2	4LM	B	401	-	-	4/15/17/17	0/1/1/1
2	4LM	D	401	-	-	4/15/17/17	0/1/1/1
2	4LM	C	401	-	-	0/15/17/17	0/1/1/1

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	4LM	CA-C	7.32	1.55	1.48
2	B	401	4LM	CA-C	6.87	1.54	1.48
2	C	401	4LM	CA-C	6.58	1.54	1.48
2	D	401	4LM	CA-C	5.51	1.53	1.48
2	A	401	4LM	CA-N	-3.43	1.37	1.41
2	B	401	4LM	C5A-C5	-3.43	1.42	1.50
2	A	401	4LM	C2-N1	3.40	1.39	1.33
2	C	401	4LM	C5A-C5	-3.28	1.42	1.50
2	C	401	4LM	C2-N1	3.14	1.39	1.33
2	A	401	4LM	C5A-C5	-3.09	1.42	1.50
2	B	401	4LM	CA-N	-3.09	1.37	1.41
2	B	401	4LM	C3-C2	-3.05	1.37	1.41
2	D	401	4LM	P-OP2	-2.74	1.44	1.54
2	D	401	4LM	C5A-C5	-2.66	1.44	1.50
2	D	401	4LM	CA-N	-2.56	1.38	1.41
2	C	401	4LM	C2A-C2	2.53	1.54	1.50
2	B	401	4LM	P-OP2	-2.36	1.46	1.54
2	A	401	4LM	P-OP2	-2.34	1.46	1.54
2	C	401	4LM	CA-N	-2.33	1.38	1.41
2	B	401	4LM	C2-N1	2.30	1.38	1.33
2	A	401	4LM	O2-C	-2.24	1.24	1.30
2	C	401	4LM	O1-C	2.19	1.28	1.22
2	D	401	4LM	C2-N1	2.10	1.37	1.33
2	A	401	4LM	C2A-C2	2.03	1.53	1.50

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	4LM	O2-C-CA	5.16	121.68	113.40
2	D	401	4LM	C4-C3-C2	5.10	123.01	120.14
2	C	401	4LM	O2-C-CA	4.95	121.34	113.40
2	B	401	4LM	O2-C-CA	4.45	120.54	113.40
2	D	401	4LM	O2-C-CA	4.24	120.21	113.40
2	A	401	4LM	C4-C3-C2	3.97	122.38	120.14
2	D	401	4LM	C3-C2-N1	-3.83	116.12	120.96
2	C	401	4LM	C3-C2-N1	-3.23	116.89	120.96
2	A	401	4LM	C3-C2-N1	-3.14	117.00	120.96
2	B	401	4LM	C3-C2-N1	-3.06	117.10	120.96
2	C	401	4LM	C4-C3-C2	3.05	121.86	120.14
2	D	401	4LM	C2A-C2-C3	2.90	124.19	120.80
2	B	401	4LM	C4-C3-C2	2.66	121.64	120.14
2	A	401	4LM	C4-C4A-N	-2.63	116.95	123.01

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	4LM	O2-C-O1	-2.60	117.71	123.90
2	A	401	4LM	O2-C-O1	-2.48	117.99	123.90
2	A	401	4LM	C2A-C2-C3	2.26	123.44	120.80
2	C	401	4LM	C2A-C2-N1	2.21	121.81	117.64
2	C	401	4LM	C4-C4A-N	-2.21	117.92	123.01
2	B	401	4LM	C6-N1-C2	2.19	123.18	119.20
2	D	401	4LM	O2-C-O1	-2.17	118.74	123.90
2	D	401	4LM	C4-C4A-N	-2.13	118.10	123.01

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	4LM	O1-C-CA-CB
2	A	401	4LM	O2-C-CA-N
2	A	401	4LM	O2-C-CA-CB
2	D	401	4LM	O1-C-CA-CB
2	D	401	4LM	O2-C-CA-N
2	D	401	4LM	O2-C-CA-CB
2	B	401	4LM	O1-C-CA-CB
2	B	401	4LM	O2-C-CA-CB
2	B	401	4LM	O1-C-CA-N
2	D	401	4LM	O1-C-CA-N
2	B	401	4LM	O2-C-CA-N


There are no ring outliers.


2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	4LM	1	0
2	C	401	4LM	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	392/398 (98%)	0.37	29 (7%)	20 19	15, 32, 62, 86	0
1	B	392/398 (98%)	-0.08	12 (3%)	51 50	15, 24, 47, 73	0
1	C	392/398 (98%)	0.05	18 (4%)	37 36	12, 27, 52, 82	1 (0%)
1	D	398/398 (100%)	0.10	30 (7%)	20 19	13, 24, 57, 90	1 (0%)
All	All	1574/1592 (98%)	0.11	89 (5%)	29 28	12, 27, 56, 90	2 (0%)

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	1	MET	5.6
1	D	361	PRO	4.9
1	C	367	TYR	4.6
1	D	398	ALA	4.5
1	A	366	HIS	4.5
1	D	358	SER	4.3
1	D	322	ILE	4.2
1	B	365	ALA	4.2
1	C	361	PRO	4.1
1	C	352	ALA	4.1
1	D	365	ALA	4.1
1	A	367	TYR	4.0
1	A	55	ALA	3.8
1	A	177	ARG	3.8
1	C	55	ALA	3.7
1	D	326	ARG	3.6
1	D	372	GLY	3.6
1	C	360	THR	3.5
1	A	361	PRO	3.4
1	D	320	GLY	3.4
1	C	398	ALA	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	350	HIS	3.3
1	C	7	LEU	3.3
1	D	319	LYS	3.2
1	C	372	GLY	3.2
1	B	55	ALA	3.1
1	D	43	VAL	3.1
1	D	55	ALA	3.0
1	D	52	GLY	3.0
1	C	45	TYR	3.0
1	A	362	GLU	2.9
1	D	362	GLU	2.9
1	A	398	ALA	2.9
1	D	321	GLY	2.9
1	A	368	GLY	2.9
1	A	150	ALA	2.8
1	B	288	LEU	2.8
1	C	47	ALA	2.8
1	C	368	GLY	2.7
1	C	365	ALA	2.7
1	B	149	PRO	2.7
1	C	362	GLU	2.7
1	A	54	GLN	2.7
1	A	288	LEU	2.7
1	B	360	THR	2.6
1	D	2	HIS	2.6
1	B	368	GLY	2.6
1	B	51	ALA	2.6
1	D	360	THR	2.6
1	C	359	TYR	2.5
1	A	281	ALA	2.5
1	A	365	ALA	2.5
1	D	48	ALA	2.5
1	D	287	GLU	2.5
1	D	45	TYR	2.5
1	B	355	THR	2.5
1	B	47	ALA	2.5
1	A	7	LEU	2.4
1	D	329	MET	2.4
1	A	57	HIS	2.4
1	D	165[A]	HIS	2.4
1	A	112	THR	2.4
1	D	284	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	52	GLY	2.4
1	A	322	ILE	2.4
1	A	51	ALA	2.3
1	C	43	VAL	2.3
1	A	371	GLU	2.3
1	A	284	PRO	2.3
1	A	355	THR	2.3
1	B	382	ASP	2.3
1	A	286	VAL	2.2
1	D	374	VAL	2.2
1	D	316	PHE	2.2
1	A	329	MET	2.2
1	C	351	PRO	2.2
1	D	354	MET	2.2
1	A	369	ILE	2.1
1	D	288	LEU	2.1
1	C	354	MET	2.1
1	A	360	THR	2.1
1	A	141	GLN	2.1
1	D	283	GLN	2.1
1	B	350	HIS	2.1
1	B	357	SER	2.1
1	D	41	PRO	2.1
1	A	325	GLY	2.1
1	D	371	GLU	2.0
1	A	372	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	H2S	A	402	1/1	0.82	0.15	69,69,69,69	0
3	H2S	C	402	1/1	0.83	0.14	71,71,71,71	0
3	H2S	D	402	1/1	0.88	0.10	62,62,62,62	0
3	H2S	B	402	1/1	0.89	0.10	63,63,63,63	0
2	4LM	A	401	22/22	0.95	0.09	21,34,46,46	0
2	4LM	B	401	22/22	0.96	0.08	19,29,46,51	0
2	4LM	C	401	22/22	0.96	0.09	17,32,47,50	0
2	4LM	D	401	22/22	0.97	0.07	16,26,40,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 4LM A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 4LM B 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 4LM C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.