



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:32 AM UTC

PDB ID : 2X4D / pdb_00002x4d
Title : Crystal structure of human phospholysine phosphohistidine inorganic pyrophosphate phosphatase LHPP
Authors : Vollmar, M.; Gileadi, C.; Guo, K.; Savitsky, P.; Muniz, J.R.C.; Yue, W.; Allerton, C.; von Delft, F.; Bountra, C.; Arrowsmith, C.H.; Weigelt, J.; Edwards, A.; Kavanagh, K.L.; Oppermann, U.
Deposited on : 2010-01-29
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	9% 93% 6%
1	B	271	45% 73% 5% 21%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	POP	A	500	-	X	-	-
2	POP	B	500	-	X	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3704 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PHOSPHOLYSINE PHOSPHOHISTIDINE INORGANIC PYROPHOSPHATE PHOSPHATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	2	0
			1986	1249	348	377	12			
1	B	213	Total	C	N	O	S	0	2	0
			1478	930	252	285	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	94	ARG	GLN	SEE REMARK 999	UNP Q9H008
A	160	ALA	LYS	engineered mutation	UNP Q9H008
A	161	ALA	GLU	engineered mutation	UNP Q9H008
B	94	ARG	GLN	SEE REMARK 999	UNP Q9H008
B	160	ALA	LYS	engineered mutation	UNP Q9H008
B	161	ALA	GLU	engineered mutation	UNP Q9H008

- Molecule 2 is PYROPHOSPHATE 2- (CCD ID: POP) (formula: $\text{H}_2\text{O}_7\text{P}_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			5	1	3	1		
4	B	1	Total	C	O	S	0	0
			5	1	3	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	185	Total	O	0	0
			185	185		
6	B	27	Total	O	0	0
			27	27		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PHOSPHOLYSINE PHOSPHOHISTIDINE INORGANIC PYROPHOSPHATE PHOSPHATASE



- Molecule 1: PHOSPHOLYSINE PHOSPHOHISTIDINE INORGANIC PYROPHOSPHATE PHOSPHATASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.21Å 39.41Å 112.92Å 90.00° 91.79° 90.00°	Depositor
Resolution (Å)	40.00 – 1.92 40.00 – 1.92	Depositor EDS
% Data completeness (in resolution range)	94.2 (40.00-1.92) 94.1 (40.00-1.92)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.5.0089	Depositor
R, R_{free}	0.227 , 0.258 (Not available) , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	26.9	Xtriage
Anisotropy	0.554	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-l	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	3704	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: POP, MG, EPE, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	1/2030 (0.0%)	0.96	2/2749 (0.1%)
1	B	0.75	0/1511	0.86	0/2049
All	All	0.92	1/3541 (0.0%)	0.92	2/4798 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	103	HIS	ND1-CE1	5.13	1.37	1.32

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	103	HIS	CB-CG-CD2	-5.81	123.64	131.20
1	A	103	HIS	CB-CG-ND1	5.06	130.29	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1986	0	1947	10	0
1	B	1478	0	1357	9	0
2	A	5	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	5	0	1	0	0
4	B	5	0	1	0	0
5	A	6	0	8	0	0
6	A	185	0	0	1	1
6	B	27	0	0	1	0
All	All	3704	0	3314	17	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:ARG:HD3	1:B:178:TYR:O	1.80	0.80
1:A:178:TYR:O	1:B:157:ARG:HD3	1.91	0.70
1:B:86:ALA:HB2	6:B:2002:HOH:O	1.91	0.69
1:A:162:THR:HG22	6:A:2014:HOH:O	1.98	0.63
1:B:61[B]:ARG:CZ	1:B:85:PRO:HG2	2.33	0.58
1:A:56:GLU:HG2	1:A:64:LEU:HD22	1.87	0.55
1:A:84:ALA:HB3	1:A:85:PRO:HD3	1.89	0.55
1:B:61[B]:ARG:NH1	1:B:85:PRO:HG2	2.22	0.54
1:B:56:GLU:HG2	1:B:64:LEU:HD22	1.92	0.52
1:A:264:LEU:O	1:A:268:ALA:HB3	2.14	0.47
1:B:94:ARG:NH2	1:B:185:GLU:OE1	2.48	0.46
1:A:94:ARG:NH2	1:A:185:GLU:OE1	2.49	0.45
1:B:84:ALA:HB3	1:B:85:PRO:HD3	1.99	0.43
1:B:97:ARG:NH1	1:B:117:ASN:O	2.43	0.43
1:A:55:ASN:O	1:A:56:GLU:C	2.61	0.43
1:A:17:ASP:OD1	2:A:500:POP:P1	2.77	0.43
1:A:97:ARG:NH1	1:A:117:ASN:O	2.50	0.42

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2028:HOH:O	6:A:2141:HOH:O[1_545]	2.04	0.16

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/271 (100%)	263 (97%)	7 (3%)	0	100	100
1	B	203/271 (75%)	197 (97%)	6 (3%)	0	100	100
All	All	473/542 (87%)	460 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/211 (94%)	196 (99%)	2 (1%)	68	64
1	B	135/211 (64%)	133 (98%)	2 (2%)	57	47
All	All	333/422 (79%)	329 (99%)	4 (1%)	63	57

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	TYR
1	A	103	HIS
1	B	23	TYR
1	B	103	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	244	HIS
1	B	244	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	POP	A	500	3	4,4,8	2.29	2 (50%)	6,6,13	2.05	4 (66%)
2	POP	B	500	3	4,4,8	2.22	2 (50%)	6,6,13	2.52	4 (66%)
5	GOL	A	503	-	5,5,5	0.62	0	5,5,5	1.08	1 (20%)
4	EPE	B	502	-	4,4,15	1.50	1 (25%)	6,6,20	1.41	1 (16%)
4	EPE	A	502	-	4,4,15	1.90	1 (25%)	6,6,20	1.04	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	503	-	-	2/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	POP	P1-O	3.89	1.66	1.54
2	B	500	POP	P1-O	3.33	1.64	1.54
4	A	502	EPE	O1S-S	2.96	1.54	1.44
2	B	500	POP	P1-O1	2.91	1.57	1.50
2	A	500	POP	P1-O1	2.35	1.56	1.50
4	B	502	EPE	O2S-S	2.11	1.51	1.44

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	POP	O2-P1-O1	4.02	125.16	110.95
2	B	500	POP	O3-P1-O2	-3.04	98.44	107.91
2	A	500	POP	O3-P1-O2	2.91	116.98	107.91
2	B	500	POP	O-P1-O2	-2.78	99.26	107.91
2	A	500	POP	O3-P1-O1	-2.73	101.31	110.95
4	A	502	EPE	O2S-S-C10	-2.18	98.91	107.88
4	B	502	EPE	O2S-S-O1S	-2.16	104.77	114.80
2	A	500	POP	O-P1-O3	2.16	114.62	107.91
2	A	500	POP	O-P1-O2	-2.10	101.38	107.91
5	A	503	GOL	C3-C2-C1	-2.06	104.26	111.80
2	B	500	POP	O-P1-O3	2.01	114.17	107.91

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	503	GOL	O1-C1-C2-C3
5	A	503	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	POP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	270/271 (99%)	0.77	25 (9%)	14 18	4, 9, 15, 27	2 (0%)
1	B	213/271 (78%)	2.37	122 (57%)	0 0	7, 9, 12, 14	2 (0%)
All	All	483/542 (89%)	1.48	147 (30%)	1 1	4, 9, 13, 27	4 (0%)

All (147) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	252	GLY	7.6
1	B	176	LEU	6.8
1	B	178	TYR	6.5
1	B	179	ALA	6.3
1	B	182	ILE	6.1
1	B	40	VAL	5.9
1	B	180	CYS	5.6
1	B	105	GLY	5.6
1	B	260	ALA	5.5
1	B	139	PHE	5.4
1	B	143	MET	5.3
1	B	33	ILE	4.9
1	B	60	SER	4.8
1	B	62	ALA	4.7
1	B	142	LEU	4.7
1	B	68	LEU	4.5
1	B	150	LEU	4.5
1	B	144	GLU	4.5
1	B	158	TYR	4.2
1	B	132	TYR	4.0
1	A	146	GLU	4.0
1	B	32	ALA	3.9
1	B	136	ASN	3.9
1	B	52	PHE	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	121	VAL	3.7
1	B	53	CYS	3.7
1	B	58	ALA	3.6
1	B	181	GLY	3.6
1	B	26	GLY	3.6
1	B	65	VAL	3.6
1	B	145	LEU	3.6
1	B	257	LEU	3.6
1	B	141	VAL	3.6
1	B	110	PHE	3.5
1	B	36	SER	3.5
1	B	81	THR	3.5
1	B	169	VAL	3.5
1	B	248	VAL	3.5
1	A	263	LEU	3.5
1	B	201	ALA	3.5
1	B	111	ASP	3.4
1	B	202	ILE	3.4
1	A	148	PRO	3.3
1	B	107	ARG	3.3
1	B	35	GLY	3.3
1	B	131	SER	3.3
1	B	151	ILE	3.3
1	B	240	PRO	3.3
1	B	23	TYR	3.2
1	B	34	ALA	3.2
1	A	144	GLU	3.2
1	B	165	LEU	3.2
1	B	195	PHE	3.2
1	B	99	TYR	3.2
1	B	133	GLN	3.1
1	B	97	ARG	3.1
1	B	135	MET	3.1
1	B	177	GLU	3.1
1	B	213	GLY	3.1
1	B	90	ILE	3.0
1	A	117	ASN	3.0
1	B	31	THR	3.0
1	B	209	ALA	3.0
1	A	93	GLU	3.0
1	B	149	VAL	2.9
1	B	186	VAL	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	59	ALA	2.9
1	B	199	LEU	2.9
1	A	162	THR	2.9
1	B	244	HIS	2.9
1	B	140	GLN	2.8
1	B	188	GLY	2.8
1	A	268	ALA	2.8
1	B	175	ALA	2.8
1	B	89	GLN	2.8
1	A	264	LEU	2.8
1	A	116	SER	2.7
1	B	164	GLY	2.7
1	B	100	LEU	2.7
1	B	227	GLY	2.6
1	B	147	LYS	2.6
1	B	167	LEU	2.6
1	A	160	ALA	2.6
1	B	122	VAL	2.6
1	B	187	VAL	2.6
1	B	254	VAL	2.6
1	B	85	PRO	2.5
1	B	243	GLU	2.5
1	B	155	LYS	2.5
1	B	183	LYS	2.5
1	A	101	LEU	2.5
1	B	64	LEU	2.5
1	B	184	ALA	2.5
1	B	194	PHE	2.5
1	B	152	SER	2.5
1	B	247	GLU	2.5
1	B	84	ALA	2.5
1	A	147	LYS	2.5
1	A	258	ALA	2.4
1	A	233	VAL	2.4
1	B	119	ASN	2.4
1	B	241	SER	2.4
1	A	87	ALA	2.4
1	A	230	ALA	2.4
1	B	37	VAL	2.4
1	B	66	GLY	2.4
1	B	130	PHE	2.3
1	B	137	ASN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	106	VAL	2.3
1	B	106	VAL	2.3
1	B	210	VAL	2.3
1	B	226	CYS	2.3
1	B	246	PRO	2.3
1	B	30	GLY	2.3
1	B	234	ARG	2.3
1	B	21	VAL	2.3
1	B	153	LEU	2.3
1	B	171	PRO	2.2
1	B	220	VAL	2.2
1	B	170	GLY	2.2
1	A	260	ALA	2.2
1	B	251	ASP	2.2
1	B	103	HIS	2.2
1	B	91	LEU	2.2
1	B	108[A]	SER	2.2
1	B	67	GLN	2.2
1	B	233	VAL	2.2
1	B	138	ALA	2.2
1	B	206	ALA	2.2
1	B	172	TYR	2.1
1	A	254	VAL	2.1
1	B	203	GLY	2.1
1	A	257	LEU	2.1
1	A	265	LEU	2.1
1	B	15	LEU	2.1
1	B	174	LYS	2.1
1	A	161	ALA	2.1
1	B	258	ALA	2.1
1	B	207	HIS	2.1
1	B	148	PRO	2.1
1	A	96	LEU	2.1
1	B	120	CYS	2.1
1	B	39	ALA	2.1
1	A	80	VAL	2.0
1	B	192	PRO	2.0
1	B	19	SER	2.0
1	B	98	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EPE	A	502	5/15	0.70	0.25	34,35,39,43	0
5	GOL	A	503	6/6	0.71	0.17	25,28,29,32	0
4	EPE	B	502	5/15	0.75	0.32	33,35,38,42	0
2	POP	B	500	5/9	0.80	0.30	28,30,34,35	0
2	POP	A	500	5/9	0.88	0.18	18,23,26,28	0
3	MG	A	501	1/1	0.93	0.08	9,9,9,9	0
3	MG	B	501	1/1	0.95	0.04	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.