



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 10:52 AM UTC

PDB ID : 2X4Q / pdb_00002x4q
Title : Crystal structure of MHC CLass I HLA-A2.1 bound to a photocleavable peptide
Authors : Celie, P.H.N.; Toebe, M.; Rodenko, B.; Ovaa, H.; Perrakis, A.; Schumacher, T.N.M.
Deposited on : 2010-02-02
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	68% 31% .
1	D	275	65% 33% .
2	B	100	78% 21% .
2	E	100	76% 22% .
3	C	9	22% 22% 11% 44%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	9	 A horizontal bar chart showing the quality of chain F. The bar is divided into four segments: green (22%), yellow (22%), orange (33%), and red (22%).

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6736 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	3	0
			2280	1424	417	430	9			
1	D	275	Total	C	N	O	S	0	1	0
			2258	1409	413	427	9			

- Molecule 2 is a protein called BETA-2-MICROGLOBULIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	Se	0	0	0
			837	533	141	159	2	2			
2	E	100	Total	C	N	O	S	Se	0	1	0
			845	538	142	160	2	3			

- Molecule 3 is a protein called HLA-A2.1-RESTRICTED INFLUENZA A MATRIX EPI-TOPE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	Se	0	0	0
			81	55	11	14	1			
3	F	9	Total	C	N	O	Se	0	0	0
			81	55	11	14	1			

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	E	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O		
			6	3	3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	85	Total 85	O 85	0	0
6	B	50	Total 50	O 50	0	0
6	C	2	Total 2	O 2	0	0
6	D	117	Total 117	O 117	0	0
6	E	55	Total 55	O 55	0	0
6	F	3	Total 3	O 3	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2.1



• Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2.1



• Molecule 2: BETA-2-MICROGLOBULIN



• Molecule 2: BETA-2-MICROGLOBULIN





● Molecule 3: HLA-A2.1-RESTRICTED INFLUENZA A MATRIX EPITOPE



● Molecule 3: HLA-A2.1-RESTRICTED INFLUENZA A MATRIX EPITOPE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.03Å 86.49Å 80.47Å 90.00° 90.21° 90.00°	Depositor
Resolution (Å)	32.43 – 1.90 32.43 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.9 (32.43-1.90) 95.8 (32.43-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 1.90Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.167 , 0.206 0.187 , 0.206	Depositor DCC
R_{free} test set	3299 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	31.0	Xtriage
Anisotropy	0.128	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 32.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.327 for h,-k,-l	Xtriage
Reported twinning fraction	0.337 for -h,-k,l	Depositor
Outliers	0 of 64817 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6736	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PRV, GOL, PRQ, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/2347	0.76	2/3184 (0.1%)
1	D	0.43	0/2323	0.80	4/3151 (0.1%)
2	B	0.42	0/859	0.74	0/1159
2	E	0.51	1/867 (0.1%)	0.75	0/1169
3	C	0.32	0/52	0.86	0/64
3	F	0.36	0/52	0.84	0/64
All	All	0.43	1/6500 (0.0%)	0.77	6/8791 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	5
3	F	0	4
All	All	0	9

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	99	MSE	C-OXT	8.65	1.40	1.23

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	269	PRO	CA-N-CD	-8.33	100.34	112.00
1	D	15	PRO	CA-N-CD	-8.28	100.41	112.00
1	A	56	GLY	CA-C-N	5.83	125.21	118.85
1	A	56	GLY	C-N-CA	5.83	125.21	118.85

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	56	GLY	CA-C-N	5.02	125.04	119.32

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	4	GLY	Peptide
3	C	5	PRV	Mainchain
3	C	7	PHE	Peptide,Mainchain
3	C	8	PRQ	Mainchain
3	F	4	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2280	0	2122	73	1
1	D	2258	0	2108	109	0
2	B	837	0	803	19	0
2	E	845	0	813	21	1
3	C	81	0	73	13	0
3	F	81	0	73	15	0
4	B	12	0	12	2	0
4	D	12	0	12	1	0
4	E	12	0	12	0	0
5	B	6	0	8	3	0
6	A	85	0	0	8	0
6	B	50	0	0	0	0
6	C	2	0	0	0	0
6	D	117	0	0	12	0
6	E	55	0	0	4	0
6	F	3	0	0	0	0
All	All	6736	0	6036	226	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 226 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:98:ASP:HA	2:B:99:MSE:C	1.44	1.31
2:B:98:ASP:CA	2:B:99:MSE:OXT	1.89	1.20
2:B:98:ASP:HA	2:B:99:MSE:OXT	1.01	1.16
1:D:267:PRO:O	1:D:268:LYS:HG3	1.49	1.12
1:A:13:SER:HA	1:A:20:PRO:HB3	1.36	1.06

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:GLU:OE1	2:E:20:SER:OG[2_656]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/275 (100%)	259 (94%)	14 (5%)	3 (1%)	11	4
1	D	274/275 (100%)	256 (93%)	16 (6%)	2 (1%)	18	10
2	B	98/100 (98%)	94 (96%)	4 (4%)	0	100	100
2	E	98/100 (98%)	96 (98%)	1 (1%)	1 (1%)	12	5
3	C	5/9 (56%)	1 (20%)	2 (40%)	2 (40%)	0	0
3	F	5/9 (56%)	2 (40%)	1 (20%)	2 (40%)	0	0
All	All	756/768 (98%)	708 (94%)	38 (5%)	10 (1%)	9	3

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	6	VAL
3	F	6	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	F	7	PHE
1	A	180	GLN
3	C	7	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/231 (101%)	232 (99%)	2 (1%)	70	73
1	D	232/231 (100%)	229 (99%)	3 (1%)	61	61
2	B	95/93 (102%)	91 (96%)	4 (4%)	26	19
2	E	96/93 (103%)	95 (99%)	1 (1%)	68	70
3	C	6/5 (120%)	5 (83%)	1 (17%)	2	0
3	F	6/5 (120%)	5 (83%)	1 (17%)	2	0
All	All	669/658 (102%)	657 (98%)	12 (2%)	51	50

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	21	ARG
1	D	115	GLN
3	F	7	PHE
1	D	181	ARG
2	B	1	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	96	GLN
2	E	13	HIS
1	D	115	GLN
2	E	42	ASN
1	D	253	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	PRV	F	5	3	11,13,14	0.98	1 (9%)	9,17,19	1.22	1 (11%)
3	PRQ	F	8	3	14,14,15	1.04	1 (7%)	13,18,20	1.11	0
3	PRV	C	5	3	11,13,14	1.02	1 (9%)	9,17,19	1.36	1 (11%)
3	PRQ	C	8	3	14,14,15	0.98	1 (7%)	13,18,20	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PRV	F	5	3	-	1/6/10/12	0/1/1/1
3	PRQ	F	8	3	-	5/9/11/12	0/1/1/1
3	PRV	C	5	3	-	2/6/10/12	0/1/1/1
3	PRQ	C	8	3	-	6/9/11/12	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	8	PRQ	CAL-NAN	-2.78	1.40	1.45
3	C	8	PRQ	CAL-NAN	-2.59	1.41	1.45
3	C	5	PRV	CD2-N1	-2.56	1.41	1.45
3	F	5	PRV	CD2-N1	-2.38	1.41	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	5	PRV	CD1-CG-CD2	2.25	119.42	117.08
3	C	5	PRV	CD1-CG-CA	-2.25	117.69	120.61

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	5	PRV	CG-CD2-N1-O2
3	F	5	PRV	C-CA-CG-CD2
3	C	8	PRQ	N-CA-CAJ-C
3	C	8	PRQ	CAK-CA-CAJ-C
3	C	8	PRQ	N-CA-CAK-CAH

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	5	PRV	1	0
3	F	8	PRQ	5	0
3	C	5	PRV	2	0
3	C	8	PRQ	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	B	1100	-	5,5,5	0.46	0	5,5,5	0.41	0
4	MES	D	1276	-	12,12,12	2.10	1 (8%)	15,16,16	6.31	10 (66%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MES	B	1099	-	12,12,12	2.14	1 (8%)	15,16,16	6.07	11 (73%)
4	MES	E	1099	-	12,12,12	2.07	1 (8%)	15,16,16	6.09	10 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	1100	-	-	2/4/4/4	-
4	MES	D	1276	-	-	2/6/14/14	0/1/1/1
4	MES	B	1099	-	-	1/6/14/14	0/1/1/1
4	MES	E	1099	-	-	1/6/14/14	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1099	MES	C8-S	-6.85	1.68	1.77
4	D	1276	MES	C8-S	-6.72	1.68	1.77
4	E	1099	MES	C8-S	-6.62	1.68	1.77

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1099	MES	O1S-S-C8	-14.98	84.09	106.73
4	E	1099	MES	O1S-S-C8	-14.49	84.83	106.73
4	D	1276	MES	O1S-S-C8	-13.86	85.78	106.73
4	B	1099	MES	O2S-S-C8	-12.40	87.99	106.73
4	D	1276	MES	O2S-S-C8	-12.24	88.23	106.73

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	1100	GOL	O1-C1-C2-C3
5	B	1100	GOL	O1-C1-C2-O2
4	B	1099	MES	C8-C7-N4-C3
4	D	1276	MES	C8-C7-N4-C3
4	D	1276	MES	C8-C7-N4-C5

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1100	GOL	3	0
4	D	1276	MES	1	0
4	B	1099	MES	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	-0.95	0 100 100	14, 40, 61, 84	3 (1%)
1	D	275/275 (100%)	-0.96	0 100 100	16, 40, 59, 89	1 (0%)
2	B	98/100 (98%)	-1.02	0 100 100	23, 36, 58, 80	0
2	E	98/100 (98%)	-1.10	0 100 100	21, 37, 59, 66	0
3	C	6/9 (66%)	-0.30	0 100 100	31, 44, 88, 105	0
3	F	6/9 (66%)	0.03	0 100 100	40, 57, 97, 100	0
All	All	758/768 (98%)	-0.97	0 100 100	14, 39, 62, 105	4 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PRV	F	5	13/14	0.93	0.10	88,99,105,106	0
3	PRV	C	5	13/14	0.94	0.09	92,113,118,118	0
3	PRQ	F	8	14/15	0.96	0.10	60,99,104,106	0
3	PRQ	C	8	14/15	0.97	0.08	39,75,90,93	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	B	1100	6/6	0.96	0.08	38,44,48,52	0
4	MES	D	1276	12/12	0.97	0.07	44,57,61,63	0
4	MES	B	1099	12/12	0.97	0.07	65,70,82,86	0
4	MES	E	1099	12/12	0.98	0.05	50,54,69,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.