



Full wwPDB NMR Structure Validation Report ⓘ

Mar 30, 2026 – 01:25 AM UTC

PDB ID : 6X46 / pdb_00006x46
BMRB ID : 30754
Title : NMR solution structure of Asterix/Gtsf1 from mouse (CHHC zinc finger domains)
Authors : Ipsaro, J.J.; O'Brien, P.A.; Bhattacharya, S.; Palmer III, A.G.; Joshua-Tor, L.
Deposited on : 2020-05-22

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 81%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	121	35% 6% 26% 34%

2 Ensemble composition and analysis

This entry contains 20 models. Model 15 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:14-A:38 (25)	0.28	15
2	A:48-A:71 (24)	0.28	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 10, 11, 12, 14, 16, 17, 18
2	2, 4, 5, 6, 8, 13, 20
3	3, 15, 19
Single-model clusters	7; 9

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1280 atoms, of which 626 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Gametocyte-specific factor 1.

Mol	Chain	Residues	Atoms						Trace
1	A	80	Total	C	H	N	O	S	0
			1278	402	626	121	124	5	

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	SER	CYS	engineered mutation	UNP Q9DAN6
A	76	SER	CYS	engineered mutation	UNP Q9DAN6
A	100	SER	CYS	engineered mutation	UNP Q9DAN6
A	103	SER	CYS	engineered mutation	UNP Q9DAN6
A	116	GLU	-	expression tag	UNP Q9DAN6
A	117	ASN	-	expression tag	UNP Q9DAN6
A	118	LEU	-	expression tag	UNP Q9DAN6
A	119	TYR	-	expression tag	UNP Q9DAN6
A	120	PHE	-	expression tag	UNP Q9DAN6
A	121	GLN	-	expression tag	UNP Q9DAN6

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	
2	A	2	Total	Zn
			2	2

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Gametocyte-specific factor 1




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: Gametocyte-specific factor 1



4.2.2 Score per residue for model 2

- Molecule 1: Gametocyte-specific factor 1





4.2.3 Score per residue for model 3

- Molecule 1: Gametocyte-specific factor 1


Chain A: 33% 7% 26% 34%



4.2.4 Score per residue for model 4

- Molecule 1: Gametocyte-specific factor 1


Chain A: 34% 7% 26% 34%



4.2.5 Score per residue for model 5


- Molecule 1: Gametocyte-specific factor 1

Chain A: 32% 8% 26% 34%




4.2.6 Score per residue for model 6

- Molecule 1: Gametocyte-specific factor 1




4.2.7 Score per residue for model 7

- Molecule 1: Gametocyte-specific factor 1




4.2.8 Score per residue for model 8

- Molecule 1: Gametocyte-specific factor 1




4.2.9 Score per residue for model 9

- Molecule 1: Gametocyte-specific factor 1




4.2.10 Score per residue for model 10

- Molecule 1: Gametocyte-specific factor 1




4.2.11 Score per residue for model 11

- Molecule 1: Gametocyte-specific factor 1




4.2.12 Score per residue for model 12

- Molecule 1: Gametocyte-specific factor 1




4.2.13 Score per residue for model 13

- Molecule 1: Gametocyte-specific factor 1




4.2.14 Score per residue for model 14

- Molecule 1: Gametocyte-specific factor 1




4.2.15 Score per residue for model 15 (medoid)

- Molecule 1: Gametocyte-specific factor 1




4.2.16 Score per residue for model 16

- Molecule 1: Gametocyte-specific factor 1




4.2.17 Score per residue for model 17

- Molecule 1: Gametocyte-specific factor 1




4.2.18 Score per residue for model 18

- Molecule 1: Gametocyte-specific factor 1




4.2.19 Score per residue for model 19

- Molecule 1: Gametocyte-specific factor 1



4.2.20 Score per residue for model 20

- Molecule 1: Gametocyte-specific factor 1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DGSA-distance geometry simulated annealing*.

Of the 500 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	structure calculation	2.3
ARIA	refinement	2.3
CNS	refinement	1.2

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	886
Number of shifts mapped to atoms	886
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	81%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	403	397	396	5±1
All	All	8100	7940	7920	107

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:30:PHE:N	1:A:31:PRO:HD2	0.53	2.18	8	10
1:A:17:CYS:CB	1:A:23:HIS:HB2	0.52	2.34	11	17
1:A:51:CYS:CB	1:A:57:HIS:HB2	0.50	2.36	2	20
1:A:51:CYS:HB3	1:A:54:ASN:O	0.50	2.06	1	20
1:A:17:CYS:HB3	1:A:20:ASP:O	0.50	2.05	8	15
1:A:20:ASP:HB3	1:A:23:HIS:CE1	0.47	2.43	9	1
1:A:56:ARG:NE	1:A:56:ARG:HA	0.47	2.25	12	1
1:A:61:ARG:O	1:A:64:ILE:HG22	0.46	2.11	8	7
1:A:64:ILE:O	1:A:68:ILE:HG12	0.46	2.10	7	2
1:A:64:ILE:O	1:A:68:ILE:HG13	0.44	2.12	19	4
1:A:20:ASP:OD2	1:A:22:ASN:HB2	0.44	2.12	18	1
1:A:54:ASN:OD1	1:A:56:ARG:HB2	0.44	2.13	3	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:21:LYS:N	1:A:21:LYS:HD2	0.44	2.28	9	1
1:A:49:ALA:HB2	1:A:61:ARG:HG2	0.43	1.88	2	1
1:A:24:GLN:C	1:A:25:ILE:HD12	0.42	2.40	18	3
1:A:51:CYS:SG	1:A:53:PHE:HB2	0.41	2.55	10	2
1:A:67:HIS:O	1:A:71:CYS:HB3	0.41	2.16	9	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	49/121 (40%)	48±1 (97±2%)	1±1 (3±2%)	0±0 (0±0%)	100	100
All	All	980/2420 (40%)	954 (97%)	26 (3%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	45/114 (39%)	45±0 (100±0%)	0±0 (0±0%)	87	98
All	All	900/2280 (39%)	899 (100%)	1 (0%)	87	98

All 1 unique residues with a non-rotameric sidechain are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	21	LYS	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers ⓘ

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 81% for the well-defined parts and 81% for the entire structure.

7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: *starch_output*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	886
Number of shifts mapped to atoms	886
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	79	-0.30 ± 0.15	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	75	0.13 ± 0.06	None needed (< 0.5 ppm)
$^{13}\text{C}'$	68	-0.07 ± 0.09	None needed (< 0.5 ppm)
^{15}N	72	-0.37 ± 0.50	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 81%, i.e. 568 atoms were assigned a chemical shift out of a possible 705. 0 out of 5 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	231/237 (97%)	94/94 (100%)	92/98 (94%)	45/45 (100%)
Sidechain	308/394 (78%)	201/256 (79%)	101/116 (87%)	6/22 (27%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	29/74 (39%)	15/38 (39%)	14/30 (47%)	0/6 (0%)
Overall	568/705 (81%)	310/388 (80%)	207/244 (85%)	51/73 (70%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 81%, i.e. 886 atoms were assigned a chemical shift out of a possible 1100. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	370/388 (95%)	151/154 (98%)	147/160 (92%)	72/74 (97%)
Sidechain	483/621 (78%)	317/398 (80%)	158/194 (81%)	8/29 (28%)
Aromatic	33/91 (36%)	17/46 (37%)	16/37 (43%)	0/8 (0%)
Overall	886/1100 (81%)	485/598 (81%)	321/391 (82%)	80/111 (72%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

