



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:42 AM UTC

PDB ID : 4X5V / pdb_00004x5v
Title : Crystal structure of the post-catalytic nick complex of DNA polymerase lambda with a templating A and incorporated 8-oxo-dGMP
Authors : Burak, M.J.; Guja, K.E.; Garcia-Diaz, M.
Deposited on : 2014-12-05
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	16% 90% 7% ..
2	D	4	75% 25%
3	P	7	86% 14%
4	T	11	82% 18%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 3019 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase lambda.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	0	0	0
			2375	1491	433	440	11			

- Molecule 2 is a DNA chain called DNA (5'-D(P*GP*CP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	4	Total	C	N	O	P	0	0	0
			83	38	16	25	4			

- Molecule 3 is a DNA chain called DNA (5'-D(*CP*AP*GP*TP*AP*CP*(8OG))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	7	Total	C	N	O	P	0	0	0
			142	68	28	40	6			

- Molecule 4 is a DNA chain called DNA (5'-D(*CP*GP*GP*CP*AP*GP*TP*AP*CP*TP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	T	11	Total	C	N	O	P	0	0	0
			224	107	43	64	10			

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			13	6	7		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	133	Total	O	0	0
			133	133		
8	D	1	Total	O	0	0
			1	1		
8	P	22	Total	O	0	0
			22	22		
8	T	24	Total	O	0	0
			24	24		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase lambda




- Molecule 2: DNA (5'-D(P*GP*CP*CP*G)-3')



- Molecule 3: DNA (5'-D(*CP*AP*GP*TP*AP*CP*(8OG))-3')



- Molecule 4: DNA (5'-D(*CP*GP*GP*CP*AP*GP*TP*AP*CP*TP*G)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.01Å 62.84Å 141.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.98 – 2.15 46.98 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.98-2.15) 99.9 (46.98-2.15)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 2.16Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.214 , 0.245 0.220 , 0.251	Depositor DCC
R_{free} test set	2000 reflections (7.16%)	wwPDB-VP
Wilson B-factor (Å ²)	35.9	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 61.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3019	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 8OG, MG, NA, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/2419	0.77	2/3273 (0.1%)
2	D	0.19	0/92	0.35	0/138
3	P	0.25	0/133	0.46	0/203
4	T	0.23	0/251	0.52	0/386
All	All	0.42	0/2895	0.72	2/4000 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	434	HIS	CA-C-N	-5.21	114.45	119.87
1	A	434	HIS	C-N-CA	-5.21	114.45	119.87

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2375	0	2281	16	0
2	D	83	0	45	1	0
3	P	142	0	80	0	0
4	T	224	0	125	2	0
5	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	1	0	0	0	0
7	A	13	0	4	0	0
8	A	133	0	0	5	0
8	D	1	0	0	0	0
8	P	22	0	0	0	0
8	T	24	0	0	1	0
All	All	3019	0	2535	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:394:GLU:OE1	8:A:701:HOH:O	1.99	0.81
1:A:558:ASP:OD1	8:A:702:HOH:O	2.05	0.74
1:A:521:LYS:NZ	4:T:4:DC:OP1	2.20	0.73
1:A:308:ARG:NH1	2:D:1:DG:OP2	2.23	0.71
1:A:574:ASP:OD2	8:A:703:HOH:O	2.07	0.71
1:A:442:GLY:O	1:A:445:SER:OG	2.14	0.61
4:T:8:DA:OP2	8:T:101:HOH:O	2.16	0.60
1:A:454:GLU:OE2	8:A:704:HOH:O	2.17	0.58
1:A:434:HIS:ND1	1:A:439:SER:HB2	2.22	0.54
1:A:434:HIS:ND1	1:A:439:SER:CB	2.73	0.52
1:A:434:HIS:HD2	1:A:435:PRO:CD	2.26	0.49
1:A:315:GLU:OE1	1:A:322:LEU:HD12	2.14	0.47
1:A:434:HIS:HD2	1:A:435:PRO:HD2	1.80	0.47
1:A:251:ASN:N	8:A:716:HOH:O	2.47	0.47
1:A:431:LEU:HD13	1:A:492:ILE:HD11	2.01	0.41
1:A:519:LEU:O	1:A:522:THR:OG1	2.37	0.41
1:A:434:HIS:O	1:A:496:TYR:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/325 (96%)	306 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/273 (85%)	226 (97%)	6 (3%)	40	43

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	263	LEU
1	A	324	LYS
1	A	411	LEU
1	A	439	SER
1	A	506	PHE
1	A	529	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	270	GLN
1	A	284	ASN
1	A	400	GLN
1	A	464	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	8OG	P	7	5,4,3	22,25,26	3.87	13 (59%)	26,37,40	2.08	9 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	8OG	P	7	5,4,3	-	1/7/21/22	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	7	8OG	C3'-C4'	-8.74	1.30	1.53
3	P	7	8OG	O4'-C4'	7.74	1.62	1.45
3	P	7	8OG	C4-N3	5.38	1.46	1.34
3	P	7	8OG	C2-N3	5.34	1.46	1.33
3	P	7	8OG	C8-N7	5.25	1.48	1.38
3	P	7	8OG	O4'-C1'	-4.80	1.31	1.42
3	P	7	8OG	C8-N9	4.10	1.47	1.40
3	P	7	8OG	C2-N2	4.00	1.43	1.34
3	P	7	8OG	C2-N1	3.24	1.45	1.37
3	P	7	8OG	C5-C4	3.05	1.41	1.37
3	P	7	8OG	O3'-C3'	2.80	1.49	1.43
3	P	7	8OG	C5-C6	2.71	1.50	1.41
3	P	7	8OG	C5-N7	2.37	1.41	1.37

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	7	8OG	C2-N3-C4	5.56	121.89	112.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	7	8OG	N9-C4-N3	3.92	131.03	126.13
3	P	7	8OG	O6-C6-C5	-3.25	119.42	127.26
3	P	7	8OG	C5-C6-N1	3.18	120.85	112.13
3	P	7	8OG	N1-C2-N3	-2.94	117.94	123.32
3	P	7	8OG	C5-N7-C8	-2.79	105.62	109.47
3	P	7	8OG	C4-C5-N7	2.65	110.91	106.06
3	P	7	8OG	C2-N1-C6	-2.45	120.66	125.11
3	P	7	8OG	N2-C2-N1	2.11	121.22	116.76

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	P	7	8OG	O4'-C1'-N9-C4

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	CIT	A	603	5	12,12,12	1.31	0	17,17,17	1.39	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	CIT	A	603	5	-	2/16/16/16	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	603	CIT	O6-C6-C3	3.47	119.80	113.14
7	A	603	CIT	O4-C5-C4	2.17	121.22	114.35

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	603	CIT	C3-C4-C5-O4
7	A	603	CIT	C3-C4-C5-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/325 (97%)	0.90	51 (16%) 4 5	20, 44, 82, 93	0
2	D	4/4 (100%)	0.00	0 100 100	47, 51, 51, 54	0
3	P	6/7 (85%)	-0.78	0 100 100	22, 25, 27, 34	0
4	T	11/11 (100%)	-0.07	0 100 100	29, 38, 46, 48	0
All	All	338/347 (97%)	0.83	51 (15%) 5 6	20, 43, 82, 93	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	406	PHE	5.1
1	A	467	ASN	4.6
1	A	255	HIS	4.4
1	A	256	ILE	4.3
1	A	443	ILE	4.1
1	A	314	ILE	3.9
1	A	313	ILE	3.7
1	A	409	GLY	3.7
1	A	441	ARG	3.6
1	A	575	TRP	3.5
1	A	316	ILE	3.4
1	A	317	LEU	3.4
1	A	540	THR	3.4
1	A	297	GLN	3.3
1	A	471	GLN	3.3
1	A	319	SER	3.3
1	A	301	SER	3.2
1	A	468	GLY	3.2
1	A	547	PRO	3.2
1	A	437	GLY	3.1
1	A	328	ILE	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	252	HIS	3.1
1	A	408	SER	3.1
1	A	321	HIS	3.0
1	A	543	CYS	3.0
1	A	302	ILE	2.9
1	A	545	VAL	2.9
1	A	286	LEU	2.7
1	A	311	GLU	2.6
1	A	368	SER	2.6
1	A	442	GLY	2.5
1	A	251	ASN	2.5
1	A	285	ALA	2.5
1	A	440	HIS	2.5
1	A	296	TYR	2.4
1	A	438	ARG	2.4
1	A	444	PHE	2.4
1	A	318	GLU	2.4
1	A	308	ARG	2.4
1	A	257	THR	2.4
1	A	535	ALA	2.3
1	A	287	LYS	2.3
1	A	511	HIS	2.2
1	A	445	SER	2.2
1	A	470	GLN	2.1
1	A	434	HIS	2.1
1	A	410	LEU	2.1
1	A	303	PRO	2.1
1	A	537	VAL	2.1
1	A	305	ILE	2.0
1	A	411	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	8OG	P	7	23/24	0.93	0.12	24,24,40,41	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NA	A	601	1/1	0.85	0.16	79,79,79,79	0
7	CIT	A	603	13/13	0.89	0.11	40,44,53,58	0
6	MG	A	602	1/1	0.96	0.04	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.