



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 02:45 PM UTC

PDB ID : 7X51 / pdb_00007x51
Title : Crystal structure of Bacteroides thetaiotaomicron glutamate decarboxylase
BTGAD-PLP-GUA complex
Authors : Liu, S.; Du, G.; Wang, L.; Wen, B.; Xin, F.
Deposited on : 2022-03-03
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	 87% 9% 5%
1	B	481	 87% 7% 5%
1	C	481	 86% 9% 5%
1	D	481	 87% 8% 5%
1	E	481	 87% 8% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	481	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GUA	D	503	-	X	-	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 23995 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate decarboxylase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	458	Total	C	N	O	S	4	0	0
			3663	2371	600	672	20			
1	B	459	Total	C	N	O	S	4	0	0
			3683	2381	602	680	20			
1	C	459	Total	C	N	O	S	4	0	0
			3667	2370	599	678	20			
1	D	458	Total	C	N	O	S	4	0	0
			3673	2375	600	677	21			
1	E	458	Total	C	N	O	S	4	0	0
			3659	2367	598	674	20			
1	F	459	Total	C	N	O	S	4	0	0
			3667	2371	600	676	20			

- Molecule 2 is MALONATE ION (CCD ID: MLI) (formula: $C_3H_2O_4$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 7 3 4	0	0
2	B	1	Total C O 7 3 4	0	0
2	C	1	Total C O 7 3 4	0	0
2	D	1	Total C O 7 3 4	0	0
2	E	1	Total C O 7 3 4	0	0

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O P 15 8 1 5 1	0	0
3	B	1	Total C N O P 15 8 1 5 1	0	0
3	C	1	Total C N O P 15 8 1 5 1	0	0
3	D	1	Total C N O P 15 8 1 5 1	0	0
3	E	1	Total C N O P 15 8 1 5 1	0	0
3	F	1	Total C N O P 15 8 1 5 1	0	0

- Molecule 4 is GLUTARIC ACID (CCD ID: GUA) (formula: $C_5H_8O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			9	5	4		
4	B	1	Total	C	O	0	0
			9	5	4		
4	C	1	Total	C	O	0	0
			9	5	4		
4	D	1	Total	C	O	0	0
			9	5	4		
4	E	1	Total	C	O	0	0
			9	5	4		
4	F	1	Total	C	O	0	0
			9	5	4		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	342	Total	O	0	0
			342	342		
5	B	364	Total	O	0	0
			364	364		
5	C	307	Total	O	0	0
			307	307		
5	D	304	Total	O	0	0
			304	304		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	262	Total 262	O 262	0	0
5	F	225	Total 225	O 225	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Glutamate decarboxylase

Chain A: 




• Molecule 1: Glutamate decarboxylase

Chain B: 



• Molecule 1: Glutamate decarboxylase


Chain C: 





• Molecule 1: Glutamate decarboxylase

Chain D: 








● Molecule 1: Glutamate decarboxylase



● Molecule 1: Glutamate decarboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	109.82Å 131.42Å 109.84Å 90.00° 100.19° 90.00°	Depositor
Resolution (Å)	36.04 – 2.00 36.04 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.04-2.00) 99.9 (36.04-2.00)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.00Å)	Xtriage
Refinement program	PHENIX (1.19.1_4122: ???)	Depositor
R, R_{free}	0.201 , 0.239 0.201 , 0.239	Depositor DCC
R_{free} test set	10067 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.454	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 26.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.407 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	23995	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLI, PLP, GUA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/3762	0.53	0/5117
1	B	0.37	1/3782 (0.0%)	0.55	3/5143 (0.1%)
1	C	0.34	0/3766	0.53	3/5126 (0.1%)
1	D	0.33	0/3772	0.53	2/5130 (0.0%)
1	E	0.31	0/3758	0.50	0/5114
1	F	0.31	0/3766	0.50	0/5125
All	All	0.34	1/22606 (0.0%)	0.52	8/30755 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	273	HIS	C-N	-5.30	1.26	1.33

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	274	LYS	O-C-N	6.29	127.68	120.58
1	D	410	SER	CA-C-N	6.01	133.01	121.54
1	D	410	SER	C-N-CA	6.01	133.01	121.54
1	B	410	SER	CA-C-N	5.61	132.25	121.54
1	B	410	SER	C-N-CA	5.61	132.25	121.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3663	0	3605	28	0
1	B	3683	0	3626	29	0
1	C	3667	0	3589	30	0
1	D	3673	0	3616	28	0
1	E	3659	0	3587	28	0
1	F	3667	0	3593	29	0
2	A	7	0	2	0	0
2	B	7	0	2	0	0
2	C	7	0	2	0	0
2	D	7	0	2	0	0
2	E	7	0	2	0	0
3	A	15	0	7	2	0
3	B	15	0	7	2	0
3	C	15	0	7	2	0
3	D	15	0	7	2	0
3	E	15	0	7	2	0
3	F	15	0	7	2	0
4	A	9	0	0	0	0
4	B	9	0	0	0	0
4	C	9	0	0	0	0
4	D	9	0	0	0	0
4	E	9	0	0	0	0
4	F	9	0	0	0	0
5	A	342	0	0	3	0
5	B	364	0	0	3	0
5	C	307	0	0	3	0
5	D	304	0	0	1	0
5	E	262	0	0	2	0
5	F	225	0	0	3	0
All	All	23995	0	21668	151	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 151 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:191:GLU:HA	1:E:194:LYS:HE2	1.63	0.80
1:F:123:SER:HB2	3:F:501:PLP:O4P	1.85	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:309:THR:HG22	1:C:311:VAL:HG23	1.68	0.75
1:E:123:SER:HB2	3:E:502:PLP:O4P	1.86	0.75
1:A:402:MET:HG2	1:B:87:GLU:HG3	1.70	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	456/481 (95%)	442 (97%)	14 (3%)	0	100	100
1	B	457/481 (95%)	442 (97%)	15 (3%)	0	100	100
1	C	457/481 (95%)	443 (97%)	14 (3%)	0	100	100
1	D	456/481 (95%)	442 (97%)	14 (3%)	0	100	100
1	E	456/481 (95%)	440 (96%)	16 (4%)	0	100	100
1	F	457/481 (95%)	442 (97%)	15 (3%)	0	100	100
All	All	2739/2886 (95%)	2651 (97%)	88 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	390/414 (94%)	390 (100%)	0	100	100
1	B	394/414 (95%)	393 (100%)	1 (0%)	86	91
1	C	390/414 (94%)	390 (100%)	0	100	100
1	D	393/414 (95%)	392 (100%)	1 (0%)	86	91
1	E	389/414 (94%)	389 (100%)	0	100	100
1	F	390/414 (94%)	390 (100%)	0	100	100
All	All	2346/2484 (94%)	2344 (100%)	2 (0%)	88	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	433	MET
1	D	301	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	310	GLN
1	F	399	HIS
1	C	302	ASN
1	C	314	ASN
1	C	398	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	MLI	C	501	-	6,6,6	1.30	0	7,7,7	1.24	0
2	MLI	E	501	-	6,6,6	1.44	0	7,7,7	1.40	1 (14%)
3	PLP	B	502	1	15,15,16	1.12	1 (6%)	21,22,23	1.59	3 (14%)
2	MLI	A	501	-	6,6,6	1.49	0	7,7,7	1.43	1 (14%)
3	PLP	C	502	1	15,15,16	1.11	1 (6%)	21,22,23	1.57	4 (19%)
3	PLP	E	502	1	15,15,16	1.12	1 (6%)	21,22,23	1.58	3 (14%)
4	GUA	C	503	-	8,8,8	1.11	0	9,9,9	2.05	3 (33%)
4	GUA	D	503	-	8,8,8	1.26	0	9,9,9	2.17	5 (55%)
3	PLP	A	502	1	15,15,16	1.12	1 (6%)	21,22,23	1.58	3 (14%)
2	MLI	B	501	-	6,6,6	1.24	0	7,7,7	1.28	0
3	PLP	F	501	1	15,15,16	1.12	1 (6%)	21,22,23	1.56	3 (14%)
4	GUA	A	503	-	8,8,8	1.35	0	9,9,9	1.29	1 (11%)
4	GUA	B	503	-	8,8,8	1.18	0	9,9,9	2.26	3 (33%)
4	GUA	E	503	-	8,8,8	1.06	0	9,9,9	1.96	3 (33%)
4	GUA	F	502	-	8,8,8	1.10	0	9,9,9	1.87	2 (22%)
2	MLI	D	501	-	6,6,6	1.28	0	7,7,7	1.31	0
3	PLP	D	502	1	15,15,16	1.13	1 (6%)	21,22,23	1.56	4 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLI	C	501	-	-	2/4/4/4	-
2	MLI	E	501	-	-	2/4/4/4	-
3	PLP	B	502	1	-	0/6/6/8	0/1/1/1
2	MLI	A	501	-	-	4/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PLP	C	502	1	-	0/6/6/8	0/1/1/1
3	PLP	E	502	1	-	1/6/6/8	0/1/1/1
4	GUA	C	503	-	-	1/6/6/6	-
4	GUA	D	503	-	-	6/6/6/6	-
3	PLP	A	502	1	-	1/6/6/8	0/1/1/1
2	MLI	B	501	-	-	0/4/4/4	-
3	PLP	F	501	1	-	0/6/6/8	0/1/1/1
4	GUA	A	503	-	-	3/6/6/6	-
4	GUA	B	503	-	-	1/6/6/6	-
4	GUA	E	503	-	-	3/6/6/6	-
4	GUA	F	502	-	-	5/6/6/6	-
2	MLI	D	501	-	-	2/4/4/4	-
3	PLP	D	502	1	-	0/6/6/8	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	PLP	C2-N1	2.65	1.38	1.33
3	E	502	PLP	C2-N1	2.65	1.38	1.33
3	A	502	PLP	C2-N1	2.63	1.38	1.33
3	F	501	PLP	C2-N1	2.63	1.38	1.33
3	D	502	PLP	C2-N1	2.62	1.38	1.33

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	503	GUA	C3-C2-C1	-4.20	103.54	114.51
4	E	503	GUA	C3-C2-C1	-3.70	104.84	114.51
4	C	503	GUA	C3-C2-C1	-3.58	105.17	114.51
3	B	502	PLP	C5-C6-N1	-3.54	118.08	123.83
3	F	501	PLP	C5-C6-N1	-3.52	118.10	123.83

There are no chirality outliers.

5 of 31 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	503	GUA	C2-C3-C4-C5
4	F	502	GUA	C2-C3-C4-C5
4	D	503	GUA	C1-C2-C3-C4
2	A	501	MLI	C3-C1-C2-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	501	MLI	C2-C1-C3-O9

There are no ring outliers.


6 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	PLP	2	0
3	C	502	PLP	2	0
3	E	502	PLP	2	0
3	A	502	PLP	2	0
3	F	501	PLP	2	0
3	D	502	PLP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand PLP C 502




Ligand PLP E 502










Ligand PLP A 502




Ligand PLP F 501









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	458/481 (95%)	-1.46	0 100 100	11, 25, 39, 60	1 (0%)
1	B	459/481 (95%)	-1.47	0 100 100	11, 24, 39, 62	1 (0%)
1	C	459/481 (95%)	-1.43	0 100 100	11, 28, 46, 68	1 (0%)
1	D	458/481 (95%)	-1.37	0 100 100	11, 31, 50, 69	1 (0%)
1	E	458/481 (95%)	-1.39	0 100 100	11, 33, 50, 63	1 (0%)
1	F	459/481 (95%)	-1.36	0 100 100	11, 34, 53, 72	1 (0%)
All	All	2751/2886 (95%)	-1.41	0 100 100	11, 29, 49, 72	6 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MLI	D	501	7/7	0.96	0.07	51,53,59,59	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MLI	E	501	7/7	0.96	0.05	52,54,55,57	0
2	MLI	C	501	7/7	0.97	0.07	29,35,39,43	0
2	MLI	A	501	7/7	0.97	0.04	35,37,42,43	0
2	MLI	B	501	7/7	0.97	0.07	31,35,37,38	0
3	PLP	F	501	15/16	0.97	0.06	18,21,24,24	0
3	PLP	E	502	15/16	0.98	0.05	18,21,24,24	0
4	GUA	A	503	9/9	0.98	0.04	17,21,28,30	0
4	GUA	E	503	9/9	0.98	0.05	25,29,41,42	0
4	GUA	F	502	9/9	0.98	0.05	27,32,41,45	0
3	PLP	B	502	15/16	0.99	0.04	18,21,24,24	0
3	PLP	C	502	15/16	0.99	0.04	18,21,24,24	0
4	GUA	B	503	9/9	0.99	0.04	19,21,29,30	0
4	GUA	C	503	9/9	0.99	0.06	21,25,35,39	0
4	GUA	D	503	9/9	0.99	0.04	25,28,36,39	0
3	PLP	D	502	15/16	0.99	0.04	18,21,24,24	0
3	PLP	A	502	15/16	0.99	0.03	18,21,24,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around PLP F 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PLP E 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GUA A 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GUA E 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GUA F 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PLP B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PLP C 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GUA B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GUA C 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GUA D 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PLP D 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around PLP A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.