



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:42 PM UTC

PDB ID : 7X5D / pdb_00007x5d
Title : Crystal Structure of the K316C mutant of Human Lamin A/C Coil 2 (residues 244-340)
Authors : Ahn, J.; Jo, I.; Ha, N.-C.
Deposited on : 2022-03-04
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	103	15% 77% 19% .
1	B	103	20% 80% 12% 9%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lamin-A/C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	99	Total	C	N	O	S	0	0	0
			801	482	153	164	2			
1	B	94	Total	C	N	O	S	0	0	0
			758	454	148	155	1			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	237	GLY	-	expression tag	UNP P02545
A	238	ALA	-	expression tag	UNP P02545
A	239	MET	-	expression tag	UNP P02545
A	240	ASP	-	expression tag	UNP P02545
A	241	PRO	-	expression tag	UNP P02545
A	242	GLU	-	expression tag	UNP P02545
A	243	PHE	-	expression tag	UNP P02545
A	316	CYS	LYS	engineered mutation	UNP P02545
B	237	GLY	-	expression tag	UNP P02545
B	238	ALA	-	expression tag	UNP P02545
B	239	MET	-	expression tag	UNP P02545
B	240	ASP	-	expression tag	UNP P02545
B	241	PRO	-	expression tag	UNP P02545
B	242	GLU	-	expression tag	UNP P02545
B	243	PHE	-	expression tag	UNP P02545
B	316	CYS	LYS	engineered mutation	UNP P02545



- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	66	Total	O	0	0
			66	66		
2	B	58	Total	O	0	0
			58	58		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lamin-A/C



- Molecule 1: Lamin-A/C



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	29.79Å 40.97Å 140.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.10 – 1.82 20.10 – 1.82	Depositor EDS
% Data completeness (in resolution range)	86.2 (20.10-1.82) 86.1 (20.10-1.82)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.46 (at 1.82Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.239 , 0.279 0.239 , 0.279	Depositor DCC
R_{free} test set	695 reflections (4.30%)	wwPDB-VP
Wilson B-factor (Å ²)	16.7	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 53.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	1683	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.21	0/806	0.37	0/1078
1	B	0.20	0/761	0.36	0/1017
All	All	0.20	0/1567	0.36	0/2095

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	801	0	791	11	0
1	B	758	0	756	9	0
2	A	66	0	0	1	0
2	B	58	0	0	1	0
All	All	1683	0	1547	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:ASP:OD1	1:A:275:ARG:NH2	2.27	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:261:LYS:HE2	1:B:261:LYS:HA	1.78	0.66
1:B:329:ARG:NH1	2:B:403:HOH:O	2.36	0.59
1:A:319:LYS:HB2	1:B:288:ALA:HB2	1.88	0.56
1:B:283:ASN:OD1	1:B:284:LEU:N	2.43	0.52
1:A:332:ASP:O	1:A:336:ARG:HG3	2.12	0.50
1:A:332:ASP:OD2	1:A:335:ARG:NH1	2.43	0.50
1:A:247:GLU:O	1:A:251:GLN:HG3	2.12	0.49
1:B:292:LEU:O	1:B:296:ARG:HG2	2.15	0.47
1:B:275:ARG:HH11	1:B:275:ARG:HB3	1.80	0.45
1:B:265:LYS:HE2	1:B:265:LYS:HA	2.00	0.44
1:B:333:THR:O	1:B:337:LEU:HD13	2.17	0.44
1:A:273:ASN:ND2	2:A:401:HOH:O	2.41	0.44
1:A:266:THR:O	1:A:270:LYS:HG2	2.19	0.43
1:A:240:ASP:HB3	1:A:241:PRO:HD3	2.02	0.42
1:A:304:ALA:O	1:A:308:GLN:HG3	2.20	0.42
1:A:327:LEU:O	1:A:331:ARG:HG3	2.19	0.41
1:A:323:LEU:HD21	1:B:285:VAL:HG22	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/103 (94%)	97 (100%)	0	0	100	100
1	B	92/103 (89%)	92 (100%)	0	0	100	100
All	All	189/206 (92%)	189 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/87 (99%)	84 (98%)	2 (2%)	44	28
1	B	81/87 (93%)	81 (100%)	0	100	100
All	All	167/174 (96%)	165 (99%)	2 (1%)	63	54

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	253	GLU
1	A	333	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	252	HIS
1	A	258	GLN
1	A	305	GLN
1	B	251	GLN
1	B	294	GLN
1	B	305	GLN
1	B	310	GLN
1	B	312	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	99/103 (96%)	0.81	15 (15%) 5 5	5, 27, 64, 77	0
1	B	94/103 (91%)	1.17	21 (22%) 2 2	6, 40, 66, 79	0
All	All	193/206 (93%)	0.98	36 (18%) 3 4	5, 36, 65, 79	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	337	LEU	6.3
1	B	285	VAL	5.1
1	B	282	SER	4.8
1	A	331	ARG	4.7
1	B	284	LEU	4.6
1	B	273	ASN	4.5
1	B	287	ALA	4.5
1	B	286	GLY	3.9
1	B	277	SER	3.3
1	B	283	ASN	3.2
1	B	278	ALA	3.1
1	B	281	ASN	3.1
1	B	298	ARG	3.1
1	A	312	GLN	2.8
1	A	318	ALA	2.8
1	A	321	ARG	2.8
1	A	315	ALA	2.7
1	B	289	HIS	2.7
1	A	326	SER	2.6
1	B	279	GLU	2.6
1	A	328	ALA	2.4
1	A	329	ARG	2.3
1	B	269	ALA	2.3
1	A	322	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	274	ALA	2.3
1	A	239	MET	2.2
1	A	308	GLN	2.2
1	B	267	TYR	2.1
1	A	332	ASP	2.1
1	B	337	LEU	2.1
1	B	244	ALA	2.1
1	B	280	ARG	2.1
1	A	323	LEU	2.1
1	B	288	ALA	2.1
1	A	319	LYS	2.1
1	B	304	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.