



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 04:51 PM UTC

PDB ID : 8X5V / pdb_00008x5v
Title : BICas9-sgRNA-target DNA complex
Authors : Nakane, T.; Nakagawa, R.; Yamashita, K.; Nishimasu, H.; Nureki, O.
Deposited on : 2023-11-19
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)
RNA backbone	3983	1000 (2.36-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	933	5% 91% 5% .
2	B	110	% 80% 15% . .
3	C	28	93% 7%
4	D	8	75% 25%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 10610 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BlCas9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	892	Total	C	N	O	S	0	5	0
			7172	4549	1269	1334	20			

- Molecule 2 is a RNA chain called RNA (110-mer).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	106	Total	C	N	O	P	0	0	0
			2258	1007	396	749	106			

- Molecule 3 is a DNA chain called DNA (28-mer).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	28	Total	C	N	O	P	0	0	0
			567	272	103	165	27			

- Molecule 4 is a DNA chain called DNA (8-mer).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	8	Total	C	N	O	P	0	0	0
			158	78	27	46	7			

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Cl	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	171	Total	O	0	0
			171	171		
8	B	105	Total	O	0	0
			105	105		
8	C	40	Total	O	0	0
			40	40		
8	D	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: BlCas9




- Molecule 2: RNA (110-mer)



- Molecule 3: DNA (28-mer)



- Molecule 4: DNA (8-mer)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	144.75Å 99.21Å 119.30Å 90.00° 97.09° 90.00°	Depositor
Resolution (Å)	118.39 – 2.00 118.39 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (118.39-2.00) 99.9 (118.39-2.00)	Depositor EDS
R_{merge}	0.25	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.198 , 0.231 0.204 , 0.235	Depositor DCC
R_{free} test set	5612 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	39.8	Xtriage
Anisotropy	0.218	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10610	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/7324	1.03	4/9882 (0.0%)
2	B	0.58	0/2521	1.12	9/3925 (0.2%)
3	C	0.41	0/635	1.08	2/977 (0.2%)
4	D	0.42	0/176	1.15	0/269
All	All	0.57	0/10656	1.06	15/15053 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	52	A	O3'-P-O5'	-8.56	91.17	104.00
2	B	83	C	O3'-P-O5'	-7.22	93.16	104.00
2	B	7	U	O3'-P-O5'	-6.53	94.21	104.00
2	B	89	C	O3'-P-O5'	-6.32	94.52	104.00
1	A	845	ASP	CA-CB-CG	6.06	118.66	112.60
2	B	99	G	O3'-P-O5'	-5.96	95.07	104.00
1	A	819	PHE	CA-CB-CG	5.93	119.73	113.80
3	C	-8	DT	O3'-P-O5'	-5.80	95.30	104.00
1	A	349	PHE	CA-CB-CG	5.45	119.25	113.80
2	B	64	C	O3'-P-O5'	-5.38	95.94	104.00
2	B	34	G	C1'-C2'-O2'	5.36	116.44	108.40
2	B	83	C	C2'-C3'-O3'	-5.28	105.78	113.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	66	G	C1'-C2'-O2'	5.25	116.28	108.40
1	A	151	ASP	CA-CB-CG	5.06	117.66	112.60
3	C	7	DC	O3'-P-O5'	-5.02	96.47	104.00

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	51	ARG	Sidechain
1	A	804	ARG	Sidechain
1	A	901	ARG	Sidechain
1	A	91	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7172	0	7108	17	1
2	B	2258	0	1140	1	0
3	C	567	0	317	0	0
4	D	158	0	93	1	0
5	A	68	0	102	0	0
5	B	12	0	18	0	0
5	C	4	0	6	0	0
6	A	45	0	0	0	0
7	A	1	0	0	0	0
8	A	171	0	0	0	0
8	B	105	0	0	0	0
8	C	40	0	0	0	0
8	D	9	0	0	0	0
All	All	10610	0	8784	19	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:719:ASP:OD2	1:A:804:ARG:NH2	2.30	0.63
1:A:79:LEU:HG	1:A:210:LEU:HD13	1.82	0.61
1:A:72:HIS:CD2	1:A:83:ILE:HD11	2.40	0.56
1:A:23:ARG:O	1:A:23:ARG:HD2	2.10	0.52
1:A:706:ARG:HD2	1:A:724:MET:HE3	1.95	0.49
1:A:292:MET:HE3	1:A:299:VAL:HG13	1.94	0.48
1:A:79:LEU:HD11	1:A:210:LEU:HB3	1.96	0.48
1:A:367:THR:HG23	1:A:399:ASN:HB2	1.96	0.48
1:A:987:ARG:NH2	1:A:1049[B]:CYS:SG	2.88	0.46
1:A:292:MET:HE3	1:A:299:VAL:CG1	2.46	0.46
1:A:440:MET:HE2	1:A:444:LYS:HB3	1.98	0.45
1:A:491:ILE:HD12	1:A:497:PRO:HD3	1.98	0.45
1:A:112:GLN:HE21	1:A:112:GLN:H	1.66	0.42
4:D:1:DT:H2'	4:D:2:DT:C6	2.55	0.42
1:A:131:ASN:ND2	1:A:229:VAL:HG11	2.35	0.42
2:B:83:C:O3'	2:B:84:U:H4'	2.20	0.41
1:A:102:ARG:HH11	1:A:121:HIS:HD1	1.68	0.41
1:A:348:ILE:HD12	1:A:416:LEU:HD11	2.01	0.41
1:A:706:ARG:HD3	1:A:725:ASP:OD1	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:GLU:OE1	1:A:156:GLU:OE1[2_656]	2.13	0.07

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	886/933 (95%)	869 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	772/836 (92%)	760 (98%)	12 (2%)	55 62

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ARG
1	A	112	GLN
1	A	180	ASN
1	A	236	GLU
1	A	354	LEU
1	A	409	LEU
1	A	463	VAL
1	A	749	ARG
1	A	766	ARG
1	A	827	LYS
1	A	836	LYS
1	A	922	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	112	GLN
1	A	182	ASN
1	A	265	GLN
1	A	276	GLN
1	A	288	GLN
1	A	302	HIS
1	A	346	ASN
1	A	350	ASN
1	A	477	ASN
1	A	696	ASN
1	A	742	ASN
1	A	816	GLN
1	A	905	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	953	ASN
1	A	1046	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	104/110 (94%)	10 (9%)	1 (0%)

All (10) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	31	C
2	B	51	A
2	B	63	C
2	B	76	G
2	B	84	U
2	B	85	A
2	B	87	C
2	B	92	U
2	B	93	U
2	B	94	U

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	B	84	U

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 31 ligands modelled in this entry, 1 is monoatomic - leaving 30 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	SO4	A	1119	-	4,4,4	0.32	0	6,6,6	0.09	0
6	SO4	A	1126	-	4,4,4	0.34	0	6,6,6	0.15	0
5	EDO	A	1108	-	3,3,3	0.26	0	2,2,2	0.14	0
5	EDO	A	1101	-	3,3,3	0.21	0	2,2,2	0.08	0
5	EDO	A	1110	-	3,3,3	0.26	0	2,2,2	0.57	0
5	EDO	A	1114	-	3,3,3	0.14	0	2,2,2	0.36	0
5	EDO	A	1116	-	3,3,3	0.18	0	2,2,2	0.29	0
5	EDO	A	1109	-	3,3,3	0.44	0	2,2,2	0.62	0
6	SO4	A	1120	-	4,4,4	0.27	0	6,6,6	0.11	0
5	EDO	A	1105	-	3,3,3	0.14	0	2,2,2	0.24	0
5	EDO	B	203	-	3,3,3	0.09	0	2,2,2	0.28	0
5	EDO	A	1111	-	3,3,3	0.03	0	2,2,2	0.08	0
6	SO4	A	1125	-	4,4,4	0.25	0	6,6,6	0.22	0
5	EDO	A	1113	-	3,3,3	0.08	0	2,2,2	0.18	0
5	EDO	B	202	-	3,3,3	0.14	0	2,2,2	0.37	0
5	EDO	A	1107	-	3,3,3	0.13	0	2,2,2	0.13	0
5	EDO	B	201	-	3,3,3	0.03	0	2,2,2	0.18	0
5	EDO	A	1117	-	3,3,3	0.10	0	2,2,2	0.17	0
6	SO4	A	1118	-	4,4,4	0.31	0	6,6,6	0.14	0
6	SO4	A	1121	-	4,4,4	0.25	0	6,6,6	0.10	0
5	EDO	A	1106	-	3,3,3	0.17	0	2,2,2	0.27	0
6	SO4	A	1124	-	4,4,4	0.31	0	6,6,6	0.08	0
5	EDO	C	201	-	3,3,3	0.12	0	2,2,2	0.42	0
6	SO4	A	1123	-	4,4,4	0.31	0	6,6,6	0.08	0
6	SO4	A	1122	-	4,4,4	0.29	0	6,6,6	0.17	0
5	EDO	A	1112	-	3,3,3	0.22	0	2,2,2	0.48	0
5	EDO	A	1103	-	3,3,3	0.17	0	2,2,2	0.39	0
5	EDO	A	1104	-	3,3,3	0.36	0	2,2,2	0.35	0
5	EDO	A	1115	-	3,3,3	0.11	0	2,2,2	0.24	0
5	EDO	A	1102	-	3,3,3	0.25	0	2,2,2	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '–' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	1108	-	-	1/1/1/1	-
5	EDO	A	1101	-	-	0/1/1/1	-
5	EDO	A	1110	-	-	0/1/1/1	-
5	EDO	A	1114	-	-	0/1/1/1	-
5	EDO	A	1116	-	-	1/1/1/1	-
5	EDO	A	1109	-	-	1/1/1/1	-
5	EDO	A	1105	-	-	0/1/1/1	-
5	EDO	B	203	-	-	1/1/1/1	-
5	EDO	A	1111	-	-	0/1/1/1	-
5	EDO	A	1113	-	-	1/1/1/1	-
5	EDO	B	202	-	-	0/1/1/1	-
5	EDO	A	1107	-	-	1/1/1/1	-
5	EDO	B	201	-	-	1/1/1/1	-
5	EDO	A	1117	-	-	1/1/1/1	-
5	EDO	A	1106	-	-	1/1/1/1	-
5	EDO	C	201	-	-	0/1/1/1	-
5	EDO	A	1112	-	-	1/1/1/1	-
5	EDO	A	1103	-	-	1/1/1/1	-
5	EDO	A	1104	-	-	0/1/1/1	-
5	EDO	A	1115	-	-	0/1/1/1	-
5	EDO	A	1102	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1107	EDO	O1-C1-C2-O2
5	A	1103	EDO	O1-C1-C2-O2
5	A	1108	EDO	O1-C1-C2-O2
5	A	1116	EDO	O1-C1-C2-O2
5	A	1113	EDO	O1-C1-C2-O2
5	B	201	EDO	O1-C1-C2-O2
5	A	1117	EDO	O1-C1-C2-O2
5	A	1109	EDO	O1-C1-C2-O2
5	A	1106	EDO	O1-C1-C2-O2
5	A	1112	EDO	O1-C1-C2-O2
5	B	203	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	892/933 (95%)	0.46	43 (4%) 35 34	29, 55, 92, 135	5 (0%)
2	B	106/110 (96%)	-0.57	1 (0%) 81 80	38, 55, 131, 147	0
3	C	28/28 (100%)	-0.71	0 100 100	39, 46, 60, 62	0
4	D	8/8 (100%)	-0.49	0 100 100	42, 48, 61, 71	0
All	All	1034/1079 (95%)	0.32	44 (4%) 40 39	29, 55, 96, 147	5 (0%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	79	LEU	6.0
1	A	78	GLY	4.6
1	A	469	ASP	4.5
1	A	1005	ILE	4.3
1	A	90	LEU	4.0
1	A	468	ILE	3.5
1	A	80	ALA	3.5
1	A	718[A]	ASN	3.5
1	A	81	VAL	3.4
1	A	83	ILE	3.4
1	A	86	LEU	3.1
1	A	1	MET	3.1
1	A	1074	THR	2.9
1	A	456	ILE	2.8
1	A	457	SER	2.8
1	A	134	THR	2.7
1	A	88	GLN	2.6
1	A	91	ARG	2.6
1	A	740	VAL	2.5
1	A	463	VAL	2.5
1	A	76	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	45	ALA	2.5
1	A	320	ASP	2.5
1	A	464	VAL	2.4
1	A	307	ILE	2.4
1	A	737	ILE	2.4
1	A	696	ASN	2.4
2	B	84	U	2.3
1	A	44	GLU	2.3
1	A	2	ALA	2.3
1	A	760	LEU	2.2
1	A	827	LYS	2.2
1	A	1003	HIS	2.2
1	A	392	SER	2.2
1	A	85	HIS	2.2
1	A	131	ASN	2.2
1	A	794	LEU	2.1
1	A	77	ASN	2.1
1	A	84	GLN	2.1
1	A	870	LEU	2.1
1	A	923	SER	2.1
1	A	1076	ASP	2.1
1	A	317	VAL	2.0
1	A	873	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	SO4	A	1121	5/5	0.68	0.10	88,89,111,119	0
6	SO4	A	1123	5/5	0.72	0.08	115,119,122,125	0
5	EDO	A	1115	4/4	0.79	0.28	81,84,88,93	0
6	SO4	A	1125	5/5	0.79	0.13	68,69,90,96	0
6	SO4	A	1120	5/5	0.82	0.10	86,86,93,106	0
5	EDO	A	1105	4/4	0.84	0.17	85,88,93,96	0
5	EDO	A	1117	4/4	0.84	0.19	55,68,73,79	0
5	EDO	A	1107	4/4	0.84	0.14	65,73,74,77	0
5	EDO	A	1116	4/4	0.86	0.12	70,79,81,84	0
5	EDO	C	201	4/4	0.87	0.14	68,69,70,81	0
6	SO4	A	1119	5/5	0.88	0.06	78,96,102,108	0
5	EDO	A	1111	4/4	0.89	0.15	77,77,81,85	0
6	SO4	A	1122	5/5	0.89	0.09	77,81,99,107	0
6	SO4	A	1126	5/5	0.89	0.09	62,73,82,82	0
6	SO4	A	1124	5/5	0.90	0.07	112,115,120,126	0
5	EDO	A	1101	4/4	0.90	0.15	58,60,61,63	0
5	EDO	B	202	4/4	0.90	0.21	56,67,74,79	0
5	EDO	A	1112	4/4	0.91	0.11	52,62,64,72	0
5	EDO	A	1104	4/4	0.92	0.13	60,63,63,63	0
5	EDO	A	1109	4/4	0.93	0.12	53,59,64,66	0
7	CL	A	1127	1/1	0.93	0.11	68,68,68,68	0
5	EDO	A	1108	4/4	0.94	0.11	48,53,58,58	0
5	EDO	B	203	4/4	0.94	0.12	73,73,75,80	0
5	EDO	A	1103	4/4	0.95	0.12	56,58,58,64	0
5	EDO	A	1114	4/4	0.95	0.11	66,68,69,69	0
5	EDO	A	1106	4/4	0.95	0.09	53,53,56,59	0
5	EDO	A	1110	4/4	0.95	0.11	41,60,62,66	0
5	EDO	A	1102	4/4	0.95	0.10	53,54,55,55	0
5	EDO	B	201	4/4	0.95	0.11	58,60,61,63	0
5	EDO	A	1113	4/4	0.96	0.10	71,77,77,78	0
6	SO4	A	1118	5/5	0.98	0.06	44,46,48,51	0

6.5 Other polymers ⓘ

There are no such residues in this entry.