



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 12:01 PM UTC

PDB ID : 4X6G / pdb_00004x6g
Title : Full-length OxyR C199D from pseudomonas aeruginosa
Authors : Jo, I.; Ha, N.C.
Deposited on : 2014-12-08
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	3% 79% 15% 6%
1	B	316	3% 81% 12% 6%
1	C	316	2% 82% 11% 7%
1	D	316	2% 77% 16% . .
1	E	316	4% 81% 15% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	316	3% 77% 18% 5%
1	G	316	13% 71% 16% • 12%
1	H	316	3% 76% 15% • 8%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 19805 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called OxyR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	S	0	1	0
			2338	1495	411	425	7			
1	B	296	Total	C	N	O	S	0	1	0
			2326	1490	412	417	7			
1	C	295	Total	C	N	O	S	0	0	0
			2307	1477	404	420	6			
1	D	302	Total	C	N	O	S	0	0	0
			2362	1511	415	429	7			
1	E	307	Total	C	N	O	S	0	1	0
			2414	1542	425	440	7			
1	F	300	Total	C	N	O	S	0	1	0
			2356	1507	415	427	7			
1	G	279	Total	C	N	O	S	0	0	0
			2176	1398	376	396	6			
1	H	292	Total	C	N	O	S	0	0	0
			2278	1461	398	412	7			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP Q9HTL4
A	-4	HIS	-	expression tag	UNP Q9HTL4
A	-3	HIS	-	expression tag	UNP Q9HTL4
A	-2	HIS	-	expression tag	UNP Q9HTL4
A	-1	HIS	-	expression tag	UNP Q9HTL4
A	0	HIS	-	expression tag	UNP Q9HTL4
A	199	ASP	CYS	engineered mutation	UNP Q9HTL4
B	-5	HIS	-	expression tag	UNP Q9HTL4
B	-4	HIS	-	expression tag	UNP Q9HTL4
B	-3	HIS	-	expression tag	UNP Q9HTL4
B	-2	HIS	-	expression tag	UNP Q9HTL4
B	-1	HIS	-	expression tag	UNP Q9HTL4
B	0	HIS	-	expression tag	UNP Q9HTL4

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	199	ASP	CYS	engineered mutation	UNP Q9HTL4
C	-5	HIS	-	expression tag	UNP Q9HTL4
C	-4	HIS	-	expression tag	UNP Q9HTL4
C	-3	HIS	-	expression tag	UNP Q9HTL4
C	-2	HIS	-	expression tag	UNP Q9HTL4
C	-1	HIS	-	expression tag	UNP Q9HTL4
C	0	HIS	-	expression tag	UNP Q9HTL4
C	199	ASP	CYS	engineered mutation	UNP Q9HTL4
D	-5	HIS	-	expression tag	UNP Q9HTL4
D	-4	HIS	-	expression tag	UNP Q9HTL4
D	-3	HIS	-	expression tag	UNP Q9HTL4
D	-2	HIS	-	expression tag	UNP Q9HTL4
D	-1	HIS	-	expression tag	UNP Q9HTL4
D	0	HIS	-	expression tag	UNP Q9HTL4
D	199	ASP	CYS	engineered mutation	UNP Q9HTL4
E	-5	HIS	-	expression tag	UNP Q9HTL4
E	-4	HIS	-	expression tag	UNP Q9HTL4
E	-3	HIS	-	expression tag	UNP Q9HTL4
E	-2	HIS	-	expression tag	UNP Q9HTL4
E	-1	HIS	-	expression tag	UNP Q9HTL4
E	0	HIS	-	expression tag	UNP Q9HTL4
E	199	ASP	CYS	engineered mutation	UNP Q9HTL4
F	-5	HIS	-	expression tag	UNP Q9HTL4
F	-4	HIS	-	expression tag	UNP Q9HTL4
F	-3	HIS	-	expression tag	UNP Q9HTL4
F	-2	HIS	-	expression tag	UNP Q9HTL4
F	-1	HIS	-	expression tag	UNP Q9HTL4
F	0	HIS	-	expression tag	UNP Q9HTL4
F	199	ASP	CYS	engineered mutation	UNP Q9HTL4
G	-5	HIS	-	expression tag	UNP Q9HTL4
G	-4	HIS	-	expression tag	UNP Q9HTL4
G	-3	HIS	-	expression tag	UNP Q9HTL4
G	-2	HIS	-	expression tag	UNP Q9HTL4
G	-1	HIS	-	expression tag	UNP Q9HTL4
G	0	HIS	-	expression tag	UNP Q9HTL4
G	199	ASP	CYS	engineered mutation	UNP Q9HTL4
H	-5	HIS	-	expression tag	UNP Q9HTL4
H	-4	HIS	-	expression tag	UNP Q9HTL4
H	-3	HIS	-	expression tag	UNP Q9HTL4
H	-2	HIS	-	expression tag	UNP Q9HTL4
H	-1	HIS	-	expression tag	UNP Q9HTL4
H	0	HIS	-	expression tag	UNP Q9HTL4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	199	ASP	CYS	engineered mutation	UNP Q9HTL4

- Molecule 2 is HYDROGEN PEROXIDE (CCD ID: PEO) (formula: H_2O_2).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O 2 2	0	0
2	B	1	Total O 2 2	0	0
2	C	1	Total O 2 2	0	0
2	D	1	Total O 2 2	0	0
2	E	1	Total O 2 2	0	0
2	F	1	Total O 2 2	0	0
2	G	1	Total O 2 2	0	0
2	H	1	Total O 2 2	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	F	1	Total	C	O	0	0
			6	3	3		
3	G	1	Total	C	O	0	0
			6	3	3		
3	H	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	H	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	162	Total	O		0	0
			162	162			
4	B	123	Total	O		0	0
			123	123			
4	C	149	Total	O		0	0
			149	149			
4	D	166	Total	O		0	0
			166	166			
4	E	164	Total	O		0	0
			164	164			
4	F	111	Total	O		0	0
			111	111			
4	G	138	Total	O		0	0
			138	138			
4	H	129	Total	O		0	0
			129	129			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: OxyR




- Molecule 1: OxyR




- Molecule 1: OxyR




- Molecule 1: OxyR





Chain H:  3% 76% 15% 8%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.64Å 151.13Å 141.58Å 90.00° 97.28° 90.00°	Depositor
Resolution (Å)	19.96 – 2.00 19.96 – 2.00	Depositor EDS
% Data completeness (in resolution range)	75.3 (19.96-2.00) 75.3 (19.96-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.46 (at 2.00Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.186 , 0.237 0.189 , 0.238	Depositor DCC
R_{free} test set	8656 reflections (3.78%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.173	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 43.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	19805	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.54 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2852e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.78	0/2392	1.03	6/3254 (0.2%)
1	B	0.76	0/2383	0.98	2/3245 (0.1%)
1	C	0.76	1/2361 (0.0%)	0.98	0/3213
1	D	0.76	0/2419	0.97	4/3295 (0.1%)
1	E	0.80	0/2470	1.01	7/3361 (0.2%)
1	F	0.77	1/2412 (0.0%)	0.96	1/3284 (0.0%)
1	G	0.80	0/2224	1.03	4/3026 (0.1%)
1	H	0.74	0/2331	1.03	8/3173 (0.3%)
All	All	0.77	2/18992 (0.0%)	1.00	32/25851 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	E	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	300	ARG	CA-C	5.58	1.58	1.52
1	C	110	ILE	CA-CB	-5.45	1.50	1.53

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	36	VAL	CA-C-N	-11.82	103.94	122.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	36	VAL	C-N-CA	-11.82	103.94	122.49
1	B	120	MET	CA-C-N	-7.26	113.33	120.52
1	B	120	MET	C-N-CA	-7.26	113.33	120.52
1	E	117	ALA	CA-C-N	7.24	127.87	119.47

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	37	LYS	Peptide
1	E	309	ILE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2338	0	2376	32	1
1	B	2326	0	2357	25	0
1	C	2307	0	2343	21	0
1	D	2362	0	2388	42	0
1	E	2414	0	2443	40	1
1	F	2356	0	2386	41	0
1	G	2176	0	2218	42	0
1	H	2278	0	2316	34	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
2	G	2	0	0	0	0
2	H	2	0	0	0	0
3	A	18	0	24	3	0
3	B	18	0	24	4	0
3	C	12	0	16	0	0
3	D	12	0	16	1	0
3	E	6	0	8	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	6	0	8	2	0
3	G	6	0	8	3	0
3	H	12	0	16	2	0
4	A	162	0	0	5	0
4	B	123	0	0	2	0
4	C	149	0	0	5	0
4	D	166	0	0	3	0
4	E	164	0	0	4	0
4	F	111	0	0	2	0
4	G	138	0	0	2	0
4	H	129	0	0	1	0
All	All	19805	0	18947	260	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 260 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:LYS:HB2	1:A:40:GLU:HB2	1.39	1.04
1:H:187:ASN:HB3	1:H:209:PRO:HG2	1.49	0.94
1:A:137:ARG:NH2	1:A:152:GLU:OE1	2.06	0.88
1:E:120:MET:HE1	1:E:285:ALA:HB1	1.56	0.86
1:D:187:ASN:ND2	1:D:209:PRO:O	2.10	0.85

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:ARG:NH1	1:E:310:ALA:OXT[1_554]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/316 (94%)	291 (98%)	6 (2%)	0	100	100
1	B	293/316 (93%)	284 (97%)	9 (3%)	0	100	100
1	C	293/316 (93%)	285 (97%)	8 (3%)	0	100	100
1	D	298/316 (94%)	289 (97%)	8 (3%)	1 (0%)	36	35
1	E	304/316 (96%)	296 (97%)	8 (3%)	0	100	100
1	F	297/316 (94%)	287 (97%)	10 (3%)	0	100	100
1	G	273/316 (86%)	266 (97%)	7 (3%)	0	100	100
1	H	288/316 (91%)	273 (95%)	13 (4%)	2 (1%)	18	14
All	All	2343/2528 (93%)	2271 (97%)	69 (3%)	3 (0%)	48	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	82	ALA
1	H	82	ALA
1	H	86	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/269 (94%)	248 (98%)	6 (2%)	43	47
1	B	252/269 (94%)	249 (99%)	3 (1%)	63	70
1	C	250/269 (93%)	245 (98%)	5 (2%)	48	54
1	D	257/269 (96%)	248 (96%)	9 (4%)	32	32
1	E	262/269 (97%)	256 (98%)	6 (2%)	44	49
1	F	256/269 (95%)	249 (97%)	7 (3%)	39	42
1	G	236/269 (88%)	225 (95%)	11 (5%)	23	22
1	H	247/269 (92%)	242 (98%)	5 (2%)	48	54

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	2014/2152 (94%)	1962 (97%)	52 (3%)	40 44

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	296	CYS
1	F	240	VAL
1	H	222	VAL
1	F	50	ARG
1	F	86	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	88	GLN
1	G	114	HIS
1	H	76	GLN
1	D	114	HIS
1	D	0	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

23 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PEO	B	401	-	1,1,1	0.17	0	-		
3	GOL	B	403	-	5,5,5	0.44	0	5,5,5	0.73	0
3	GOL	H	402	-	5,5,5	0.40	0	5,5,5	0.54	0
2	PEO	F	401	-	1,1,1	0.15	0	-		
2	PEO	H	401	-	1,1,1	0.18	0	-		
2	PEO	E	401	-	1,1,1	0.11	0	-		
3	GOL	C	403	-	5,5,5	0.31	0	5,5,5	0.51	0
3	GOL	H	403	-	5,5,5	0.38	0	5,5,5	0.48	0
3	GOL	A	403	-	5,5,5	0.31	0	5,5,5	1.03	0
2	PEO	C	401	-	1,1,1	0.22	0	-		
3	GOL	C	402	-	5,5,5	0.70	0	5,5,5	0.85	0
2	PEO	D	401	-	1,1,1	0.00	0	-		
3	GOL	B	404	-	5,5,5	0.43	0	5,5,5	0.54	0
3	GOL	D	403	-	5,5,5	0.41	0	5,5,5	0.51	0
3	GOL	B	402	-	5,5,5	0.40	0	5,5,5	0.81	0
3	GOL	A	404	-	5,5,5	0.48	0	5,5,5	0.41	0
3	GOL	E	402	-	5,5,5	0.29	0	5,5,5	0.77	0
3	GOL	F	402	-	5,5,5	0.45	0	5,5,5	1.21	1 (20%)
2	PEO	G	401	-	1,1,1	0.18	0	-		
3	GOL	G	402	-	5,5,5	0.34	0	5,5,5	0.76	0
2	PEO	A	401	-	1,1,1	0.20	0	-		
3	GOL	A	402	-	5,5,5	0.36	0	5,5,5	0.93	0
3	GOL	D	402	-	5,5,5	0.40	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	F	402	-	-	3/4/4/4	-
3	GOL	B	403	-	-	2/4/4/4	-
3	GOL	C	403	-	-	4/4/4/4	-
3	GOL	H	402	-	-	0/4/4/4	-
3	GOL	A	403	-	-	2/4/4/4	-
3	GOL	H	403	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	402	-	-	2/4/4/4	-
3	GOL	G	402	-	-	0/4/4/4	-
3	GOL	B	404	-	-	2/4/4/4	-
3	GOL	D	403	-	-	2/4/4/4	-
3	GOL	B	402	-	-	2/4/4/4	-
3	GOL	A	402	-	-	4/4/4/4	-
3	GOL	D	402	-	-	2/4/4/4	-
3	GOL	A	404	-	-	4/4/4/4	-
3	GOL	E	402	-	-	4/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	402	GOL	O1-C1-C2	2.22	120.36	110.38

There are no chirality outliers.

5 of 35 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	GOL	O1-C1-C2-C3
3	A	402	GOL	C1-C2-C3-O3
3	A	404	GOL	O1-C1-C2-C3
3	A	404	GOL	C1-C2-C3-O3
3	A	404	GOL	O2-C2-C3-O3

There are no ring outliers.

10 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	H	402	GOL	1	0
3	H	403	GOL	1	0
3	B	404	GOL	3	0
3	B	402	GOL	1	0
3	A	404	GOL	1	0
3	E	402	GOL	2	0
3	F	402	GOL	2	0
3	G	402	GOL	3	0
3	A	402	GOL	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	402	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/316 (94%)	-0.32	11 (3%) 45 44	11, 28, 75, 102	1 (0%)
1	B	296/316 (93%)	-0.21	9 (3%) 52 51	13, 31, 66, 85	1 (0%)
1	C	295/316 (93%)	-0.42	6 (2%) 65 64	11, 28, 66, 103	0
1	D	302/316 (95%)	-0.37	7 (2%) 61 60	12, 27, 63, 87	0
1	E	307/316 (97%)	-0.19	12 (3%) 43 42	13, 30, 76, 92	1 (0%)
1	F	300/316 (94%)	-0.10	11 (3%) 45 44	12, 34, 74, 96	1 (0%)
1	G	279/316 (88%)	0.10	40 (14%) 6 5	10, 30, 100, 110	0
1	H	292/316 (92%)	-0.30	8 (2%) 56 55	14, 30, 70, 88	0
All	All	2369/2528 (93%)	-0.23	104 (4%) 39 38	10, 30, 75, 110	4 (0%)

The worst 5 of 104 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	48	PHE	5.4
1	G	8	TYR	4.4
1	D	120	MET	4.3
1	H	209	PRO	4.1
1	G	60	VAL	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	404	6/6	0.77	0.16	51,60,64,65	0
3	GOL	C	403	6/6	0.78	0.12	54,58,65,68	0
3	GOL	A	403	6/6	0.81	0.13	44,49,51,59	0
3	GOL	G	402	6/6	0.83	0.14	39,41,46,55	0
3	GOL	E	402	6/6	0.84	0.14	40,51,57,59	0
3	GOL	B	404	6/6	0.86	0.13	30,59,60,64	0
3	GOL	C	402	6/6	0.87	0.14	39,42,45,48	0
3	GOL	D	402	6/6	0.88	0.11	28,38,41,43	0
3	GOL	B	403	6/6	0.88	0.11	37,39,41,44	0
3	GOL	B	402	6/6	0.88	0.12	39,44,60,64	0
3	GOL	H	402	6/6	0.88	0.09	33,37,50,52	0
3	GOL	A	402	6/6	0.89	0.09	32,45,46,47	0
3	GOL	F	402	6/6	0.90	0.09	30,45,47,49	0
3	GOL	D	403	6/6	0.92	0.14	41,51,53,54	0
3	GOL	H	403	6/6	0.92	0.11	29,49,55,57	0
2	PEO	D	401	2/2	0.97	0.07	15,15,15,22	0
2	PEO	B	401	2/2	0.98	0.06	14,14,14,26	0
2	PEO	C	401	2/2	0.98	0.10	11,11,11,33	0
2	PEO	A	401	2/2	0.98	0.05	15,15,15,24	0
2	PEO	F	401	2/2	0.98	0.04	20,20,20,20	0
2	PEO	H	401	2/2	0.98	0.06	16,16,16,24	0
2	PEO	E	401	2/2	0.99	0.05	8,8,8,26	0
2	PEO	G	401	2/2	0.99	0.05	13,13,13,24	0

6.5 Other polymers

There are no such residues in this entry.