



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 29, 2026 – 05:39 PM UTC

PDB ID : 1X79 / pdb_00001x79
Title : Crystal structure of human GGA1 GAT domain complexed with the GAT-binding domain of Rabaptin5
Authors : Zhu, G.; Zhang, X.C.
Deposited on : 2004-08-13
Resolution : 2.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6562 (2.44-2.40)
Ramachandran outliers	187476	6481 (2.44-2.40)
Sidechain outliers	187428	6482 (2.44-2.40)
RSRZ outliers	180081	6066 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	98	
2	B	112	
2	C	112	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	DTT	B	395	X	X	-	-
4	DTT	C	394	X	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2243 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribosylation factor binding protein GGA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	82	Total	C	N	O	S	0	0	0
			664	407	118	134	5			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	205	GLY	-	cloning artifact	UNP Q9UJY5
A	206	PRO	-	cloning artifact	UNP Q9UJY5
A	207	LEU	-	cloning artifact	UNP Q9UJY5
A	208	GLY	-	cloning artifact	UNP Q9UJY5
A	209	SER	-	cloning artifact	UNP Q9UJY5

- Molecule 2 is a protein called Rab GTPase binding effector protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	90	Total	C	N	O	S	0	0	0
			727	440	130	153	4			
2	C	88	Total	C	N	O	S	0	0	0
			709	430	128	147	4			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	550	MET	-	initiating methionine	UNP Q15276
C	550	MET	-	initiating methionine	UNP Q15276

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (CCD ID: DTT) (formula: $C_4H_{10}O_2S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	O	S	0	0
			8	4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	O	S	0	0
			8	4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	28	Total	O	0	0
			28	28		
5	B	42	Total	O	0	0
			42	42		
5	C	47	Total	O	0	0
			47	47		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-ribosylation factor binding protein GGA1



- Molecule 2: Rab GTPase binding effector protein 1



- Molecule 2: Rab GTPase binding effector protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	155.22Å 155.22Å 53.05Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.81 – 2.41 50.81 – 2.41	Depositor EDS
% Data completeness (in resolution range)	82.5 (50.81-2.41) 82.8 (50.81-2.41)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.86 (at 2.39Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.233 , 0.283 0.225 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	41.9	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 63.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.055 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2243	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DTT, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/667	0.96	1/896 (0.1%)
2	B	0.83	0/728	0.90	2/970 (0.2%)
2	C	0.90	1/710 (0.1%)	0.93	1/946 (0.1%)
All	All	0.79	1/2105 (0.0%)	0.93	4/2812 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	575	MET	SD-CE	6.20	1.95	1.79

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	627	GLN	N-CA-C	-6.81	103.86	111.28
2	B	636	GLU	N-CA-C	-6.51	104.22	111.71
2	C	588	SER	N-CA-C	-6.12	104.69	111.36
1	A	280	ILE	CB-CA-C	-5.46	104.98	111.81

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	664	0	666	50	0
2	B	727	0	731	20	0
2	C	709	0	719	38	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
4	B	8	0	8	0	0
4	C	8	0	8	1	0
5	A	28	0	0	2	0
5	B	42	0	0	3	0
5	C	47	0	0	2	0
All	All	2243	0	2132	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 100 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270:THR:HG21	1:A:277:LEU:HB2	1.47	0.97
2:C:638:VAL:HG23	2:C:639:SER:H	1.40	0.85
2:B:574:THR:HG22	2:C:574:THR:HG22	1.64	0.80
2:B:628:MET:HG2	2:B:632:MET:HE3	1.65	0.78
1:A:211:ILE:O	1:A:215:VAL:HG23	1.84	0.78

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	78/98 (80%)	66 (85%)	8 (10%)	4 (5%)	1 1
2	B	88/112 (79%)	82 (93%)	6 (7%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	86/112 (77%)	76 (88%)	9 (10%)	1 (1%)	10	15
All	All	252/322 (78%)	224 (89%)	23 (9%)	5 (2%)	6	7

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	272	ASP
1	A	274	ASP
1	A	234	SER
2	C	639	SER
1	A	231	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/84 (92%)	70 (91%)	7 (9%)	9	13
2	B	83/103 (81%)	73 (88%)	10 (12%)	5	6
2	C	81/103 (79%)	72 (89%)	9 (11%)	6	8
All	All	241/290 (83%)	215 (89%)	26 (11%)	6	9

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	633	GLN
2	C	564	LEU
2	C	635	ARG
2	B	636	GLU
2	C	571	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	568	ASN
2	C	570	GLN
1	A	284	ASN
2	B	561	GLN
2	B	566	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	C	392	-	4,4,4	0.38	0	6,6,6	0.13	0
3	SO4	B	391	-	4,4,4	0.37	0	6,6,6	0.12	0
4	DTT	B	395	-	7,7,7	6.46	5 (71%)	4,8,8	5.36	4 (100%)
4	DTT	C	394	-	7,7,7	6.81	5 (71%)	4,8,8	5.84	4 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTT	B	395	-	2/2/2/2	3/8/8/8	-
4	DTT	C	394	-	2/2/2/2	4/8/8/8	-

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	394	DTT	O2-C2	-16.41	1.09	1.43
4	B	395	DTT	O2-C2	-15.30	1.11	1.43
4	C	394	DTT	O3-C3	-4.32	1.34	1.43
4	B	395	DTT	C3-C2	-4.24	1.42	1.53
4	B	395	DTT	C4-C3	-3.85	1.40	1.51

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	395	DTT	C3-C4-S4	-8.65	90.26	114.43
4	C	394	DTT	C3-C4-S4	-8.21	91.49	114.43
4	C	394	DTT	C2-C1-S1	-6.63	95.92	114.43
4	B	395	DTT	C2-C1-S1	-4.10	102.99	114.43
4	B	395	DTT	O2-C2-C3	4.07	118.13	109.57

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	395	DTT	C3
4	B	395	DTT	C2
4	C	394	DTT	C3
4	C	394	DTT	C2

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	395	DTT	C1-C2-C3-O3
4	C	394	DTT	S1-C1-C2-O2
4	C	394	DTT	S1-C1-C2-C3
4	C	394	DTT	C2-C3-C4-S4
4	C	394	DTT	O3-C3-C4-S4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	394	DTT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	82/98 (83%)	0.70	9 (10%) 10 8	35, 71, 109, 119	0
2	B	90/112 (80%)	0.04	5 (5%) 30 26	29, 46, 106, 113	0
2	C	88/112 (78%)	0.13	5 (5%) 29 25	28, 45, 104, 117	0
All	All	260/322 (80%)	0.28	19 (7%) 21 17	28, 52, 108, 119	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	553	THR	5.5
1	A	211	ILE	4.5
1	A	236	SER	3.8
2	C	638	VAL	3.3
1	A	248	LEU	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	C	392	5/5	0.93	0.09	87,90,92,93	0
4	DTT	C	394	8/8	0.93	0.11	57,66,83,86	0
4	DTT	B	395	8/8	0.94	0.15	74,83,88,97	0
3	SO4	B	391	5/5	0.95	0.08	79,81,81,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.