



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:46 AM UTC

PDB ID : 4X7R / pdb_00004x7r
Title : Crystal structure of S. aureus TarM G117R mutant in complex with Fondaparinux, alpha-GlcNAc-glycerol and UDP
Authors : Worrall, L.J.; Sobhanifar, S.; Strynadka, N.C.
Deposited on : 2014-12-09
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	493	4% 92% 8%
1	B	493	5% 92% 7%
2	C	5	80% 20%
2	D	5	60% 40%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TarM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	493	Total	C	N	O	S	0	0	0
			4040	2585	684	752	19			
1	B	493	Total	C	N	O	S	0	0	0
			4040	2585	684	752	19			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	117	ARG	GLY	engineered mutation	UNP H0AM96
B	117	ARG	GLY	engineered mutation	UNP H0AM96

- Molecule 2 is an oligosaccharide called 2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-4)-2-deoxy-3,6-di-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-methyl 2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranoside.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	5	Total	C	N	O	S	0	0	0
			91	31	3	49	8			
2	D	5	Total	C	N	O	S	0	0	0
			91	31	3	49	8			

- Molecule 3 is URIDINE-5'-DIPHOSPHATE (CCD ID: UDP) (formula: C₉H₁₄N₂O₁₂P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	B	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

- Molecule 4 is (2S)-2,3-dihydroxypropyl 2-acetamido-2-deoxy- α -D-glucopyranoside (CCD ID: 3YW) (formula: C₁₁H₂₁NO₈).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			20	11	1	8		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			20	11	1	8		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	183	Total	O	0	0
			183	183		
5	B	182	Total	O	0	0
			182	182		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TarM



- Molecule 1: TarM



- Molecule 2: 2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-4)-2-deoxy-3,6-di-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-methyl 2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranoside



- Molecule 2: 2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-4)-2-deoxy-3,6-di-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-methyl 2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranoside





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	43.47Å 92.14Å 96.11Å 65.89° 83.66° 84.12°	Depositor
Resolution (Å)	87.48 – 2.15 87.47 – 2.15	Depositor EDS
% Data completeness (in resolution range)	97.9 (87.48-2.15) 97.9 (87.47-2.15)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.14Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.179 , 0.200 (Not available) , 0.208	Depositor DCC
R_{free} test set	3707 reflections (4.06%)	wwPDB-VP
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.351	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.106 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8717	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SGN, UDP, ZDO, BDP, IDS, SUS, 3YW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	1/4117 (0.0%)	1.25	7/5525 (0.1%)
1	B	0.84	1/4117 (0.0%)	1.24	5/5525 (0.1%)
All	All	0.84	2/8234 (0.0%)	1.25	12/11050 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	449	MET	SD-CE	-7.32	1.61	1.79
1	B	449	MET	SD-CE	-7.18	1.61	1.79

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	326	ARG	N-CA-C	-6.19	100.67	109.96
1	A	326	ARG	N-CA-C	-6.14	100.75	109.96
1	B	491	PHE	CA-CB-CG	5.79	119.59	113.80
1	A	491	PHE	CA-CB-CG	5.63	119.43	113.80
1	A	14	ASN	CA-CB-CG	5.62	118.22	112.60
1	B	393	PHE	CA-CB-CG	5.49	119.29	113.80
1	A	393	PHE	CA-CB-CG	5.39	119.19	113.80
1	A	142	GLU	N-CA-C	5.20	116.97	109.07
1	A	17	GLY	N-CA-C	5.18	118.94	112.73
1	B	15	LYS	N-CA-C	5.17	118.11	110.46
1	B	7	MET	N-CA-C	5.07	117.51	109.24
1	A	121	ALA	N-CA-C	5.03	116.83	109.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4040	0	4067	19	0
1	B	4040	0	4067	19	0
2	C	91	0	27	1	0
2	D	91	0	27	2	0
3	A	25	0	11	1	0
3	B	25	0	11	0	0
4	A	20	0	6	2	0
4	B	20	0	6	0	0
5	A	183	0	0	1	0
5	B	182	0	0	2	0
All	All	8717	0	8222	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:339:VAL:HG21	1:B:449:MET:HE2	1.70	0.74
1:A:339:VAL:HG21	1:A:449:MET:HE2	1.70	0.73
1:A:343:VAL:HG12	1:A:350:VAL:HG11	1.72	0.71
1:B:343:VAL:HG12	1:B:350:VAL:HG11	1.72	0.69
1:B:306:VAL:CG2	1:B:414:ILE:HD11	2.29	0.63
1:A:7:MET:HE1	1:A:234:MET:HB2	1.84	0.60
1:A:306:VAL:CG2	1:A:414:ILE:HD11	2.31	0.60
1:B:7:MET:HE1	1:B:234:MET:HB2	1.85	0.59
1:B:395:LEU:HD22	1:B:456:LEU:HB3	1.89	0.54
1:A:395:LEU:HD22	1:A:456:LEU:HB3	1.89	0.53
1:A:229:GLY:HA3	2:C:3:SUS:O7S	2.11	0.51
1:B:395:LEU:HD21	1:B:457:VAL:HG13	1.95	0.49
1:A:395:LEU:HD21	1:A:457:VAL:HG13	1.94	0.48
1:A:399:THR:CG2	1:A:449:MET:HE1	2.44	0.47
1:B:399:THR:CG2	1:B:449:MET:HE1	2.46	0.46
1:A:327:MET:HE2	1:A:354:ILE:HG22	1.98	0.46
1:B:123:ILE:HG12	1:B:132:ILE:HG12	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:323:HIS:HE1	5:A:610:HOH:O	1.99	0.45
1:A:18:MET:HB2	4:A:503:3YW:C4A	2.47	0.45
1:B:327:MET:HE2	1:B:354:ILE:HG22	1.97	0.45
1:B:229:GLY:HA3	2:D:3:SUS:O7S	2.17	0.44
1:B:323:HIS:HE1	5:B:619:HOH:O	2.00	0.44
1:B:306:VAL:HG22	1:B:414:ILE:HD11	2.00	0.44
1:B:416:LYS:HE3	1:B:471:ARG:HD2	1.99	0.43
1:A:441:ILE:HG21	1:A:449:MET:HB2	2.01	0.43
1:B:77:SER:HB2	1:B:195:VAL:HG11	1.99	0.43
1:B:113:ASP:O	1:B:117:ARG:HA	2.19	0.43
1:A:77:SER:HB2	1:A:195:VAL:HG11	2.01	0.42
1:B:233:LYS:NZ	5:B:604:HOH:O	2.52	0.42
1:B:441:ILE:HG21	1:B:449:MET:HB2	2.01	0.42
1:A:156:THR:HB	1:A:165:TYR:HB3	2.00	0.42
1:A:306:VAL:HG22	1:A:414:ILE:HD11	1.99	0.42
1:A:391:GLU:HA	1:A:417:ARG:HH21	1.85	0.42
1:A:116:THR:HG23	1:A:118:THR:HG22	2.02	0.42
3:A:501:UDP:O1B	4:A:503:3YW:O1	2.39	0.41
1:B:203:ASN:HB3	2:D:2:IDS:O2S	2.20	0.41
1:B:339:VAL:HG11	1:B:449:MET:HE2	2.02	0.41
1:A:59:LYS:NZ	1:A:382:TYR:OH	2.53	0.41
1:A:339:VAL:HG11	1:A:449:MET:HE2	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	491/493 (100%)	473 (96%)	17 (4%)	1 (0%)	43	44
1	B	491/493 (100%)	473 (96%)	17 (4%)	1 (0%)	43	44
All	All	982/986 (100%)	946 (96%)	34 (4%)	2 (0%)	43	44

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	298	ASN
1	A	298	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/450 (76%)	339 (99%)	4 (1%)	63	70
1	B	450/450 (100%)	442 (98%)	8 (2%)	51	58
All	All	793/900 (88%)	781 (98%)	12 (2%)	57	64

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	TYR
1	A	131	VAL
1	A	264	LYS
1	A	395	LEU
1	B	47	TYR
1	B	118	THR
1	B	131	VAL
1	B	213	LYS
1	B	264	LYS
1	B	348	ASN
1	B	363	LYS
1	B	395	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	HIS
1	B	182	ASN
1	B	447	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ZDO	C	1	2	21,21,21	4.53	5 (23%)	26,32,32	1.29	4 (15%)
2	IDS	C	2	2	16,16,17	1.35	2 (12%)	16,24,26	1.21	1 (6%)
2	SUS	C	3	2	23,23,24	4.23	4 (17%)	26,36,38	1.31	1 (3%)
2	BDP	C	4	2	12,12,13	1.33	2 (16%)	14,17,19	0.66	0
2	SGN	C	5	2	19,19,20	4.48	6 (31%)	23,29,31	1.21	1 (4%)
2	ZDO	D	1	2	21,21,21	4.67	7 (33%)	26,32,32	1.44	2 (7%)
2	IDS	D	2	2	16,16,17	1.45	2 (12%)	16,24,26	1.24	2 (12%)
2	SUS	D	3	2	23,23,24	4.15	3 (13%)	26,36,38	1.39	3 (11%)
2	BDP	D	4	2	12,12,13	1.19	1 (8%)	14,17,19	0.76	0
2	SGN	D	5	2	19,19,20	3.98	4 (21%)	23,29,31	1.19	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ZDO	C	1	2	-	1/13/33/33	0/1/1/1
2	IDS	C	2	2	-	0/9/26/29	0/1/1/1
2	SUS	C	3	2	-	1/16/33/36	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BDP	C	4	2	-	0/4/21/24	0/1/1/1
2	SGN	C	5	2	-	1/11/28/31	0/1/1/1
2	ZDO	D	1	2	-	2/13/33/33	0/1/1/1
2	IDS	D	2	2	-	0/9/26/29	0/1/1/1
2	SUS	D	3	2	-	1/16/33/36	0/1/1/1
2	BDP	D	4	2	-	0/4/21/24	0/1/1/1
2	SGN	D	5	2	-	4/11/28/31	0/1/1/1

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	ZDO	S2-N2	14.41	1.77	1.59
2	D	3	SUS	S1-N2	14.07	1.77	1.59
2	C	3	SUS	S1-N2	13.73	1.76	1.59
2	C	1	ZDO	S2-N2	13.19	1.76	1.59
2	C	5	SGN	S1-N2	11.26	1.73	1.59
2	C	5	SGN	O2S-S1	10.91	1.54	1.42
2	C	1	ZDO	OSB-S2	10.74	1.54	1.42
2	D	5	SGN	O2S-S1	10.42	1.53	1.42
2	C	1	ZDO	OSA-S2	10.40	1.53	1.42
2	D	1	ZDO	OSA-S2	10.29	1.53	1.42
2	D	5	SGN	O1S-S1	10.17	1.53	1.42
2	C	5	SGN	O1S-S1	10.07	1.53	1.42
2	D	1	ZDO	OSB-S2	9.86	1.53	1.42
2	C	3	SUS	O3S-S1	9.80	1.53	1.42
2	C	3	SUS	O1S-S1	9.64	1.53	1.42
2	D	3	SUS	O3S-S1	9.26	1.52	1.42
2	D	3	SUS	O1S-S1	9.15	1.52	1.42
2	D	5	SGN	S1-N2	7.51	1.69	1.59
2	D	1	ZDO	O1-C1	3.88	1.46	1.40
2	C	1	ZDO	O1-C1	3.08	1.45	1.40
2	D	2	IDS	O5-C1	2.92	1.48	1.43
2	D	2	IDS	C1-C2	2.75	1.56	1.51
2	C	5	SGN	C1-C2	2.71	1.56	1.52
2	D	1	ZDO	C4-C5	2.59	1.58	1.53
2	D	5	SGN	C1-C2	2.53	1.55	1.52
2	C	4	BDP	C4-C5	2.43	1.57	1.53
2	D	4	BDP	C4-C5	2.41	1.57	1.53
2	C	4	BDP	C4-C3	2.32	1.58	1.52
2	C	5	SGN	O5-C5	2.20	1.47	1.43
2	C	2	IDS	C3-C2	2.19	1.57	1.53
2	C	2	IDS	C1-C2	2.11	1.55	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3	SUS	C4-C5	2.09	1.57	1.53
2	D	1	ZDO	OS2-S6	2.04	1.54	1.45
2	D	1	ZDO	O5-C1	2.02	1.47	1.41
2	C	5	SGN	O4S-S2	2.01	1.54	1.45
2	C	1	ZDO	OS1-S6	2.00	1.54	1.45

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	SUS	O1S-S1-O3S	-5.11	109.00	120.36
2	C	5	SGN	O1S-S1-O2S	-4.86	109.56	120.36
2	D	5	SGN	O1S-S1-O2S	-4.50	110.35	120.36
2	D	3	SUS	O1S-S1-O3S	-4.48	110.40	120.36
2	D	1	ZDO	OSA-S2-OSB	-4.25	110.91	120.36
2	C	2	IDS	O2-C2-C3	3.32	111.58	106.95
2	C	1	ZDO	OSA-S2-OSB	-3.25	113.13	120.36
2	D	2	IDS	O2-C2-C3	3.03	111.19	106.95
2	D	3	SUS	O5-C5-C6	-2.81	101.51	107.59
2	D	5	SGN	O6S-S2-O6	2.69	112.55	106.37
2	D	1	ZDO	O6-C6-C5	2.33	111.72	107.57
2	C	1	ZDO	OS3-S6-O6	2.24	111.52	106.37
2	D	2	IDS	C4-C3-C2	2.19	114.20	110.23
2	C	1	ZDO	O1-C1-C2	2.18	111.43	108.13
2	D	3	SUS	C1-O5-C5	2.13	115.05	112.19
2	C	1	ZDO	C4-C3-C2	2.11	113.47	110.40

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	1	ZDO	C2-N2-S2-OSA
2	D	5	SGN	C6-O6-S2-O6S
2	C	3	SUS	C2-N2-S1-O1S
2	D	3	SUS	C2-N2-S1-O1S
2	D	5	SGN	C2-N2-S1-O2S
2	D	5	SGN	C6-O6-S2-O5S
2	C	1	ZDO	C5-C6-O6-S6
2	D	5	SGN	C5-C6-O6-S2
2	C	5	SGN	C2-N2-S1-O1S
2	D	1	ZDO	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	3	SUS	1	0
2	D	2	IDS	1	0
2	C	3	SUS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	3YW	B	503	-	20,20,20	0.33	0	26,27,27	1.01	3 (11%)
3	UDP	A	501	-	25,26,26	0.60	1 (4%)	38,40,40	0.56	0
4	3YW	A	503	-	20,20,20	0.24	0	26,27,27	0.98	2 (7%)
3	UDP	B	501	-	25,26,26	0.89	1 (4%)	38,40,40	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	3YW	B	503	-	-	2/13/33/33	0/1/1/1
3	UDP	A	501	-	-	2/16/32/32	0/2/2/2
4	3YW	A	503	-	-	3/13/33/33	0/1/1/1
3	UDP	B	501	-	-	4/16/32/32	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	501	UDP	PA-O3A	-3.79	1.55	1.59
3	A	501	UDP	PA-O3A	-2.56	1.56	1.59

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	503	3YW	O5-C1-C2	2.65	115.60	110.59
4	A	503	3YW	O1-C4A-C5A	2.64	113.80	108.81
4	A	503	3YW	O5-C1-C2	2.51	115.34	110.59
4	B	503	3YW	C1-C2-C3	2.34	116.65	110.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	503	3YW	O1-C4A-C5A	2.18	112.94	108.81

There are no chirality outliers.

All (11) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	501	UDP	PB-O3A-PA-O5'
4	A	503	3YW	C2-C1-O1-C4A
4	B	503	3YW	C2-C1-O1-C4A
4	A	503	3YW	O1-C4A-C5A-C6A
3	B	501	UDP	PB-O3A-PA-O5'
4	A	503	3YW	O1-C4A-C5A-O2
3	B	501	UDP	C5'-O5'-PA-O1A
3	B	501	UDP	O4'-C4'-C5'-O5'
3	A	501	UDP	O4'-C4'-C5'-O5'
3	B	501	UDP	PA-O3A-PB-O1B
4	B	503	3YW	C5A-C4A-O1-C1



There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	UDP	1	0
4	A	503	3YW	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	493/493 (100%)	0.31	21 (4%)	40	45	31, 54, 93, 118	0
1	B	493/493 (100%)	0.36	25 (5%)	33	37	32, 54, 95, 132	0
All	All	986/986 (100%)	0.34	46 (4%)	36	41	31, 54, 95, 132	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	295	ASP	5.8
1	B	295	ASP	3.9
1	B	1	MET	3.5
1	B	493	SER	3.4
1	B	116	THR	3.2
1	A	116	THR	3.1
1	B	311	ALA	3.0
1	B	260	GLY	3.0
1	A	115	THR	2.9
1	B	368	ILE	2.8
1	A	1	MET	2.8
1	A	91	HIS	2.8
1	A	119	TYR	2.8
1	B	105	SER	2.7
1	B	256	PHE	2.7
1	A	139	LYS	2.7
1	A	106	LYS	2.6
1	B	313	LYS	2.6
1	B	351	LYS	2.6
1	A	306	VAL	2.6
1	A	14	ASN	2.5
1	B	350	VAL	2.5
1	A	313	LYS	2.5
1	B	365	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	363	LYS	2.5
1	B	259	THR	2.5
1	B	100	ILE	2.5
1	B	114	ILE	2.5
1	B	102	ILE	2.3
1	A	112	PHE	2.3
1	B	108	ILE	2.3
1	B	257	ASP	2.3
1	A	62	ARG	2.2
1	B	106	LYS	2.2
1	A	375	ARG	2.2
1	A	351	LYS	2.2
1	B	160	ASP	2.2
1	A	108	ILE	2.1
1	A	308	ILE	2.1
1	B	14	ASN	2.1
1	B	136	LYS	2.1
1	B	366	LYS	2.1
1	B	183	ALA	2.1
1	A	444	HIS	2.1
1	A	493	SER	2.1
1	A	107	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZDO	D	1	21/21	0.82	0.15	56,69,85,87	21
2	BDP	D	4	12/13	0.85	0.12	50,54,55,57	12
2	ZDO	C	1	21/21	0.86	0.15	56,62,87,89	21
2	BDP	C	4	12/13	0.87	0.11	50,54,56,59	12
2	SUS	D	3	23/24	0.94	0.10	40,46,60,61	23
2	SGN	D	5	19/20	0.94	0.09	45,53,63,65	19


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	IDS	C	2	16/17	0.95	0.08	44,48,51,51	16
2	SGN	C	5	19/20	0.95	0.09	41,50,64,66	19
2	SUS	C	3	23/24	0.95	0.09	44,47,61,64	23
2	IDS	D	2	16/17	0.96	0.07	40,44,49,49	16

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	3YW	B	503	20/20	0.88	0.16	37,52,63,67	20
4	3YW	A	503	20/20	0.92	0.13	36,51,55,57	20
3	UDP	A	501	25/25	0.96	0.07	45,50,55,57	0
3	UDP	B	501	25/25	0.96	0.07	46,50,56,57	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 3YW B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around 3YW A 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around UDP A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.