



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:59 PM UTC

PDB ID : 8X70 / pdb_00008x70
Title : The Crystal Structure of IFI16 from Biortus.
Authors : Wang, F.; Cheng, W.; Lv, Z.; Meng, Q.; Wang, J.
Deposited on : 2023-11-22
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	4% 87% 8% 5%
1	B	204	2% 86% 5% 9%
1	C	204	2% 93% • 5%
1	D	204	6% 88% 5% 6%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7045 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-interferon-inducible protein 16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	1	0
			1583	1009	272	293	9			
1	B	186	Total	C	N	O	S	0	0	0
			1505	962	254	280	9			
1	C	193	Total	C	N	O	S	0	1	0
			1573	1005	267	292	9			
1	D	191	Total	C	N	O	S	0	0	0
			1547	990	262	286	9			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	190	GLY	-	expression tag	UNP Q16666
A	191	GLY	-	expression tag	UNP Q16666
B	190	GLY	-	expression tag	UNP Q16666
B	191	GLY	-	expression tag	UNP Q16666
C	190	GLY	-	expression tag	UNP Q16666
C	191	GLY	-	expression tag	UNP Q16666
D	190	GLY	-	expression tag	UNP Q16666
D	191	GLY	-	expression tag	UNP Q16666

- Molecule 2 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Br	0	0
			1	1		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total	K	0	0
			6	6		
4	B	2	Total	K	0	0
			2	2		
4	C	4	Total	K	0	0
			4	4		
4	D	2	Total	K	0	0
			2	2		

- Molecule 5 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			10	6	4		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	227	Total	O	0	0
			227	227		
6	B	201	Total	O	0	0
			201	201		
6	C	217	Total	O	0	0
			217	217		
6	D	147	Total	O	0	0
			147	147		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gamma-interferon-inducible protein 16




- Molecule 1: Gamma-interferon-inducible protein 16



- Molecule 1: Gamma-interferon-inducible protein 16



- Molecule 1: Gamma-interferon-inducible protein 16



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.40Å 87.42Å 111.21Å 90.00° 99.27° 90.00°	Depositor
Resolution (Å)	46.52 – 1.70 46.52 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.52-1.70) 99.8 (46.52-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.201 , 0.247 0.207 , 0.253	Depositor DCC
R_{free} test set	4254 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 32.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.025 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7045	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BR, K, PGE, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	0/1612	1.11	0/2164
1	B	0.93	0/1530	1.11	0/2054
1	C	0.92	0/1601	1.11	0/2149
1	D	0.95	0/1572	1.17	0/2109
All	All	0.94	0/6315	1.13	0/8476

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1583	0	1634	10	0
1	B	1505	0	1546	10	0
1	C	1573	0	1626	4	0
1	D	1547	0	1600	6	0
2	A	1	0	0	0	0
3	A	8	0	12	0	0
3	C	8	0	12	0	0
3	D	4	0	6	2	0
4	A	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	2	0	0	0	0
4	C	4	0	0	0	0
4	D	2	0	0	0	0
5	C	10	0	14	2	0
6	A	227	0	0	1	0
6	B	201	0	0	2	0
6	C	217	0	0	1	0
6	D	147	0	0	0	0
All	All	7045	0	6450	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:227:ILE:HG22	1:A:247:THR:HG21	1.42	1.00
1:A:221:PRO:HD3	1:A:382:MET:HE3	1.74	0.70
1:B:268:ASP:HA	3:D:401:EDO:H21	1.76	0.67
1:D:234:ALA:HB3	1:D:290:VAL:HG21	1.79	0.63
1:C:310:GLN:O	5:C:402:PGE:H12	2.02	0.58
1:A:268:ASP:HB3	1:A:315:ILE:HG13	1.89	0.54
1:A:221:PRO:HD3	1:A:382:MET:CE	2.38	0.53
1:A:356:CYS:HB3	1:A:388:ILE:HD12	1.92	0.52
1:B:373:ASN:N	6:B:503:HOH:O	2.37	0.47
1:D:362:LEU:HD23	1:D:362:LEU:N	2.31	0.46
1:B:270:LEU:C	1:B:270:LEU:HD12	2.40	0.46
1:B:314:ASN:HA	3:D:401:EDO:H22	1.98	0.46
1:A:311:ALA:HA	5:C:402:PGE:H6	1.97	0.45
1:B:388:ILE:O	1:B:389:LYS:CB	2.64	0.45
1:D:362:LEU:HD22	1:D:388:ILE:CD1	2.47	0.45
1:B:243:LYS:HD3	1:B:366:CYS:SG	2.57	0.44
1:D:302:LEU:HD23	1:D:302:LEU:HA	1.86	0.44
1:B:353:ASN:O	1:B:353:ASN:CG	2.62	0.43
1:C:333:ILE:N	1:C:333:ILE:HD12	2.34	0.43
1:A:229:PHE:CZ	1:A:244:VAL:HG11	2.53	0.43
1:A:385:PHE:CE2	1:A:387:GLN:HG3	2.54	0.43
1:D:308:HIS:HE1	1:D:342:LYS:O	2.02	0.43
1:A:372:LYS:O	1:A:373:ASN:C	2.63	0.42
1:B:324:LYS:NZ	6:B:506:HOH:O	2.47	0.42
1:B:311:ALA:HB3	1:B:314:ASN:ND2	2.35	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:306:ILE:O	1:C:310:GLN:HG2	2.20	0.41
1:D:315:ILE:HD11	1:D:368:ARG:CZ	2.51	0.41
1:B:388:ILE:O	1:B:389:LYS:HB2	2.19	0.41
1:C:325:LYS:NZ	6:C:519:HOH:O	2.54	0.41
1:A:275[A]:GLU:HG3	6:A:674:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/204 (95%)	187 (97%)	6 (3%)	0	100	100
1	B	182/204 (89%)	181 (100%)	1 (0%)	0	100	100
1	C	190/204 (93%)	188 (99%)	2 (1%)	0	100	100
1	D	187/204 (92%)	185 (99%)	2 (1%)	0	100	100
All	All	752/816 (92%)	741 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/189 (96%)	181 (100%)	1 (0%)	81	76

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	173/189 (92%)	173 (100%)	0	100	100
1	C	182/189 (96%)	182 (100%)	0	100	100
1	D	178/189 (94%)	175 (98%)	3 (2%)	53	38
All	All	715/756 (95%)	711 (99%)	4 (1%)	78	72

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	223	MET
1	D	310	GLN
1	D	325	LYS
1	D	362	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	296	ASN
1	A	310	GLN
1	A	337	GLN
1	A	373	ASN
1	B	241	HIS
1	C	241	HIS
1	C	292	ASN
1	C	310	GLN
1	C	373	ASN
1	C	387	GLN
1	D	241	HIS
1	D	308	HIS
1	D	314	ASN
1	D	363	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 21 ligands modelled in this entry, 15 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	403	-	3,3,3	0.40	0	2,2,2	0.07	0
5	PGE	C	402	-	9,9,9	0.38	0	8,8,8	0.40	0
3	EDO	C	403	-	3,3,3	0.40	0	2,2,2	0.24	0
3	EDO	D	401	-	3,3,3	0.09	0	2,2,2	0.19	0
3	EDO	A	402	-	3,3,3	0.13	0	2,2,2	0.28	0
3	EDO	C	404	-	3,3,3	0.01	0	2,2,2	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	403	-	-	1/1/1/1	-
5	PGE	C	402	-	-	2/7/7/7	-
3	EDO	C	403	-	-	1/1/1/1	-
3	EDO	D	401	-	-	1/1/1/1	-
3	EDO	A	402	-	-	1/1/1/1	-
3	EDO	C	404	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	EDO	O1-C1-C2-O2
3	D	401	EDO	O1-C1-C2-O2
5	C	402	PGE	O2-C3-C4-O3
3	A	402	EDO	O1-C1-C2-O2
3	C	403	EDO	O1-C1-C2-O2
5	C	402	PGE	C3-C4-O3-C5

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	402	PGE	2	0
3	D	401	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	194/204 (95%)	0.23	8 (4%) 41 45	16, 23, 45, 78	2 (1%)
1	B	186/204 (91%)	0.39	4 (2%) 62 66	17, 26, 41, 53	2 (1%)
1	C	193/204 (94%)	0.35	5 (2%) 57 61	16, 27, 41, 49	3 (1%)
1	D	191/204 (93%)	0.59	12 (6%) 26 28	18, 29, 50, 64	2 (1%)
All	All	764/816 (93%)	0.39	29 (3%) 44 48	16, 27, 44, 78	9 (1%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	199	VAL	4.6
1	D	294	ILE	3.2
1	D	292	ASN	3.0
1	D	296	ASN	2.9
1	A	374	GLN	2.8
1	D	293	LYS	2.7
1	D	302	LEU	2.7
1	D	375	MET	2.7
1	D	221	PRO	2.7
1	A	388	ILE	2.7
1	D	291	PRO	2.6
1	C	254	ASN	2.6
1	A	284	PRO	2.6
1	D	288	PHE	2.5
1	D	290	VAL	2.5
1	C	392	THR	2.5
1	B	331	THR	2.4
1	C	198	ASN	2.4
1	A	196	ARG	2.4
1	B	315	ILE	2.4
1	A	285	ASN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	373	ASN	2.3
1	A	372	LYS	2.3
1	D	198	ASN	2.2
1	B	375	MET	2.2
1	A	287	THR	2.1
1	C	257	LYS	2.1
1	C	284	PRO	2.0
1	B	269	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	C	404	4/4	0.77	0.14	45,46,46,46	0
3	EDO	A	402	4/4	0.79	0.19	35,38,39,45	0
3	EDO	D	401	4/4	0.83	0.12	40,41,42,46	0
3	EDO	A	403	4/4	0.86	0.16	27,33,34,43	0
3	EDO	C	403	4/4	0.86	0.16	26,33,38,42	0
5	PGE	C	402	10/10	0.86	0.14	28,34,37,39	0
4	K	D	403	1/1	0.95	0.08	36,36,36,36	0
4	K	D	402	1/1	0.96	0.06	35,35,35,35	0
4	K	C	401	1/1	0.97	0.06	30,30,30,30	0
4	K	A	406	1/1	0.97	0.11	27,27,27,27	0
4	K	A	407	1/1	0.97	0.07	37,37,37,37	0
4	K	A	409	1/1	0.97	0.06	30,30,30,30	0
4	K	C	407	1/1	0.98	0.04	23,23,23,23	0
4	K	B	402	1/1	0.98	0.05	30,30,30,30	0
4	K	A	404	1/1	0.98	0.06	22,22,22,22	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	K	C	405	1/1	0.98	0.06	23,23,23,23	0
4	K	A	405	1/1	0.99	0.06	18,18,18,18	0
4	K	A	408	1/1	0.99	0.04	25,25,25,25	0
2	BR	A	401	1/1	0.99	0.03	24,24,24,24	0
4	K	C	406	1/1	0.99	0.03	24,24,24,24	0
4	K	B	401	1/1	1.00	0.04	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.