



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:26 PM UTC

PDB ID : 4X8A / pdb_00004x8a
Title : NavMS pore and C-terminal domain grown from protein purified in LiCl
Authors : Naylor, C.E.; Bagneris, C.; Wallace, B.A.
Deposited on : 2014-12-10
Resolution : 3.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3131 (3.04-3.00)
Clashscore	190562	3444 (3.04-3.00)
Ramachandran outliers	187476	3319 (3.04-3.00)
Sidechain outliers	187428	3322 (3.04-3.00)
RSRZ outliers	180081	3130 (3.04-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	9% 54% 7% 38%
1	B	149	11% 52% 10% 36%
1	C	149	9% 52% 9% 39%
1	D	149	11% 52% 9% 39%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3034 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ion transport protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	92	Total	C	N	O	S	0	0	0
			717	488	106	118	5			
1	B	95	Total	C	N	O	S	0	0	0
			735	497	109	124	5			
1	C	91	Total	C	N	O	S	0	0	0
			715	485	105	120	5			
1	D	91	Total	C	N	O	S	0	0	0
			712	482	105	120	5			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	126	GLY	-	expression tag	UNP A0L5S6
A	127	SER	-	expression tag	UNP A0L5S6
A	128	HIS	-	expression tag	UNP A0L5S6
A	129	MET	-	expression tag	UNP A0L5S6
B	126	GLY	-	expression tag	UNP A0L5S6
B	127	SER	-	expression tag	UNP A0L5S6
B	128	HIS	-	expression tag	UNP A0L5S6
B	129	MET	-	expression tag	UNP A0L5S6
C	126	GLY	-	expression tag	UNP A0L5S6
C	127	SER	-	expression tag	UNP A0L5S6
C	128	HIS	-	expression tag	UNP A0L5S6
C	129	MET	-	expression tag	UNP A0L5S6
D	126	GLY	-	expression tag	UNP A0L5S6
D	127	SER	-	expression tag	UNP A0L5S6
D	128	HIS	-	expression tag	UNP A0L5S6
D	129	MET	-	expression tag	UNP A0L5S6

- Molecule 2 is HEGA-10 (CCD ID: 2CV) (formula: C₁₈H₃₇NO₇).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 20	C 15	N 1	O 4	0	0
2	B	1	Total 26	C 18	N 1	O 7	0	0
2	B	1	Total 12	C 9	N 1	O 2	0	0
2	C	1	Total 19	C 14	N 1	O 4	0	0
2	D	1	Total 17	C 13	N 1	O 3	0	0

- Molecule 3 is NONAETHYLENE GLYCOL (CCD ID: 2PE) (formula: $C_{18}H_{38}O_{10}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	8	5		
3	B	1	Total	C	O	0	0
			13	8	5		
3	D	1	Total	C	O	0	0
			13	8	5		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Na	0	0
			3	3		
4	C	2	Total	Na	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total	O	0	0
			5	5		
5	B	5	Total	O	0	0
			5	5		
5	C	4	Total	O	0	0
			4	4		
5	D	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Ion transport protein




• Molecule 1: Ion transport protein



• Molecule 1: Ion transport protein



• Molecule 1: Ion transport protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	80.09Å 332.35Å 80.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.56 – 3.02 45.56 – 3.02	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.56-3.02) 99.9 (45.56-3.02)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.03 (at 3.01Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.217 , 0.244 0.223 , 0.244	Depositor DCC
R_{free} test set	1118 reflections (2.50%)	wwPDB-VP
Wilson B-factor (Å ²)	40.8	Xtriage
Anisotropy	1.208	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3034	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2CV, 2PE, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	0/739	1.50	1/1013 (0.1%)
1	B	0.90	0/757	1.51	3/1038 (0.3%)
1	C	0.89	0/737	1.47	2/1010 (0.2%)
1	D	0.90	0/734	1.48	2/1006 (0.2%)
All	All	0.90	0/2967	1.49	8/4067 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	200	ILE	N-CA-CB	7.20	116.07	110.45
1	D	199	PHE	N-CA-C	6.08	117.91	111.28
1	C	200	ILE	N-CA-CB	6.03	115.15	110.45
1	C	199	PHE	N-CA-C	5.96	117.78	111.28
1	B	200	ILE	N-CA-CB	5.66	114.31	110.52
1	A	200	ILE	N-CA-CB	5.30	114.07	110.52
1	B	161	PHE	CA-C-N	5.14	125.69	119.98
1	B	161	PHE	C-N-CA	5.14	125.69	119.98

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	717	0	726	5	0
1	B	735	0	738	6	0
1	C	715	0	724	4	0
1	D	712	0	715	6	0
2	A	20	0	29	1	0
2	B	38	0	50	0	0
2	C	19	0	24	0	0
2	D	17	0	22	0	0
3	A	13	0	16	0	0
3	B	13	0	16	0	0
3	D	13	0	16	0	0
4	A	3	0	0	0	0
4	C	2	0	0	0	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
5	C	4	0	0	0	0
5	D	3	0	0	0	0
All	All	3034	0	3076	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:157:PHE:CZ	1:B:188:VAL:HA	2.44	0.53
1:D:157:PHE:CZ	1:D:188:VAL:HA	2.42	0.52
1:C:181:SER:HB3	1:D:173:GLN:HG3	1.93	0.50
1:B:200:ILE:HG22	1:B:204:MET:HE2	1.94	0.50
1:C:184:ILE:C	1:C:187:PRO:HD2	2.38	0.48
1:D:184:ILE:C	1:D:187:PRO:HD2	2.38	0.48
1:A:157:PHE:CZ	1:A:188:VAL:HA	2.49	0.48
2:A:301:2CV:H351	2:A:301:2CV:H271	1.59	0.46
1:D:189:MET:HA	1:D:192:HIS:O	2.16	0.46
1:B:142:PHE:HZ	1:B:172:PHE:HA	1.81	0.45
1:C:157:PHE:CZ	1:C:188:VAL:HA	2.52	0.45
1:C:189:MET:HA	1:C:192:HIS:O	2.17	0.45
1:B:184:ILE:C	1:B:187:PRO:HD2	2.42	0.44
1:B:197:VAL:O	1:B:201:PRO:HG2	2.16	0.44
1:A:200:ILE:HG22	1:A:204:MET:HE2	2.00	0.44
1:A:142:PHE:HZ	1:A:172:PHE:HA	1.85	0.42
1:A:179:SER:HB3	1:A:183:GLY:HA3	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179:SER:HB3	1:B:183:GLY:HA3	2.01	0.41
1:D:189:MET:HG2	1:D:195:ALA:HB3	2.02	0.41
1:D:196:TRP:HB2	1:D:200:ILE:HD12	2.03	0.40
1:A:175:MET:SD	1:A:210:VAL:HG21	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	90/149 (60%)	85 (94%)	5 (6%)	0	100	100
1	B	93/149 (62%)	87 (94%)	5 (5%)	1 (1%)	11	41
1	C	89/149 (60%)	86 (97%)	3 (3%)	0	100	100
1	D	89/149 (60%)	86 (97%)	3 (3%)	0	100	100
All	All	361/596 (61%)	344 (95%)	16 (4%)	1 (0%)	36	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	131	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/129 (60%)	74 (96%)	3 (4%)	28	61
1	B	78/129 (60%)	74 (95%)	4 (5%)	21	54
1	C	78/129 (60%)	72 (92%)	6 (8%)	12	38
1	D	77/129 (60%)	74 (96%)	3 (4%)	28	61
All	All	310/516 (60%)	294 (95%)	16 (5%)	21	53

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	142	PHE
1	A	144	ILE
1	A	211	LEU
1	B	142	PHE
1	B	163	ASP
1	B	190	ASN
1	B	211	LEU
1	C	142	PHE
1	C	156	THR
1	C	159	GLU
1	C	165	SER
1	C	184	ILE
1	C	211	LEU
1	D	142	PHE
1	D	156	THR
1	D	165	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	151	ASN
1	A	194	ASN
1	A	212	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 5 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	2CV	B	302	-	11,11,25	0.38	0	12,12,30	0.66	0
2	2CV	C	301	-	18,18,25	0.22	0	20,20,30	1.00	2 (10%)
2	2CV	D	301	-	16,16,25	0.27	0	17,17,30	0.81	1 (5%)
3	2PE	D	302	-	12,12,27	0.23	0	11,11,26	0.18	0
3	2PE	A	302	-	12,12,27	0.23	0	11,11,26	0.18	0
2	2CV	B	301	-	25,25,25	0.23	0	29,30,30	0.81	1 (3%)
2	2CV	A	301	-	19,19,25	0.19	0	21,21,30	0.59	1 (4%)
3	2PE	B	303	-	12,12,27	0.16	0	11,11,26	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2CV	B	302	-	-	1/12/12/34	-
2	2CV	C	301	-	-	3/21/21/34	-
2	2CV	D	301	-	-	6/18/18/34	-
3	2PE	D	302	-	-	4/10/10/25	-
3	2PE	A	302	-	-	5/10/10/25	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2CV	B	301	-	-	8/34/34/34	-
2	2CV	A	301	-	-	3/22/22/34	-
3	2PE	B	303	-	-	5/10/10/25	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	301	2CV	C37-C36-N33	3.52	116.94	113.10
2	D	301	2CV	C37-C36-N33	2.59	118.71	112.66
2	C	301	2CV	C35-N33-C36	-2.15	113.70	116.14
2	A	301	2CV	C37-C36-N33	2.15	115.45	113.10
2	B	301	2CV	C36-C37-C40	2.11	115.39	109.66

There are no chirality outliers.

All (35) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	301	2CV	N33-C36-C37-C40
2	B	301	2CV	N33-C36-C37-O47
2	D	301	2CV	N33-C36-C37-O47
2	D	301	2CV	N33-C35-C60-O63
2	B	301	2CV	O51-C41-C42-C43
2	B	301	2CV	C40-C41-C42-C43
2	C	301	2CV	N33-C35-C60-O63
3	B	303	2PE	O10-C11-C12-O13
2	B	302	2CV	C21-C24-C27-C30
2	B	301	2CV	O51-C41-C42-O53
2	B	301	2CV	C40-C41-C42-O53
2	A	301	2CV	N33-C35-C60-O63
2	C	301	2CV	O34-C30-N33-C36
2	D	301	2CV	O34-C30-N33-C36
3	A	302	2PE	C8-C9-O10-C11
3	B	303	2PE	C5-C6-O7-C8
3	B	303	2PE	C9-C8-O7-C6
2	A	301	2CV	C18-C21-C24-C27
2	D	301	2CV	C37-C36-N33-C30
3	A	302	2PE	C11-C12-O13-C14
3	B	303	2PE	C15-C14-O13-C12
2	A	301	2CV	C21-C24-C27-C30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	303	2PE	C12-C11-O10-C9
3	D	302	2PE	C8-C9-O10-C11
2	B	301	2CV	N33-C35-C60-O63
3	A	302	2PE	C15-C14-O13-C12
2	D	301	2CV	C37-C36-N33-C35
3	A	302	2PE	O10-C11-C12-O13
2	C	301	2CV	C21-C24-C27-C30
3	A	302	2PE	C12-C11-O10-C9
2	B	301	2CV	O47-C37-C40-C41
3	D	302	2PE	O7-C8-C9-O10
2	D	301	2CV	C21-C24-C27-C30
3	D	302	2PE	C11-C12-O13-C14
3	D	302	2PE	C5-C6-O7-C8


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	2CV	1	0




The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand 2CV B 302



Ligand 2CV C 301







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	92/149 (61%)	0.81	14 (15%) 5 3	26, 38, 104, 138	0
1	B	95/149 (63%)	0.92	16 (16%) 4 2	26, 37, 121, 167	0
1	C	91/149 (61%)	0.99	14 (15%) 5 3	28, 37, 104, 145	0
1	D	91/149 (61%)	0.99	16 (17%) 4 2	26, 37, 105, 146	0
All	All	369/596 (61%)	0.93	60 (16%) 4 3	26, 37, 110, 167	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	222	MET	6.6
1	A	220	ASP	6.4
1	C	214	PHE	6.0
1	D	214	PHE	5.5
1	B	222	MET	4.7
1	D	130	GLY	4.4
1	B	133	SER	4.2
1	C	130	GLY	4.2
1	B	131	VAL	4.1
1	A	131	VAL	4.1
1	B	224	ILE	3.9
1	C	215	ILE	3.8
1	B	223	ALA	3.8
1	D	215	ILE	3.7
1	B	215	ILE	3.6
1	B	220	ASP	3.5
1	A	133	SER	3.5
1	D	211	LEU	3.4
1	C	218	ILE	3.4
1	B	218	ILE	3.3
1	C	211	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	219	VAL	3.2
1	B	219	VAL	3.2
1	B	132	GLY	3.2
1	D	219	VAL	3.2
1	C	217	ILE	3.2
1	C	220	ASP	3.1
1	D	218	ILE	3.1
1	D	132	GLY	3.1
1	D	131	VAL	3.1
1	D	159	GLU	3.0
1	A	215	ILE	2.9
1	A	218	ILE	2.9
1	B	134	VAL	2.9
1	B	130	GLY	2.7
1	D	213	LEU	2.7
1	C	132	GLY	2.7
1	C	159	GLU	2.6
1	A	221	ALA	2.6
1	C	131	VAL	2.6
1	D	220	ASP	2.5
1	A	204	MET	2.5
1	D	212	ASN	2.5
1	A	211	LEU	2.4
1	D	210	VAL	2.4
1	C	213	LEU	2.4
1	B	159	GLU	2.4
1	B	204	MET	2.3
1	C	207	THR	2.3
1	B	211	LEU	2.3
1	A	134	VAL	2.3
1	A	219	VAL	2.3
1	C	210	VAL	2.3
1	D	217	ILE	2.2
1	B	221	ALA	2.1
1	A	188	VAL	2.1
1	A	217	ILE	2.1
1	D	207	THR	2.0
1	D	137	LEU	2.0
1	A	214	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	2PE	D	302	13/28	0.75	0.31	70,72,81,82	0
2	2CV	B	302	12/26	0.79	0.28	70,79,80,80	0
3	2PE	B	303	13/28	0.86	0.20	56,61,63,64	0
2	2CV	B	301	26/26	0.87	0.17	37,53,78,79	0
2	2CV	A	301	20/26	0.87	0.19	38,49,62,64	0
3	2PE	A	302	13/28	0.88	0.19	60,66,72,72	0
2	2CV	D	301	17/26	0.89	0.17	39,49,54,55	0
2	2CV	C	301	19/26	0.90	0.16	47,58,66,69	0
4	NA	A	303	1/1	0.93	0.54	52,52,52,52	1
4	NA	A	305	1/1	0.94	0.24	30,30,30,30	1
4	NA	C	303	1/1	0.95	0.39	26,26,26,26	1
4	NA	A	304	1/1	0.97	0.39	37,37,37,37	1
4	NA	C	302	1/1	0.99	0.33	38,38,38,38	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 2PE D 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 2CV B 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 2PE B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 2CV B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 2CV A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 2PE A 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 2CV D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 2CV C 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.