



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 02:56 PM UTC

PDB ID : 5X8F / pdb_00005x8f
Title : Ternary complex structure of a double mutant I454RA456K of o-Succinylbenzoate CoA Synthetase (MenE) from Bacillus Subtilis bound with AMP and its product analogue OSB-NCoA at 1.76 angstrom
Authors : Chen, Y.; Guo, Z.
Deposited on : 2017-03-02
Resolution : 1.76 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	485	3% 91% 9%
1	B	485	4% 94% 6%
1	C	485	2% 94% 6%
1	D	485	4% 91% 9%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 17221 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 2-succinylbenzoate--CoA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	484	Total	C	N	O	S	0	7	1
			3702	2363	615	701	23			
1	B	485	Total	C	N	O	S	0	4	0
			3678	2351	612	692	23			
1	C	485	Total	C	N	O	S	0	3	1
			3707	2365	626	693	23			
1	D	484	Total	C	N	O	S	0	6	0
			3692	2362	617	690	23			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	454	ARG	ILE	engineered mutation	UNP P23971
A	456	LYS	ALA	engineered mutation	UNP P23971
B	454	ARG	ILE	engineered mutation	UNP P23971
B	456	LYS	ALA	engineered mutation	UNP P23971
C	454	ARG	ILE	engineered mutation	UNP P23971
C	456	LYS	ALA	engineered mutation	UNP P23971
D	454	ARG	ILE	engineered mutation	UNP P23971
D	456	LYS	ALA	engineered mutation	UNP P23971

- Molecule 2 is o-succinylbenzoyl-N-coenzyme A (CCD ID: S0N) (formula: $C_{32}H_{45}N_8O_{20}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			63	32	8	20	3		
2	B	1	Total	C	N	O	P	0	0
			63	32	8	20	3		
2	C	1	Total	C	N	O	P	0	0
			63	32	8	20	3		
2	D	1	Total	C	N	O	P	0	0
			63	32	8	20	3		

- Molecule 3 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	8	Total	Mg	0	0
			8	8		
4	B	8	Total	Mg	0	0
			8	8		
4	C	4	Total	Mg	0	0
			4	4		
4	D	7	Total	Mg	0	0
			7	7		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total	Na	0	0
			5	5		
5	B	9	Total	Na	0	0
			9	9		
5	C	8	Total	Na	0	0
			8	8		
5	D	11	Total	Na	0	0
			11	11		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		
6	A	1	Total	C	O	0	0
			7	4	3		
6	B	1	Total	C	O	0	0
			7	4	3		
6	B	1	Total	C	O	0	0
			7	4	3		
6	C	1	Total	C	O	0	0
			7	4	3		
6	C	1	Total	C	O	0	0
			7	4	3		
6	D	1	Total	C	O	0	0
			7	4	3		



- Molecule 7 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		
7	B	1	Total	Ca	0	0
			1	1		
7	C	1	Total	Ca	0	0
			1	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	509	Total 509	O 509	0	0
8	B	496	Total 496	O 496	0	0
8	C	528	Total 528	O 528	0	0
8	D	499	Total 499	O 499	0	0

- Molecule 1: 2-succinylbenzoate--CoA ligase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	71.87Å 96.36Å 98.07Å 80.44° 77.96° 81.11°	Depositor
Resolution (Å)	29.96 – 1.76 29.96 – 1.76	Depositor EDS
% Data completeness (in resolution range)	92.9 (29.96-1.76) 92.9 (29.96-1.76)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 1.76Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.163 , 0.194 0.163 , 0.193	Depositor DCC
R_{free} test set	1995 reflections (0.86%)	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 66.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.049 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	17221	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CA, NA, MG, S0N, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/3794	0.53	3/5146 (0.1%)
1	B	0.31	0/3762	0.52	0/5106
1	C	0.29	0/3785	0.49	0/5131
1	D	0.30	0/3785	0.51	0/5134
All	All	0.31	0/15126	0.51	3/20517 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	482	LYS	N-CA-C	5.55	118.27	109.50
1	A	292	CYS	CA-C-N	5.05	131.19	121.54
1	A	292	CYS	C-N-CA	5.05	131.19	121.54

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	480	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3702	0	3608	31	0
1	B	3678	0	3588	20	0
1	C	3707	0	3637	20	0
1	D	3692	0	3628	29	0
2	A	63	0	40	2	0
2	B	63	0	40	2	0
2	C	63	0	40	0	0
2	D	63	0	40	0	0
3	A	23	0	12	4	0
3	C	23	0	12	2	0
4	A	8	0	0	0	0
4	B	8	0	0	0	0
4	C	4	0	0	0	0
4	D	7	0	0	0	0
5	A	5	0	0	0	0
5	B	9	0	0	0	0
5	C	8	0	0	0	0
5	D	11	0	0	0	0
6	A	14	0	20	0	0
6	B	14	0	20	4	0
6	C	14	0	20	1	0
6	D	7	0	10	1	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
8	A	509	0	0	9	1
8	B	496	0	0	4	0
8	C	528	0	0	5	0
8	D	499	0	0	11	1
All	All	17221	0	14715	103	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 103 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:445:CYS:SG	8:C:974:HOH:O	2.39	0.81
1:D:460:VAL:HG23	1:D:485:LEU:HD11	1.61	0.80
1:A:312:LYS:NZ	8:A:602:HOH:O	2.16	0.78
1:B:312:LYS:NZ	8:B:601:HOH:O	2.18	0.76
1:D:418:GLU:OE2	8:D:601:HOH:O	2.05	0.72

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:622:HOH:O	8:D:683:HOH:O[1_556]	2.03	0.17

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	489/485 (101%)	478 (98%)	10 (2%)	1 (0%)	43	27
1	B	487/485 (100%)	477 (98%)	9 (2%)	1 (0%)	43	27
1	C	486/485 (100%)	478 (98%)	7 (1%)	1 (0%)	43	27
1	D	489/485 (101%)	479 (98%)	9 (2%)	1 (0%)	43	27
All	All	1951/1940 (101%)	1912 (98%)	35 (2%)	4 (0%)	43	27

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	293	SER
1	B	293	SER
1	C	293	SER
1	D	293	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/414 (93%)	385 (100%)	0	100	100
1	B	383/414 (92%)	383 (100%)	0	100	100
1	C	387/414 (94%)	387 (100%)	0	100	100
1	D	386/414 (93%)	386 (100%)	0	100	100
All	All	1541/1656 (93%)	1541 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	359	GLN
1	D	95	GLN
1	D	393	ASN
1	D	226	HIS
1	B	284	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 76 ligands modelled in this entry, 63 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PEG	A	517	-	6,6,6	0.54	0	5,5,5	0.23	0
2	S0N	C	501	4	64,66,66	2.59	16 (25%)	92,97,97	1.59	14 (15%)
6	PEG	A	516	-	6,6,6	0.55	0	5,5,5	0.30	0
6	PEG	B	519	-	6,6,6	0.52	0	5,5,5	0.24	0
6	PEG	C	516	-	6,6,6	0.55	0	5,5,5	0.39	0
3	AMP	C	502	-	25,25,25	1.40	3 (12%)	37,38,38	1.89	12 (32%)
6	PEG	D	520	5	6,6,6	0.57	0	5,5,5	0.30	0
3	AMP	A	502	-	25,25,25	1.45	3 (12%)	37,38,38	1.91	10 (27%)
6	PEG	B	520	-	6,6,6	0.54	0	5,5,5	0.21	0
6	PEG	C	515	-	6,6,6	0.55	0	5,5,5	0.17	0
2	S0N	A	501	4	64,66,66	2.54	15 (23%)	92,97,97	1.72	17 (18%)
2	S0N	B	501	4	64,66,66	2.57	15 (23%)	92,97,97	1.68	18 (19%)
2	S0N	D	501	4	64,66,66	2.54	15 (23%)	92,97,97	1.73	16 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	A	517	-	-	1/4/4/4	-
2	S0N	C	501	4	-	11/64/80/80	0/4/4/4
6	PEG	A	516	-	-	2/4/4/4	-
6	PEG	B	519	-	-	2/4/4/4	-
6	PEG	C	516	-	-	2/4/4/4	-
3	AMP	C	502	-	-	5/10/26/26	0/3/3/3
6	PEG	D	520	5	-	2/4/4/4	-
3	AMP	A	502	-	-	2/10/26/26	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	B	520	-	-	2/4/4/4	-
6	PEG	C	515	-	-	0/4/4/4	-
2	S0N	A	501	4	-	11/64/80/80	0/4/4/4
2	S0N	B	501	4	-	11/64/80/80	0/4/4/4
2	S0N	D	501	4	-	14/64/80/80	0/4/4/4

The worst 5 of 67 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	S0N	O16-C17	9.22	1.63	1.42
2	D	501	S0N	O16-C17	9.10	1.63	1.42
2	B	501	S0N	O16-C17	9.02	1.62	1.42
2	A	501	S0N	O16-C17	8.92	1.62	1.42
2	C	501	S0N	C28-C17	-8.16	1.27	1.53

The worst 5 of 87 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	S0N	C41-C42-C43	-5.78	102.77	112.39
2	D	501	S0N	N26-C25-N24	-5.75	119.88	128.58
3	A	502	AMP	C5-C4-N3	-5.71	118.86	126.72
2	A	501	S0N	C21-C27-N26	-5.63	118.97	126.72
2	D	501	S0N	C41-C42-C43	-5.61	103.04	112.39

There are no chirality outliers.

5 of 65 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	S0N	C04-O05-P06-O08
2	A	501	S0N	C14-O13-P10-O09
2	A	501	S0N	C14-O13-P10-O11
2	A	501	S0N	C14-O13-P10-O12
2	B	501	S0N	C14-O13-P10-O09

There are no ring outliers.

8 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	519	PEG	2	0
3	C	502	AMP	2	0



Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	520	PEG	1	0
3	A	502	AMP	4	0
6	B	520	PEG	2	0
6	C	515	PEG	1	0
2	A	501	S0N	2	0
2	B	501	S0N	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







Ligand S0N A 501



Ligand S0N B 501





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	484/485 (99%)	-0.05	13 (2%) 56 62	10, 24, 50, 85	7 (1%)
1	B	485/485 (100%)	-0.03	20 (4%) 41 47	11, 23, 51, 69	4 (0%)
1	C	485/485 (100%)	-0.13	8 (1%) 70 77	9, 25, 45, 64	3 (0%)
1	D	484/485 (99%)	0.02	17 (3%) 47 53	10, 26, 54, 87	6 (1%)
All	All	1938/1940 (99%)	-0.05	58 (2%) 52 59	9, 25, 51, 87	20 (1%)

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	LEU	4.7
1	A	480	ALA	4.6
1	A	327	VAL	4.5
1	B	2	LEU	4.5
1	A	468	ALA	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	AMP	A	502	23/23	0.69	0.23	25,37,58,62	23
5	NA	A	514	1/1	0.70	0.22	48,48,48,48	0
3	AMP	C	502	23/23	0.71	0.20	24,34,54,66	23
5	NA	B	515	1/1	0.77	0.25	52,52,52,52	0
5	NA	C	513	1/1	0.77	0.19	73,73,73,73	0
5	NA	D	512	1/1	0.78	0.20	50,50,50,50	0
5	NA	D	516	1/1	0.78	0.23	69,69,69,69	0
5	NA	B	514	1/1	0.79	0.18	70,70,70,70	0
5	NA	D	509	1/1	0.79	0.16	69,69,69,69	0
4	MG	B	507	1/1	0.81	0.24	71,71,71,71	0
5	NA	C	512	1/1	0.81	0.30	62,62,62,62	0
5	NA	C	509	1/1	0.82	0.16	53,53,53,53	0
5	NA	A	515	1/1	0.82	0.16	60,60,60,60	0
6	PEG	B	520	7/7	0.82	0.14	31,40,65,68	0
5	NA	C	508	1/1	0.83	0.20	76,76,76,76	0
5	NA	B	513	1/1	0.83	0.22	58,58,58,58	0
5	NA	D	519	1/1	0.84	0.13	60,60,60,60	0
5	NA	D	510	1/1	0.86	0.19	61,61,61,61	0
5	NA	D	518	1/1	0.86	0.17	61,61,61,61	0
7	CA	B	521	1/1	0.86	0.12	64,64,64,64	0
4	MG	B	509	1/1	0.87	0.17	42,42,42,42	0
5	NA	C	510	1/1	0.87	0.13	50,50,50,50	0
5	NA	D	511	1/1	0.87	0.15	62,62,62,62	0
5	NA	C	511	1/1	0.88	0.18	59,59,59,59	0
4	MG	B	506	1/1	0.88	0.14	59,59,59,59	0
6	PEG	C	516	7/7	0.89	0.12	36,43,49,57	0
6	PEG	C	515	7/7	0.89	0.13	38,53,66,67	0
5	NA	B	511	1/1	0.90	0.15	64,64,64,64	0
6	PEG	B	519	7/7	0.90	0.12	30,37,44,47	0
4	MG	B	508	1/1	0.91	0.13	57,57,57,57	0
5	NA	B	517	1/1	0.91	0.10	40,40,40,40	0
5	NA	A	511	1/1	0.92	0.10	52,52,52,52	0
5	NA	A	513	1/1	0.92	0.12	46,46,46,46	0
5	NA	D	514	1/1	0.92	0.13	49,49,49,49	0
4	MG	A	507	1/1	0.92	0.12	46,46,46,46	0
5	NA	D	517	1/1	0.92	0.21	67,67,67,67	0
5	NA	C	507	1/1	0.92	0.16	54,54,54,54	0
6	PEG	A	516	7/7	0.93	0.10	40,43,47,50	0
2	S0N	D	501	63/63	0.94	0.09	10,21,46,51	0
4	MG	A	504	1/1	0.94	0.18	39,39,39,39	0
2	S0N	C	501	63/63	0.94	0.09	13,22,51,55	0
5	NA	B	518	1/1	0.94	0.09	52,52,52,52	0
4	MG	A	510	1/1	0.94	0.12	50,50,50,50	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MG	C	504	1/1	0.94	0.14	65,65,65,65	0
5	NA	B	512	1/1	0.94	0.09	38,38,38,38	0
5	NA	D	513	1/1	0.94	0.13	47,47,47,47	0
4	MG	C	506	1/1	0.94	0.13	37,37,37,37	0
6	PEG	D	520	7/7	0.94	0.10	27,37,40,47	0
5	NA	D	515	1/1	0.94	0.09	48,48,48,48	0
5	NA	B	516	1/1	0.95	0.13	39,39,39,39	0
4	MG	A	509	1/1	0.95	0.09	40,40,40,40	0
5	NA	C	514	1/1	0.95	0.11	48,48,48,48	0
4	MG	A	505	1/1	0.95	0.08	43,43,43,43	0
4	MG	D	504	1/1	0.95	0.06	36,36,36,36	0
6	PEG	A	517	7/7	0.95	0.09	33,39,48,54	0
5	NA	A	512	1/1	0.96	0.07	46,46,46,46	0
4	MG	B	502	1/1	0.96	0.08	27,27,27,27	0
4	MG	C	503	1/1	0.96	0.06	22,22,22,22	0
2	S0N	B	501	63/63	0.96	0.07	14,21,43,48	0
5	NA	B	510	1/1	0.96	0.09	33,33,33,33	0
4	MG	D	508	1/1	0.97	0.09	33,33,33,33	0
2	S0N	A	501	63/63	0.97	0.06	13,20,32,36	0
4	MG	A	506	1/1	0.97	0.07	37,37,37,37	0
4	MG	B	503	1/1	0.97	0.10	32,32,32,32	0
4	MG	B	505	1/1	0.97	0.09	37,37,37,37	0
4	MG	D	506	1/1	0.97	0.10	38,38,38,38	0
7	CA	C	517	1/1	0.97	0.19	62,62,62,62	0
4	MG	D	507	1/1	0.98	0.05	37,37,37,37	0
4	MG	D	502	1/1	0.98	0.04	18,18,18,18	0
4	MG	A	503	1/1	0.98	0.04	22,22,22,22	0
4	MG	A	508	1/1	0.98	0.10	36,36,36,36	0
4	MG	D	503	1/1	0.99	0.06	30,30,30,30	0
4	MG	B	504	1/1	0.99	0.05	23,23,23,23	0
7	CA	A	518	1/1	0.99	0.19	34,34,34,34	0
4	MG	D	505	1/1	0.99	0.09	23,23,23,23	0
4	MG	C	505	1/1	0.99	0.05	22,22,22,22	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around AMP A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around AMP C 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around S0N D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around S0N C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around S0N B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around S0N A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.