



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:45 AM UTC

PDB ID : 2X9P / pdb_00002x9p
Title : X-ray structure of the substrate-free cytochrome P450 PimD - a polyene macrolide antibiotic pimaricin epoxidase
Authors : Kells, P.M.; Ouellet, H.; Santos-Aberturas, J.; Aparicio, J.F.; Podust, L.M.
Deposited on : 2010-03-23
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	404	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PIMD PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	385	2955	1877	518	550	10	0	3	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	expression tag	UNP Q9EW92
A	-5	ARG	-	expression tag	UNP Q9EW92
A	-4	GLY	-	expression tag	UNP Q9EW92
A	-3	SER	-	expression tag	UNP Q9EW92
A	-2	HIS	-	expression tag	UNP Q9EW92
A	-1	HIS	-	expression tag	UNP Q9EW92
A	0	HIS	-	expression tag	UNP Q9EW92
A	1	HIS	-	expression tag	UNP Q9EW92
A	2	HIS	-	expression tag	UNP Q9EW92
A	3	HIS	-	expression tag	UNP Q9EW92
A	4	GLY	-	expression tag	UNP Q9EW92

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	177	Total 177	O 177	0	0

- Molecule 1: PIMD PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	118.60Å 139.38Å 70.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	90.17 – 2.10 90.17 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.6 (90.17-2.10) 99.6 (90.17-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.76 (at 2.11Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.186 , 0.239 0.183 , 0.232	Depositor DCC
R_{free} test set	1725 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	31.1	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3190	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.36	9/3021 (0.3%)	1.19	13/4119 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	313	LEU	N-CA	7.13	1.52	1.46
1	A	256	GLN	CG-CD	-7.10	1.34	1.52
1	A	64	GLY	N-CA	6.02	1.51	1.44
1	A	124	ALA	CA-CB	5.92	1.62	1.53
1	A	122	GLU	N-CA	5.30	1.52	1.46
1	A	256	GLN	CA-CB	-5.23	1.44	1.53
1	A	146	THR	CB-CG2	-5.16	1.35	1.52
1	A	27	ALA	CA-CB	5.15	1.61	1.53
1	A	358	THR	CB-CG2	-5.08	1.35	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	256	GLN	CA-CB-CG	-8.42	97.25	114.10
1	A	146	THR	CA-CB-CG2	7.24	122.81	110.50
1	A	374	LEU	CA-C-N	-5.96	114.12	120.03
1	A	374	LEU	C-N-CA	-5.96	114.12	120.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	THR	OG1-CB-CG2	5.46	120.22	109.30
1	A	118	GLN	CA-C-N	-5.46	113.28	119.28
1	A	118	GLN	C-N-CA	-5.46	113.28	119.28
1	A	286	PRO	CA-C-N	-5.46	114.34	119.85
1	A	286	PRO	C-N-CA	-5.46	114.34	119.85
1	A	294	MET	CB-CA-C	-5.27	100.44	109.65
1	A	358	THR	N-CA-CB	5.22	117.89	110.16
1	A	146	THR	N-CA-CB	-5.19	102.48	110.16
1	A	89	ALA	N-CA-C	5.01	116.76	110.24

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	146	THR	CB

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	5	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2955	0	2895	29	0
2	A	43	0	30	6	0
3	A	15	0	0	0	0
4	A	177	0	0	1	0
All	All	3190	0	2925	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:THR:HG21	1:A:244:ASP:OD1	1.77	0.85
1:A:239:VAL:HA	2:A:1398:HEM:HBB1	1.73	0.68
1:A:354:LEU:O	1:A:358:THR:HG23	1.95	0.65
1:A:48:VAL:HG21	1:A:57:LEU:HD12	1.77	0.64
1:A:142:HIS:HA	1:A:146:THR:HG22	1.79	0.63
1:A:108:LEU:HD13	1:A:210:ILE:HG23	1.82	0.61
1:A:239:VAL:HA	2:A:1398:HEM:CBB	2.34	0.57
1:A:118:GLN:HA	1:A:354:LEU:HD21	1.88	0.56
1:A:175:ILE:HD11	1:A:191:LEU:HD22	1.88	0.54
1:A:256:GLN:NE2	1:A:329:ALA:H	2.06	0.54
1:A:66:THR:HG22	1:A:87:SER:HB3	1.91	0.53
1:A:257:ARG:HD2	1:A:364:PHE:CD1	2.44	0.53
1:A:51:HIS:HA	1:A:314:PRO:HB3	1.91	0.51
1:A:348:GLY:HA3	2:A:1398:HEM:C3C	2.46	0.51
1:A:354:LEU:O	1:A:358:THR:CG2	2.59	0.51
1:A:54:LEU:HD13	1:A:314:PRO:HG2	1.93	0.50
1:A:63:ILE:HG22	1:A:288:ARG:HG2	1.93	0.49
1:A:239:VAL:HG22	2:A:1398:HEM:CBB	2.43	0.49
1:A:54:LEU:C	1:A:54:LEU:HD23	2.38	0.49
1:A:284:VAL:HG12	1:A:285:LEU:O	2.13	0.49
2:A:1398:HEM:O2D	4:A:2175:HOH:O	2.20	0.47
1:A:54:LEU:HD23	1:A:54:LEU:O	2.13	0.47
1:A:256:GLN:HE22	1:A:329:ALA:H	1.63	0.46
1:A:367:LEU:HD22	1:A:397:TRP:CD1	2.51	0.44
1:A:239:VAL:CG2	2:A:1398:HEM:HBB2	2.49	0.43
1:A:57:LEU:HD23	1:A:57:LEU:HA	1.94	0.42
1:A:115:LEU:HA	1:A:115:LEU:HD12	1.78	0.42
1:A:48:VAL:HG21	1:A:57:LEU:CD1	2.48	0.41
1:A:38:ARG:HH21	1:A:38:ARG:HD3	1.75	0.41
1:A:191:LEU:HD23	1:A:229:MET:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/404 (95%)	375 (98%)	8 (2%)	1 (0%)	36	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	179	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/328 (92%)	281 (94%)	19 (6%)	16	14

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ARG
1	A	93	SER
1	A	115	LEU
1	A	141	LEU
1	A	146	THR
1	A	157	VAL
1	A	168	LEU
1	A	175	ILE
1	A	179	ASP
1	A	182	GLU
1	A	279	ARG
1	A	288	ARG
1	A	295	GLU
1	A	302	ARG
1	A	358	THR
1	A	371	ARG
1	A	379	LEU
1	A	381	LEU
1	A	394	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	GLN
1	A	188	GLN
1	A	256	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	1399	-	4,4,4	0.42	0	6,6,6	0.65	0
3	SO4	A	1400	-	4,4,4	0.35	0	6,6,6	0.56	0
3	SO4	A	1401	-	4,4,4	0.37	0	6,6,6	0.44	0
2	HEM	A	1398	1	50,50,50	1.74	11 (22%)	67,82,82	1.64	13 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	1398	1	-	2/14/54/54	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1398	HEM	C3D-C2D	6.32	1.50	1.36
2	A	1398	HEM	FE-NA	3.46	2.06	1.95
2	A	1398	HEM	CAB-C3B	2.96	1.55	1.47
2	A	1398	HEM	CMC-C2C	2.73	1.56	1.50
2	A	1398	HEM	CAC-C3C	2.70	1.54	1.47
2	A	1398	HEM	C3B-C2B	-2.63	1.31	1.37
2	A	1398	HEM	C4C-NC	-2.57	1.34	1.39
2	A	1398	HEM	FE-NC	2.41	2.03	1.95
2	A	1398	HEM	FE-NB	2.40	2.02	1.94
2	A	1398	HEM	C3C-C2C	-2.28	1.32	1.37
2	A	1398	HEM	CAD-C3D	2.13	1.56	1.51

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1398	HEM	CBA-CAA-C2A	-5.83	96.40	112.53
2	A	1398	HEM	CAD-C3D-C4D	4.33	132.25	124.70
2	A	1398	HEM	CAB-C3B-C2B	-3.26	117.83	128.43
2	A	1398	HEM	C4D-ND-C1D	3.16	108.94	105.21
2	A	1398	HEM	CHD-C1D-ND	2.61	127.24	124.42
2	A	1398	HEM	O2A-CGA-CBA	2.55	122.05	114.00
2	A	1398	HEM	CMB-C2B-C1B	2.45	128.86	125.03
2	A	1398	HEM	CMA-C3A-C2A	2.26	130.42	125.62
2	A	1398	HEM	CAB-C3B-C4B	2.21	134.15	124.39
2	A	1398	HEM	C4D-C3D-C2D	-2.17	103.73	106.89
2	A	1398	HEM	CHB-C1B-NB	2.17	127.05	124.37
2	A	1398	HEM	CAD-C3D-C2D	-2.08	123.96	127.87
2	A	1398	HEM	O2D-CGD-CBD	2.01	120.35	114.00

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1398	HEM	C3D-CAD-CBD-CGD
2	A	1398	HEM	CAD-CBD-CGD-O2D

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1398	HEM	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	385/404 (95%)	-0.34	9 (2%) 61 64	11, 29, 55, 68	3 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	SER	4.0
1	A	82	LEU	3.4
1	A	178	LEU	3.3
1	A	381	LEU	3.0
1	A	184	ALA	2.6
1	A	81	PHE	2.3
1	A	6	HIS	2.2
1	A	376	VAL	2.2
1	A	180	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	1399	5/5	0.87	0.14	62,64,65,67	0
3	SO4	A	1400	5/5	0.93	0.10	63,63,64,64	0
2	HEM	A	1398	43/43	0.99	0.05	11,18,35,42	0
3	SO4	A	1401	5/5	0.99	0.04	24,24,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.