



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:47 AM UTC

PDB ID : 7X9G / pdb_00007x9g
Title : Crystal structure of human EDA and EDAR
Authors : Yu, K.; Wan, F.; Huang, C.; Wu, J.; Lei, M.
Deposited on : 2022-03-15
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	4% 77% 11% 9%
1	B	159	14% 66% 23% 10%
2	C	123	26% 43% 18% 35%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2853 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ectodysplasins-A, secreted form.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	144	Total	C	N	O	S	0	0	0
			1124	719	190	209	6			
1	B	143	Total	C	N	O	S	0	0	0
			1119	716	189	208	6			


- Molecule 2 is a protein called Tumor necrosis factor receptor superfamily member EDAR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	80	Total	C	N	O	S	0	0	0
			610	377	101	120	12			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ectodysplasins-A, secreted form



- Molecule 1: Ectodysplasins-A, secreted form



- Molecule 2: Tumor necrosis factor receptor superfamily member EDAR



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	87.17Å 87.17Å 156.35Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.71 – 2.80 34.71 – 2.80	Depositor EDS
% Data completeness (in resolution range)	90.9 (34.71-2.80) 90.9 (34.71-2.80)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.248 , 0.282 0.248 , 0.277	Depositor DCC
R_{free} test set	407 reflections (3.74%)	wwPDB-VP
Wilson B-factor (Å ²)	41.7	Xtriage
Anisotropy	0.187	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 49.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.058 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	2853	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/1148	1.21	10/1556 (0.6%)
1	B	0.65	0/1143	1.04	0/1549
2	C	0.74	0/626	1.21	3/844 (0.4%)
All	All	0.69	0/2917	1.15	13/3949 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	261	GLN	N-CA-C	-16.29	81.86	108.41
1	A	262	VAL	N-CA-CB	-11.52	94.88	110.54
1	A	380	PHE	CA-CB-CG	8.06	121.86	113.80
2	C	132	GLY	N-CA-C	7.57	121.65	112.49
1	A	262	VAL	N-CA-C	6.15	116.90	110.62
2	C	82	GLY	N-CA-C	6.04	119.92	111.10
2	C	97	PHE	CA-CB-CG	5.86	119.66	113.80
1	A	344	ASN	CA-CB-CG	5.84	118.44	112.60
1	A	299	GLY	N-CA-C	5.41	118.64	110.75
1	A	314	PHE	CA-CB-CG	5.35	119.15	113.80
1	A	261	GLN	CA-C-N	5.28	127.69	120.46
1	A	261	GLN	C-N-CA	5.28	127.69	120.46
1	A	316	ASP	CA-CB-CG	5.15	117.75	112.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	98	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1124	0	1118	10	0
1	B	1119	0	1113	31	0
2	C	610	0	549	32	0
All	All	2853	0	2780	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:81:LYS:HG2	2:C:108:GLU:HA	1.37	1.02
1:B:325:ASP:CG	1:B:358:GLN:HE22	1.73	0.96
1:B:292:GLU:OE1	1:B:359:LYS:HD3	1.65	0.96
1:B:341:THR:CG2	2:C:94:GLU:HG3	2.03	0.88
1:B:341:THR:HG21	2:C:94:GLU:O	1.79	0.82
1:B:292:GLU:CD	1:B:359:LYS:HD3	2.05	0.81
2:C:98:ARG:HA	2:C:116:CYS:HB3	1.62	0.79
1:B:341:THR:HG23	2:C:94:GLU:HG3	1.66	0.77
1:B:325:ASP:CG	1:B:358:GLN:NE2	2.43	0.75
2:C:52:PRO:HB2	2:C:107:MET:HE3	1.72	0.71
1:A:344:ASN:HD22	1:A:344:ASN:N	1.90	0.68
1:B:325:ASP:CB	1:B:358:GLN:HE22	2.07	0.67
1:B:332:CYS:HB2	1:B:348:THR:HB	1.79	0.64
1:A:276:ARG:HD2	2:C:73:PRO:HD2	1.82	0.60
2:C:54:GLU:HB3	2:C:71:CYS:SG	2.43	0.59
2:C:107:MET:C	2:C:109:ASN:N	2.57	0.58
1:B:252:HIS:ND1	1:B:304:TYR:OH	2.29	0.57
1:B:341:THR:HG23	2:C:94:GLU:CG	2.33	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:52:PRO:HB2	2:C:107:MET:HB2	1.86	0.56
2:C:75:PRO:HB2	2:C:78:LYS:HG3	1.88	0.54
1:B:369:ILE:HG22	1:B:371:ILE:HG13	1.89	0.54
2:C:107:MET:O	2:C:107:MET:HG3	2.07	0.54
1:B:325:ASP:HB2	1:B:358:GLN:HE22	1.72	0.53
1:B:262:VAL:HG13	1:B:266:LEU:HD12	1.92	0.52
2:C:52:PRO:HB3	2:C:108:GLU:HG3	1.91	0.52
2:C:98:ARG:NH2	2:C:132:GLY:O	2.43	0.52
1:B:325:ASP:CB	1:B:358:GLN:NE2	2.70	0.52
1:B:341:THR:HG21	2:C:94:GLU:HG3	1.88	0.52
1:A:272:ASN:HA	1:A:291:GLY:HA3	1.93	0.51
2:C:98:ARG:CG	2:C:116:CYS:HB3	2.40	0.51
2:C:107:MET:O	2:C:108:GLU:C	2.53	0.51
2:C:98:ARG:HG2	2:C:116:CYS:HB3	1.93	0.51
1:B:325:ASP:HB2	1:B:358:GLN:NE2	2.26	0.51
1:A:344:ASN:N	1:A:344:ASN:ND2	2.59	0.50
2:C:102:LEU:HB2	2:C:112:GLU:HB3	1.93	0.50
1:B:311:TYR:HB3	1:B:370:SER:HB2	1.93	0.49
1:A:272:ASN:HA	1:A:291:GLY:CA	2.42	0.49
1:B:332:CYS:SG	1:B:346:CYS:HB3	2.53	0.49
2:C:81:LYS:CG	2:C:108:GLU:HA	2.26	0.49
1:B:311:TYR:CE2	2:C:94:GLU:HG2	2.50	0.47
2:C:98:ARG:CA	2:C:116:CYS:HB3	2.38	0.47
2:C:107:MET:O	2:C:109:ASN:N	2.48	0.47
1:A:332:CYS:SG	1:A:346:CYS:HB3	2.55	0.46
1:A:374:SER:HB3	1:A:377:THR:OG1	2.16	0.46
1:A:253:LEU:HB3	1:A:274:TRP:HB3	1.98	0.46
1:B:263:LYS:HD2	1:B:367:ALA:HB1	1.96	0.46
2:C:98:ARG:HA	2:C:116:CYS:CB	2.38	0.45
2:C:90:HIS:CE1	2:C:104:PRO:HB2	2.52	0.45
1:B:312:ILE:HD11	1:B:318:ALA:HB2	1.99	0.45
2:C:77:GLU:C	2:C:78:LYS:HG2	2.41	0.45
1:A:276:ARG:NH1	2:C:73:PRO:O	2.50	0.45
2:C:81:LYS:HG2	2:C:108:GLU:CA	2.26	0.45
1:B:374:SER:HB3	1:B:377:THR:OG1	2.17	0.44
1:B:250:VAL:HG12	1:B:383:ILE:HG23	2.00	0.44
1:A:312:ILE:HG23	1:A:338:THR:HA	1.99	0.43
1:B:258:SER:O	1:B:373:MET:HB2	2.19	0.42
2:C:52:PRO:CB	2:C:107:MET:HE3	2.47	0.42
1:B:292:GLU:OE1	1:B:359:LYS:CD	2.51	0.42
1:B:300:THR:HG23	1:B:388:ALA:HA	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:319:SER:OG	1:B:331:GLN:NE2	2.45	0.42
2:C:107:MET:C	2:C:109:ASN:H	2.28	0.41
1:B:353:LEU:HD21	1:B:389:PRO:HD2	2.02	0.41
2:C:131:TYR:H	2:C:135:CYS:HB3	1.86	0.41
1:B:277:ILE:HG13	1:B:278:THR:N	2.35	0.40
1:B:310:TYR:HA	1:B:342:ASN:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	142/159 (89%)	131 (92%)	10 (7%)	1 (1%)	18	47
1	B	141/159 (89%)	135 (96%)	6 (4%)	0	100	100
2	C	74/123 (60%)	64 (86%)	10 (14%)	0	100	100
All	All	357/441 (81%)	330 (92%)	26 (7%)	1 (0%)	36	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	314	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/133 (92%)	116 (94%)	7 (6%)	18	49
1	B	123/133 (92%)	119 (97%)	4 (3%)	33	69
2	C	68/104 (65%)	61 (90%)	7 (10%)	7	23
All	All	314/370 (85%)	296 (94%)	18 (6%)	18	49

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	276	ARG
1	A	279	MET
1	A	282	LYS
1	A	289	ARG
1	A	314	PHE
1	A	316	ASP
1	A	344	ASN
1	B	315	THR
1	B	316	ASP
1	B	342	ASN
1	B	348	THR
2	C	47	CYS
2	C	84	TYR
2	C	89	ARG
2	C	97	PHE
2	C	116	CYS
2	C	135	CYS
2	C	136	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	256	GLN
1	A	261	GLN
1	A	272	ASN
1	A	342	ASN
1	A	344	ASN
1	A	366	HIS
1	B	261	GLN
1	B	280	ASN
1	B	358	GLN
1	B	372	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	144/159 (90%)	0.36	7 (4%) 35 27	12, 36, 64, 81	0
1	B	143/159 (89%)	1.00	22 (15%) 5 4	20, 52, 96, 112	0
2	C	80/123 (65%)	1.63	32 (40%) 1 0	39, 67, 113, 122	0
All	All	367/441 (83%)	0.88	61 (16%) 4 3	12, 51, 97, 122	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	136	TYR	5.3
2	C	130	ILE	4.7
1	B	273	ASP	4.0
2	C	134	VAL	3.7
1	B	389	PRO	3.7
2	C	127	PRO	3.6
2	C	59	SER	3.6
2	C	133	MET	3.5
1	B	385	LEU	3.5
2	C	104	PRO	3.4
1	B	247	GLN	3.2
2	C	94	GLU	3.2
2	C	106	ASP	3.0
2	C	47	CYS	3.0
2	C	132	GLY	2.9
2	C	58	LEU	2.9
2	C	102	LEU	2.8
2	C	128	ARG	2.7
1	B	284	PHE	2.7
2	C	105	GLY	2.7
2	C	108	GLU	2.7
1	B	354	LEU	2.7
1	B	314	PHE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	113	CYS	2.6
1	B	358	GLN	2.6
2	C	101	VAL	2.5
2	C	137	SER	2.5
2	C	97	PHE	2.5
1	B	281	PRO	2.5
1	B	278	THR	2.5
1	B	264	ASN	2.5
2	C	100	THR	2.5
2	C	92	ASP	2.4
2	C	110	ASP	2.4
1	B	289	ARG	2.4
2	C	99	ALA	2.4
1	B	368	ASP	2.4
1	A	273	ASP	2.4
2	C	118	PRO	2.4
2	C	81	LYS	2.3
1	B	353	LEU	2.3
2	C	131	TYR	2.3
1	B	312	ILE	2.3
1	A	256	GLN	2.3
1	A	289	ARG	2.3
1	B	388	ALA	2.2
1	B	315	THR	2.2
1	A	277	ILE	2.2
1	B	301	TYR	2.2
2	C	71	CYS	2.2
2	C	93	CYS	2.2
2	C	111	ALA	2.2
1	A	261	GLN	2.2
1	B	283	VAL	2.1
1	B	357	ARG	2.1
1	A	247	GLN	2.1
2	C	85	GLN	2.1
1	B	257	GLY	2.1
1	A	258	SER	2.0
2	C	117	LEU	2.0
1	B	336	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.