



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 10:52 PM UTC

PDB ID : 2XAJ / pdb_00002xaj
Title : Crystal structure of LSD1-CoREST in complex with (-)-trans-2- phenylcyclopropyl-1-amine
Authors : Binda, C.; Valente, S.; Romanenghi, M.; Pilotto, S.; Cirilli, R.; Karytinis, A.; Ciossani, G.; Botrugno, O.A.; Forneris, F.; Tardugno, M.; Edmondson, D.E.; Minucci, S.; Mattevi, A.; Mai, A.
Deposited on : 2010-03-31
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtrriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	852	% 45% 28% 5% 22%
2	B	482	2% 11% 13% . 72%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6356 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called LYSINE-SPECIFIC HISTONE DEMETHYLASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	666	Total	C	N	O	S	0	0	0
			5217	3324	906	967	20			

- Molecule 2 is a protein called REST COREPRESSOR 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	133	Total	C	N	O	S	0	0	0
			1076	676	194	203	3			

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is PHENYLETHYLENECARBOXYLIC ACID (CCD ID: TCA) (formula: $C_9H_8O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	9	1		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LYSINE-SPECIFIC HISTONE DEMETHYLASE 1



Chain B:

Index	Label	Category
1	GLU	GLN
2	THR	ALA
3	GLY	ASN
4	GLY	ASN
5	PRO	ALA
6	GLY	ALA
7	GLY	ALA
8	GLY	ALA
9	GLY	ALA
10	GLY	ALA
11	GLY	ALA
12	GLY	ALA
13	GLY	ALA
14	GLY	ALA
15	GLY	ALA
16	GLY	ALA
17	GLY	ALA
18	GLY	ALA
19	GLY	ALA
20	GLY	ALA
21	GLY	ALA
22	GLY	ALA
23	GLY	ALA
24	GLY	ALA
25	GLY	ALA
26	GLY	ALA
27	GLY	ALA
28	GLY	ALA
29	GLY	ALA
30	GLY	ALA
31	GLY	ALA
32	GLY	ALA
33	GLY	ALA
34	GLY	ALA
35	GLY	ALA
36	GLY	ALA
37	GLY	ALA
38	GLY	ALA
39	GLY	ALA
40	GLY	ALA
41	GLY	ALA
42	GLY	ALA
43	GLY	ALA
44	GLY	ALA
45	GLY	ALA
46	GLY	ALA
47	GLY	ALA
48	GLY	ALA
49	GLY	ALA
50	GLY	ALA
51	GLY	ALA
52	GLY	ALA
53	GLY	ALA
54	GLY	ALA
55	GLY	ALA
56	GLY	ALA
57	GLY	ALA
58	GLY	ALA
59	GLY	ALA
60	GLY	ALA
61	GLY	ALA
62	GLY	ALA
63	GLY	ALA
64	GLY	ALA
65	GLY	ALA
66	GLY	ALA
67	GLY	ALA
68	GLY	ALA
69	GLY	ALA
70	GLY	ALA
71	GLY	ALA
72	GLY	ALA
73	GLY	ALA
74	GLY	ALA
75	GLY	ALA
76	GLY	ALA
77	GLY	ALA
78	GLY	ALA
79	GLY	ALA
80	GLY	ALA
81	GLY	ALA
82	GLY	ALA
83	GLY	ALA
84	GLY	ALA
85	GLY	ALA
86	GLY	ALA
87	GLY	ALA
88	GLY	ALA
89	GLY	ALA
90	GLY	ALA
91	GLY	ALA
92	GLY	ALA
93	GLY	ALA
94	GLY	ALA
95	GLY	ALA
96	GLY	ALA
97	GLY	ALA
98	GLY	ALA
99	GLY	ALA
100	GLY	ALA

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	119.08Å 179.78Å 235.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.45 – 3.30 71.45 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (71.45-3.30) 99.8 (71.45-3.30)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 3.33Å)	Xtriage
Refinement program	REFMAC 5.5.0090	Depositor
R, R_{free}	0.199 , 0.227 0.199 , 0.223	Depositor DCC
R_{free} test set	747 reflections (1.95%)	wwPDB-VP
Wilson B-factor (Å ²)	78.2	Xtriage
Anisotropy	0.097	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 59.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6356	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, TCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.91	0/5331	1.17	23/7232 (0.3%)
2	B	0.77	0/1091	1.05	4/1471 (0.3%)
All	All	0.88	0/6422	1.15	27/8703 (0.3%)

There are no bond length outliers.

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	754	ASP	CA-C-N	8.48	129.31	119.47
1	A	754	ASP	C-N-CA	8.48	129.31	119.47
1	A	172	SER	N-CA-C	7.58	120.89	108.99
1	A	202	GLY	CA-C-N	7.47	129.17	119.84
1	A	202	GLY	C-N-CA	7.47	129.17	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5217	0	5252	183	0
2	B	1076	0	1091	73	0
3	A	53	0	31	4	0
4	A	10	0	7	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6356	0	6381	238	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 238 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:755:PRO:HA	1:A:758:ARG:NH1	1.61	1.14
1:A:286:SER:O	1:A:291:LEU:HD11	1.56	1.03
1:A:794:PRO:HD2	1:A:828:GLN:HE22	1.25	0.98
1:A:453:GLU:OE1	1:A:453:GLU:HA	1.68	0.92
1:A:755:PRO:HA	1:A:758:ARG:HH12	1.30	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	664/852 (78%)	566 (85%)	75 (11%)	23 (4%)	3 18
2	B	131/482 (27%)	102 (78%)	21 (16%)	8 (6%)	1 9
All	All	795/1334 (60%)	668 (84%)	96 (12%)	31 (4%)	2 16

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	425	ASP
1	A	737	SER
1	A	801	GLU
2	B	326	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	341	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	566/699 (81%)	486 (86%)	80 (14%)	3	14
2	B	117/395 (30%)	92 (79%)	25 (21%)	1	5
All	All	683/1094 (62%)	578 (85%)	105 (15%)	2	12

5 of 105 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	632	GLN
1	A	727	CYS
2	B	408	VAL
1	A	645	GLU
1	A	693	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	388	GLN
2	B	423	ASN
1	A	564	HIS
1	A	638	GLN
1	A	680	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TCA	A	901	3	10,10,11	3.33	2 (20%)	11,11,13	3.32	2 (18%)
3	FAD	A	900	4	58,58,58	2.38	13 (22%)	85,89,89	1.99	24 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TCA	A	901	3	-	3/4/4/5	0/1/1/1
3	FAD	A	900	4	-	3/34/50/50	0/5/6/6

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	901	TCA	C2-C3	9.57	1.52	1.34
3	A	900	FAD	C2A-N1A	8.30	1.48	1.33
3	A	900	FAD	C4X-N5	7.25	1.46	1.30
3	A	900	FAD	C2A-N3A	7.05	1.46	1.33
3	A	900	FAD	O4-C4	6.93	1.36	1.23

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	901	TCA	C31-C3-C2	-10.01	109.78	127.11
3	A	900	FAD	N3A-C2A-N1A	-9.59	114.07	128.58
3	A	900	FAD	C9A-N10-C10	-4.45	113.96	120.75
3	A	900	FAD	C5X-C9A-N10	4.02	121.60	117.97
3	A	900	FAD	N9A-C8A-N7A	-3.89	108.42	113.94

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	900	FAD	PA-O3P-P-O5'
4	A	901	TCA	C1-C2-C3-C31
3	A	900	FAD	C2'-C1'-N10-C10
4	A	901	TCA	C2-C3-C31-C36
4	A	901	TCA	C2-C3-C31-C32

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	901	TCA	3	0
3	A	900	FAD	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	666/852 (78%)	-0.17	10 (1%) 72 53	34, 65, 93, 107	0
2	B	133/482 (27%)	0.48	8 (6%) 27 19	66, 94, 110, 118	0
All	All	799/1334 (59%)	-0.06	18 (2%) 61 42	34, 70, 101, 118	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	836	LEU	5.3
1	A	171	PRO	4.5
2	B	375	VAL	4.1
2	B	376	ILE	3.5
1	A	403	ASN	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FAD	A	900	53/53	0.98	0.06	37,43,59,61	0
4	TCA	A	901	10/11	0.98	0.11	59,69,71,71	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.